Iranian Journal of Science & Technology, Transaction A, Vol. 34, No. A1 Printed in the Islamic Republic of Iran, 2010 © Shiraz University

SIMULTANEOUS CONTROL OF THE SOURCE TERMS IN A VIBRATIONAL STRING PROBLEM^{*}

T. YELOGLU¹ AND M. SUBASI^{2**}

¹Mustafa Kemal University, Faculty of Science and Literature, Department of Mathematics, Hatay, Turkey Email: tyeloglu@mku.edu.tr ²Ataturk University, Faculty of Science, Department of Mathematics, 25240, Erzurum, Turkey Email: msubasi@atauni.edu.tr

Abstract – In this paper, simultaneous control of source terms is considered in a vibrational string problem. In the considered problem, the terms to be controlled are the force and the initial velocity functions. We state the generalized (weak) solution about the considered problem. The existence and uniqueness of the solution for optimal control problem is investigated. The Frechet derivative of the functional and the Lipschitz continuity of the gradient are investigated. Minimizing sequence is obtained by the method of the projection of the gradient.

Keywords - Optimal control problem, frechet derivative, projection of the gradient

1. INTRODUCTION

Simultaneous determination of the functions in inverse hyperbolic problems has been investigated by some researchers. Bamberger, Chavent and Lailly investigated the parameter functions $(\rho(z), \mu(z))$ in

$$\rho(z)\frac{\partial^2 y}{\partial t^2} - \frac{\partial}{\partial z} \left(\mu(z)\frac{\partial y}{\partial z}\right) = 0, \ z > 0, \ t > 0$$

from the only boundary measurement of y(z,t)/z = 0 in [1]. V. Isakov dealt with the problem of determining the other two coefficients functions in [2]. Gugat, Leugering and Sklyar investigated the problem of determination of $f_1(t)$, $f_2(t)$ from the condition $y(x,T) = y_0(x)$, $y_t(x,T) = y_1(x)$ in the problem

$$y_{tt}(x,t) = c^{2} y_{xx}(x,t), (x,t) \in [0,L] \times [0,T]$$

$$y(x,0) = 0, y_{t}(x,0) = 0, x \in [0,L]$$

$$y(0,t) = f_{1}(t), y(L,t) = f_{2}(t), t \in [0,T]$$

in [3]. A. Hasanov examined the determination of the pair $w := \{F(x,t), f(t)\}$ from the final state $u(x,T) = \mu(x)$ in the problem;

^{*}Received by the editor May 3, 2010 and in final revised form July 26, 2010

^{**}Corresponding author

T. Yeloglu / M. Subasi

$$u_{tt} = (k(x)u_x)_x + F(x,t), (x,t) \in (0,L) \times (0,T)$$

$$u(x,0) = u_0(x), u_t(x,0) = u_1(x), x \in (0,L)$$

$$-k(0)u_x(0,t) = f(t), k(l)u_x(l,t) = 0, t \in (0,T)$$

in [4].

Considering the inverse source problem by a control method, the reconstruction formula and regularization methods have been obtained in [5] and [6]. In [5], Mordukhovich and Raymond considered the problem of minimizing the integral functional

$$J(y,u) = \int_{\Omega} \varphi(x, y(T)) dx + \int_{Q} g(x,t,y) dx dt + \int_{\Sigma} h(s,t,u) ds dt$$

for the problem

 $y_{tt} - \Delta y = f \qquad \text{in } Q := \Omega \times]0, T[$ $y = u \qquad \text{on } \Sigma := \Gamma \times]0, T[$ $y(0) = y_0, y_t(0) = y_1 \qquad \text{in } \Omega$

M. Yamamoto in [6] considered the determination of f(x) from $\left[\frac{\partial u(f)}{\partial n}\right](x,t)$ in the problem

$$u''(x,t) = \Delta u(x,t) + \sigma(t) f(x), \quad (x \in \Omega, 0 < t < T)$$
$$u(x,0) = u'(x,0) = 0, \quad (x \in \Omega)$$
$$u(x,t) = 0, \quad (x \in \partial\Omega, 0 < t < T).$$

The stress in this paper is the force and initial velocity functions in the vibrational string problem can be controlled simultaneously from the final state of the string. The paper is organized as follows: In section 2, we give the statement of the problem with its physical motivation and state the definition of the generalized (weak) solution. In section 3, we prove the existence and uniqueness of the solution for optimal control problem by Weierstrass theorem and strict convexity of the functional. In section 4, we get the Frechet differentiability of the functional and Lipschitz constant of the gradient. In the last section, we constitute a minimizing sequence using the Lipschitz constant by the method of the projection of the gradient and prove its convergence to the optimal solution.

2. STATEMENT OF THE PROBLEM

In this paper, we consider the problem of simultaneous control of the functions $w := \{f(x,t), h(x)\}$ in the problem

$$\rho(x)u_{tt} = (k(x)u_{x})_{x} + f(x,t), \ (x,t) \in \Omega_{T} = (0,l) \times (0,T] \subset \mathbb{R}^{2}$$
(1)

$$u(x,0) = g(x), \ u_t(x,0) = h(x), \ x \in (0,l)$$
⁽²⁾

$$u(0,t) = 0, \ u(l,t) = 0, \ t \in (0,T].$$
(3)

Using the functional

Iranian Journal of Science & Technology, Trans. A, Volume 34, Number A1

Simultaneous control of the source terms in...

$$J_{\alpha}(w) = \int_{0}^{t} \left[u(x,T;w) - y(x) \right]^{2} dx + \alpha \|w\|_{W}^{2}.$$
 (4)

For $\alpha > 0$. we will deal with the problem

$$J_{\alpha}\left(w_{*}\right) = \inf_{w \in W} J_{\alpha}\left(w\right).$$
⁽⁵⁾

39

The problem (1)-(3) models the vibration of a flexible string made of nonhomogeneous material. Here the functions $\rho(x) > 0$ and k(x) > 0 represent the density and flexural rigidity of the string. f(x,t) is the applied force function to the string. According to the Hooke's law, the function $T(x,t) := k(x)u_x(x,t)$ defines the tension of the flexible string. g(x) and h(x) state the initial position and the initial velocity, respectively. The x = 0 and x = l ends of the string are fixed. Here, the functions satisfy the conditions;

$$f(x,t) \in L_2(\Omega_T), \ g(x) \in W_2^1[0,l], \ h(x) \in L_2[0,l], \ y(x) \in L_2(0,l)$$
(6)

and

$$0 < \rho_0 \le \rho(x) \le \rho_1, \quad 0 < k_0 \le k(x) \le k_1, \quad \stackrel{0}{\forall} x \in [0, l].$$
(7)

The space W_2^1 is a Sobolev space whose functions with their first order generalized derivatives belong to L_2 space.

The admissible control set is defined as follows:

$$W := \left\{ w = \{f, h\} : \|w\|_{W} \le b \right\} \subset L_{2}(\Omega_{T}) \times L_{2}[0, l].$$
(8)

The inner product in the space W is defined as

$$\left(w_{1},w_{2}\right)_{W} := \int_{0}^{T} \int_{0}^{l} f_{1}\left(x,t\right) f_{2}\left(x,t\right) dx dt + \int_{0}^{l} h_{1}\left(x\right) h_{2}\left(x\right) dx, \ \forall \ w_{1},w_{2} \in W,$$
(9)

and the norm of an element is

$$\|w\|_{W} = \left(\int_{0}^{T} \int_{0}^{1} \left[f(x,t)\right]^{2} dx dt + \int_{0}^{1} \left[h(x)\right]^{2} dx\right)^{1/2}.$$
(10)

Definition 2.1. The function $u \in W_2^{0^{-1}}$ with u(x,0) = g(x), which satisfies the following equation for all $\eta \in W_2^{0^{-1}}$, is called the generalized solution of the problem (1)-(3),

$$\iint_{\Omega_T} \left(-\rho(x)u_t \eta_t + k(x)u_x \eta_x \right) dx dt = \int_0^t \rho(x)h(x)\eta(x,0) dx + \iint_{\Omega_T} f(x,t)\eta dx dt .$$
(11)

Here, the space $\bigcup_{0}^{0} W_2^1$ consists of those elements of W_2^1 which vanish for t = 0 and W_2^1 consists of those elements of W_2 which vanish for t = T.

Now, for $w \in W$ the following theorem is valid;

Theorem 2.2. Let the conditions (6), (7) hold. Then for each $w \in W$, there is a generalized solution of the problem (1)-(3), and for this solution the estimate

Winter 2010

T. Yeloglu / M. Subasi

$$\|u\|_{H^{1}(\Omega_{T})}^{2} \leq c_{5}\left(\|g\|_{H^{1}[0,l]}^{2} + \|w\|_{W}^{2}\right)$$
(12)

is valid.

Proof: The proof follows from the Galerkin method [7].

Since the problem is ill-posed for $\alpha = 0$, we use the parameter $\alpha > 0$ as the regularization parameter, which is the strong convexity constant, which guarantees the uniqueness and stability of the regularized solution. This parameter can be found for numerical investigations by regularization method such as Tikhonov regularization ([8]-[9]).

3. THE EXISTENCE AND UNIQUENESS OF THE OPTIMAL SOLUTION

In this section, we prove that the problem (5) has a solution. Here, we show that the problem satisfies the conditions of Weierstrass theorem. The space $L_2(\Omega_T) \times L_2[0,l]$ is a Hilbert space and the set W is a weakly compact set in this space. If we can show that the functional $J_{\alpha}(w)$ is weakly semicontinuous from below on the set W, then according to the generalized Weierstrass theorem[10], we will prove the existence of the minimum. In addition, if the functional $J_{\alpha}(w)$ is strictly convex then the minimum will be unique (Theorem 38.C. in [10]).

Let's consider the functional

$$J_{0}(w) = \int_{0}^{t} \left[u(x,T;w) - y(x) \right]^{2} dx$$

and

$$J_{\alpha}(w) = J_{0}(w) + \alpha \left\|w\right\|_{W}^{2}.$$

If the functional $J_{\alpha}(w)$ is convex and continuous on W then it is weakly semicontinuous from below. Now, let's prove the continuity of $J_{\alpha}(w)$ on W. To do this it is enough to show the continuity of $J_{0}(w)$.

Theorem 3.1. Let $w \in W$ be an arbitrary element and Δw be an increment for w such that $w + \Delta w \in W$. So, the following inequality is valid;

$$\left|\Delta J_{0}\left(w\right)\right| = \left|J_{0}\left(w + \Delta w\right) - J_{0}\left(w\right)\right| \le s_{0}\left(\left\|\Delta w\right\|_{w} + \left\|\Delta w\right\|_{w}^{2}\right).$$

$$(13)$$

Here the number s_0 is independent of Δw .

Proof: For $w + \Delta w = \{f + \Delta f, h + \Delta h\}$, we show by $u_{\Delta}(x,t;w) = u(x,t;w + \Delta w)$ and the increment function $\Delta u(x,t;w) = u_{\Delta}(x,t;w) - u(x,t;w)$ satisfies the problem;

$$\rho \Delta u_{tt} = \left(k \Delta u_x\right)_x + \Delta f \tag{14}$$

$$\Delta u(x,0) = 0 \quad \Delta u_t(x,0) = \Delta h \tag{15}$$

$$\Delta u(0,t) = \Delta u(l,t) = 0.$$
⁽¹⁶⁾

The variation of $J_0(w)$ is

Iranian Journal of Science & Technology, Trans. A, Volume 34, Number A1

Simultaneous control of the source terms in...

$$\Delta J_{0}(w) = J_{0}(w + \Delta w) - J_{0}(w) = \int_{0}^{l} \left\{ \left[u_{\Delta}(x, T; w) - y \right]^{2} - \left[u(x, T; w) - y \right]^{2} \right\} dx$$

$$= \int_{0}^{l} 2 \left[u(x, T; w) - y \right] \Delta u(x, T) dx + \int_{0}^{l} \left[\Delta u(x, T) \right]^{2} dx$$
(17)

With Cauchy-Bunyakovsky inequality, we can write that

$$\left|\Delta J_{0}(w)\right| \leq 2\left\|u(x,T;w) - y\right\|_{L_{2}[0,l]} \left\|\Delta u(x,T)\right\|_{L_{2}[0,l]} + \left\|\Delta u(x,T)\right\|_{L_{2}[0,l]}^{2}$$
(18)

Lemma 3.2. For the function $\Delta u(x,T)$, the following estimate is valid;

$$\left\|\Delta u\left(x,T\right)\right\|_{L_{2}[0,l]}^{2} \leq s_{2}\left(\left\|\Delta h\right\|_{L_{2}[0,l]}^{2} + \left\|\Delta f\right\|_{L_{2}(\Omega_{T})}^{2}\right) = s_{2}\left\|\Delta w\right\|_{W}^{2}, \quad s_{2} = 3s_{1}^{2}$$
(19)

Here, $s_1 = \max\left\{T\sqrt{\frac{r_0}{\rho_0}}, \frac{T\sqrt{T}}{\rho_0\sqrt{3}}\right\}$ and $r_0 = \max\left\{\rho_1, k_1\right\}$.

Proof: Proof can be obtained via the processes in [4]. Since $y(x) \in L_2[0, l]$, applying ε – Cauchy inequality for (18), we get

$$\left|\Delta J_{0}(w)\right| \leq s_{0}\left(\left\|\Delta w\right\|_{w}+\left\|\Delta w\right\|_{w}^{2}\right).$$

Here the number s_0 is independent of Δw . Theorem 3.1 has been proven.

So, the functional $J_{\alpha}(w)$ is continuous on W, then it is weakly semicontinuous from below.

Theorem 3. 2. The functional $J_{\alpha}(w)$ is strongly convex on W.

Proof: We know that the set W is convex [11]. From the linearity of the boundary value problem and uniqueness of the solution, we can write that

$$u(x,t;\beta w_{1}+(1-\beta)w_{2}) = \beta u(x,t;w_{1})+(1-\beta)u(x,t;w_{2})$$

for each $w_1, w_2 \in W$ and $\beta \in [0,1]$. Now, we will show that the following inequality is valid with the constant $\chi > 0$,

$$J_{\alpha}(\beta w_{1} + (1-\beta)w_{2}) \leq \beta J_{\alpha}(w_{1}) + (1-\beta)J_{\alpha}(w_{2}) - \chi\beta(1-\beta)||w_{1} - w_{2}||_{W}^{2}$$

for $\forall w_1, w_2 \in W$ and for $\forall \beta \in [0,1]$. From the convexity of the function $|u - y|^2$ to u, we have

$$J_{\alpha} \left(\beta w_{1} + (1-\beta)w_{2}\right) = \int_{0}^{l} \left[\psi\left(x,T;\beta w_{1} + (1-\beta)w_{2}\right) - y(x)\right]^{2} dx + \alpha \left\|\beta w_{1} + (1-\beta)w_{2}\right\|_{W}^{2}$$

$$\leq \beta \left\{\int_{0}^{l} \left[\psi\left(x,T;w_{1}\right) - y(x)\right]^{2} dx + \alpha \left\|w_{1}\right\|_{W}^{2}\right\}$$

$$+ (1-\beta) \left\{\int_{0}^{l} \left[\psi\left(x,T;w_{2}\right) - y(x)\right]^{2} dx + \alpha \left\|w_{2}\right\|_{W}^{2}\right\}$$

$$- \alpha \beta (1-\beta) \left\|w_{1} - w_{2}\right\|_{W}^{2}.$$

Iranian Journal of Science & Technology, Trans. A, Volume 34, Number A1 www.SID.ir

Winter 2010

T. Yeloglu / M. Subasi

So, the cost functional is strongly convex with $\chi = \alpha$.

We proved that the cost functional is strongly convex (so strictly convex) for strong convexity constant $\alpha > 0$ on the weakly compact set W. So according to the generalized Weierstrass theorem [8], the optimal solution $w_* \in W$ to the problem

$$J_{\alpha}(w_{*}) = \inf_{w \in W} J_{\alpha}(w)$$

exists and is unique.

4. FRECHET DIFFERENTIABILITY OF THE FUNCTIONAL AND LIPSCHITZ CONTINUITY OF GRADIENT

In this section, we use the definition

$$\Delta J_{\alpha}(w) = J_{\alpha}(w + \Delta w) - J_{\alpha}(w) = \left(J_{\alpha}'(w), \Delta w\right)_{w} + O\left(\left\|\Delta w\right\|_{w}^{2}\right)$$

of Frechet differentiability of $J_{\alpha}(w)$,

Theorem 4.1. Let the conditions (6)–(7) hold. Therefore, the functional $J_{\alpha}(w) \in C^{1}(W)$ can be Frechet differentiable. Moreover, this derivation can be defined as

$$J'_{\alpha}(w) = \left\{ \eta(x,t;w) + 2\alpha f, \ p(x)\eta(x,0;w) + 2\alpha h \right\}$$
(20)

via the solution of adjoint boundary value problem;

$$\rho \eta_{tt} = \left(k\eta_x\right)_x \tag{21}$$

$$\eta(x,T) = 0 \tag{22}$$

$$\rho \eta_t (x,T) = -2 \left[u(x,T) - y \right]$$
(23)

$$k(0)\eta(0,t) = 0, \ k(l)\eta(l,t) = 0$$
 (24)

Proof: Let's consider the increment of the functional $J_{\alpha}(w)$;

$$\Delta J_{\alpha}(w) = J_{\alpha}(w + \Delta w) - J_{\alpha}(w) = \int_{0}^{l} \left\{ \left[u_{\Delta}(x,T;w) - y \right]^{2} - \left[u(x,T;w) - y \right]^{2} \right\} dx$$

+ $\left\| w + \Delta w \right\|_{W}^{2} - \left\| w \right\|_{W}^{2}$
= $\int_{0}^{l} 2 \left[u(x,T;w) - y \right] \Delta u(x,T) dx + \int_{0}^{l} \left[\Delta u(x,T) \right]^{2} dx$
+ $\int_{0}^{T} \int_{0}^{l} 2\alpha f \Delta f dx dt + \int_{0}^{l} 2\alpha h \Delta h dx + \alpha \left\| \Delta w \right\|_{W}^{2}.$

Now we must estimate the term $2\int_{0}^{\infty} \left[u(x,T;w)-y\right]\Delta u(x,T)dx$, whose relation with $\Delta J_{0}(w)$ can be seen in (17).

If we multiply the adjoint boundary value problem by Δu and integrate over Ω_r , we find

Iranian Journal of Science & Technology, Trans. A, Volume 34, Number A1

Simultaneous control of the source terms in...

$$\int_{0}^{T} \int_{0}^{T} \rho \eta_{tt} \Delta u dt dx = \int_{0}^{T} \int_{0}^{T} (k \eta_{x})_{x} \Delta u dt dx.$$

By integration by parts, we get

$$\iint \left[\rho \Delta u_{tt} - \left(k \Delta u_x \right)_x \right] \eta dx dt + \int_0^t \rho \eta \left(x, 0 \right) \Delta h dx = \int_0^t 2 \left[u \left(x, T \right) - y \right] \Delta u \left(x, T \right) dx.$$
(25)

After multiplying (14)- (16) by $\eta(x,t)$ and integrating over Ω_T , if we subtract this from (25), we have

$$\int_{0}^{t} \rho \eta(x,0) \Delta h dx + \int_{0}^{T} \int_{0}^{t} \Delta f \eta dx dt = 2 \int_{0}^{t} \left[u(x,T;w) - y \right] \Delta u(x,T) dx$$

Considering (19) here, we write

$$\Delta J_{\alpha}(w) = \int_{0}^{T} \int_{0}^{l} \eta \Delta f dx dt + \int_{0}^{l} \rho \eta(x, 0; w) \Delta h dx + \int_{0}^{T} \int_{0}^{l} 2\alpha f \Delta f dx dt + \int_{0}^{l} 2\alpha h \Delta h dx + o\left(\left\|\Delta w\right\|_{w}^{2}\right)$$

so the Frechet derivative is

$$J'_{\alpha}(w) = \left\{ \eta(x,t;w) + 2\alpha f, \ \rho(x)\eta(x,0;w) + 2\alpha h \right\}$$

Hence, theorem 4.1 has been proven.

Theorem 4.2. Let the conditions of Theorem 4.1 hold. Then the estimate

$$\left\|J_{\alpha}'\left(w+\Delta w\right)-J_{\alpha}'\left(w\right)\right\|_{w} \leq L\left\|\Delta w\right\|_{w}, \quad \forall w,w+\Delta w \in W$$

$$\tag{26}$$

is valid. Here,

$$L = \sqrt{2qs_2 + 8\alpha^2}, \ q := \max\left\{\frac{4l^2T}{\rho_0}, \frac{4\rho_1^2l^2}{\rho_0}\right\}, \ s_2 = 3\left(\max\left\{T\sqrt{\frac{r_0}{\rho_0}}, \frac{T\sqrt{T}}{\rho_0\sqrt{3}}\right\}\right)^2$$
(27)

and

$$\left\|J_{\alpha}'(w+\Delta w)-J_{\alpha}'(w)\right\|_{w}^{2}=\iint_{\Omega_{f}}\left[\Delta\eta(x,t;w)+2\alpha\Delta f\right]^{2}dxdt+\int_{0}^{1}\left[\rho(x)\Delta\eta(x,0;w)+2\alpha\Delta h\right]^{2}dx$$
(28)

Proof: We can rewrite (28) with the inequality $(a+b)^2 \le 2(a^2+b^2)$ such that

$$\begin{aligned} \left\|J_{\alpha}'(w+\Delta w)-J_{\alpha}'(w)\right\|_{w}^{2} &= \iint_{\Omega_{r}}\left[\Delta\eta(x,t;w)+2\alpha\Delta f\right]^{2}dxdt + \int_{0}^{t}\left[\rho(x)\Delta\eta(x,0;w)+2\alpha\Delta h\right]^{2}dx\\ &\leq 2\iint_{\Omega_{r}}\left[\Delta\eta(x,t;w)\right]^{2}dxdt + 2\iint_{\Omega_{r}}\left[2\alpha\Delta f\right]^{2}dxdt\\ &+ 2\int_{0}^{t}\left[\rho(x)\Delta\eta(x,0;w)\right]^{2}dx + 2\int_{0}^{t}\left[2\alpha\Delta h\right]^{2}dx.\end{aligned}$$

Iranian Journal of Science & Technology, Trans. A, Volume 34, Number A1 www.SID.ir

Winter 2010

T. Yeloglu / M. Subasi

If we use the increment $w + \Delta w$ in the solution of adjoint boundary value problem and show by $\eta_{\Delta} = \eta(x,t;w + \Delta w)$, we get the following problem for $\Delta \eta = \eta(x,t;w + \Delta w) - \eta(x,t;w)$;

$$\rho \Delta \eta_{tt} = \left(k \Delta \eta_x\right)_x \tag{29}$$

$$\Delta \eta \left(x, T \right) = 0 \tag{30}$$

$$\rho \Delta \eta_t \left(x, T \right) = -2\Delta u \left(x, T; w \right) \tag{31}$$

$$k(0)\Delta\eta(0,t) = 0 \quad k(l)\Delta\eta(l,t) = 0.$$
(32)

If we multiply the both sides of (29) by $\Delta \eta_t$, integrate over [0, l] and use boundary conditions, we have

$$\int_{0}^{l} \left[\rho \Delta \eta_{tt} - (k \Delta \eta_{x})_{x} \right] \Delta \eta_{t} dx = 0$$

$$\int_{0}^{l} \left[\rho \Delta \eta_{tt} \Delta \eta_{t} + k \Delta \eta_{x} \Delta \eta_{xt} \right] dx = \int_{0}^{l} (k \Delta \eta_{x} \Delta \eta_{t})_{x} dx$$

$$\frac{1}{2} \frac{d}{dt} \left\{ \int_{0}^{l} \left[\rho \left(\Delta \eta_{t} \right)^{2} + k \left(\Delta \eta_{x} \right)^{2} \right] dx \right\} = k (l) \Delta \eta_{x} (l, t) \Delta \eta_{t} (l, t) - k (0) \Delta \eta_{x} (0, t) \Delta \eta_{t} (0, t).$$

Integrating the last equality over [t, T] we obtain,

$$\frac{1}{2}\int_{0}^{l} \left\{ \rho \left[\Delta \eta_{t}(x,T) \right]^{2} + k \left[\Delta \eta_{x}(x,T) \right]^{2} \right\} dx = \frac{1}{2}\int_{0}^{l} \left\{ \rho \left[\Delta \eta_{t}(x,t) \right]^{2} + k \left[\Delta \eta_{x}(x,t) \right]^{2} \right\} dx$$

$$\int_{0}^{l} \left(\rho \Delta \eta_{t}^{2} + k \Delta \eta_{x}^{2} \right) dx = \int_{0}^{l} \frac{4 \left[\Delta u(x,T) \right]^{2}}{\rho(x)} dx$$

$$\int_{0}^{l} \Delta \eta_{x}^{2} dx \leq \frac{4}{\rho_{0}} \int_{0}^{l} \left[\Delta u(x,T) \right]^{2} dx.$$
(33)

For the term $\Delta \eta_x$, we have the following inequality;

$$\frac{2}{l^2} \left\| \Delta \eta \left(x, t \right) \right\|_{L_2[0,l]}^2 \leq \left\| \Delta \eta_x \right\|_{L_2[0,l]}^2, \, \forall t \in [0,T].$$

Substituting this into (33), we write

$$\frac{2}{l^{2}}\int_{0}^{l} \Delta \eta^{2} dx \leq \frac{4}{\rho_{0}}\int_{0}^{l} \left[\Delta u(x,T)\right]^{2} dx$$
$$\left\|\Delta \eta\right\|_{L_{2}[0,l]}^{2} \leq \frac{2l^{2}}{\rho_{0}}\left\|\Delta u(x,T)\right\|_{L_{2}[0,l]}^{2}, \forall t \in [0,T]$$

and integrating over [0,T], we get

$$\int_{0}^{T} \int_{0}^{l} \left[\Delta \eta \left(x, t; w \right) \right]^{2} dx dt \leq \frac{2l^{2}T}{\rho_{0}} \left\| \Delta u \left(x, T \right) \right\|_{L_{2}[0, l]}^{2}$$

Moreover, we can write that

Iranian Journal of Science & Technology, Trans. A, Volume 34, Number A1

Simultaneous control of the source terms in...

$$\int_{0}^{l} \left[\rho(x) \Delta \eta(x,0) \right]^{2} dx \leq \rho_{1}^{2} \int_{0}^{l} \left[\Delta \eta(x,0) \right]^{2} dx \leq \frac{2\rho_{1}^{2}l^{2}}{\rho_{0}} \left\| \Delta u(x,T) \right\|_{L_{2}[0,l]}^{2}$$

Hence

$$2\int_{0}^{T}\int_{0}^{l} \left[\Delta\eta(x,t;w)\right]^{2} dxdt + 2\int_{0}^{l} \left[\rho(x)\Delta\eta(x,0)\right]^{2} dx \leq \frac{4l^{2}T}{\rho_{0}} \left\|\Delta u(x,T)\right\|_{L_{2}[0,l]}^{2} + \frac{4\rho_{1}^{2}l^{2}}{\rho_{0}} \left\|\Delta u(x,T)\right\|_{L_{2}[0,l]}^{2}$$
(34)
and taking as $q := \max\left\{\frac{4l^{2}T}{\rho_{0}}, \frac{4\rho_{1}^{2}l^{2}}{\rho_{0}}\right\}$, we obtain
 $\left\|J_{\alpha}'(w + \Delta w) - J_{\alpha}'(w)\right\|_{w}^{2} \leq q \left\|\Delta u(x,T)\right\|_{L_{2}[0,l]}^{2} + 8\alpha^{2} \left\|\Delta w\right\|_{w}^{2}$

With (19) one can write

$$\left\|J_{\alpha}'\left(w+\Delta w\right)-J_{\alpha}'\left(w\right)\right\|_{w}^{2} \leq 2qs_{2}\left\|\Delta w\right\|_{w}^{2}+8\alpha^{2}\left\|\Delta w\right\|_{w}^{2}$$

and for $L = \sqrt{2qs_2 + 8\alpha^2}$, we have

$$\left\|J_{\alpha}'\left(w+\Delta w\right)-J_{\alpha}'\left(w\right)\right\|_{w}\leq L\left\|\Delta w\right\|_{w}$$

Hence, theorem 4.2 has been proven.

5. MINIMIZING SEQUENCE AND ITS CONVERGENCE

We set the minimizing sequence according to the method of projection of the gradient [10], by

$$w_{k+1} = P_W \left(w_k - \beta_k J'_{\alpha} \left(w_k \right) \right), \quad k = 0, 1, 2, \dots$$
(35)

where $P_W\left(w_k - \beta_k J'_{\alpha}(w_k)\right)$ is the projection of the element $w_k - \beta_k J'_{\alpha}(w_k)$ in the set W.

Theorem 5.1. Let $w_0 \in W$ be the initial point. The sequence $\{w_k\}$ defined by (35) converges to the unique minimum element w_* of the functional $J_{\alpha}(w)$ for $\beta_k = \beta \in (0, 4\alpha L^{-2})$. Here, *L* is the Lipschitz constant for $J'_{\alpha}(w)$. Moreover, for the minimizing sequence the following inequality holds;

$$||w_k - w_*|| \le ||w_0 - w_*||.q^k, k = 0, 1, ...$$

where $q = (1 - 4\alpha\beta + \beta^2 L^2)^{1/2} \in (0, 1).$

Proof: We define the mapping $A: W \to W$ with

$$Aw = P_{W}(w - \beta J'_{\alpha}(w)).$$

It can easily be shown that the mapping P_W is a contraction mapping, namely;

$$\left\|P_{W}\left(u\right)-P_{W}\left(v\right)\right\|\leq\left\|u-v\right\|; \ \forall u,v\in W.$$

We must show that the mapping A also holds this property, while $0 < \beta < 4\alpha L^{-2}$. We know that

$$\langle J'_{\alpha}(u) - J'_{\alpha}(v), u - v \rangle \ge 2\alpha \|u - v\|^2, \quad \forall u, v \in W.$$
 (36)

So, using (36) and Lipschitz continuity of the functional, we write the following:Winter 2010Iranian Journal of Science & Technology, Trans. A, Volume 34, Number A1

www.SID.ir

T. Yeloglu / M. Subasi

$$\begin{aligned} \left\|Au - Av\right\|^{2} &= \left\|P_{W}\left(u - \beta J_{\alpha}'\left(u\right)\right) - P_{W}\left(v - \beta J_{\alpha}'\left(v\right)\right)\right\|^{2} \\ &\leq \left\|\left(u - v\right) - \beta\left(J_{\alpha}'\left(u\right) - J_{\alpha}'\left(v\right)\right)\right\|^{2} \\ &= \left\|u - v\right\|^{2} - 2\beta\left\langle J_{\alpha}'\left(u\right) - J_{\alpha}'\left(v\right), u - v\right\rangle + \beta^{2}\left\|J_{\alpha}'\left(u\right) - J_{\alpha}'\left(v\right)\right\|^{2} \\ &\leq \left(1 + \beta^{2}L^{2} - 4\beta\alpha\right)\left\|u - v\right\|^{2} \end{aligned}$$

Then we have

$$\left\|Au - Av\right\| \le q \left\|u - v\right\|$$

by $q = (1 + \beta^2 L^2 - 4\beta\alpha)^{1/2}$. The condition $\beta < 4\alpha L^{-2}$ gives $q \in (0,1)$.

The statement (35) can be written as $w_{k+1} = Aw_k$. Then by the contraction mapping principle, the sequence $\{w_k\}$ converges to the fixed point $w_* = Aw_*$ of the operator A with the factor q. We know that the minimum element w_* is unique. So we get the following:

$$\|w_{k} - w_{*}\| = \|Aw_{k-1} - Aw_{*}\| \leq q \|w_{k-1} - w_{*}\|$$

$$\leq q^{2} \|w_{k-2} - w_{*}\|$$

$$\vdots$$

$$\leq q^{k} \|w_{0} - w_{*}\|$$

Hence, the theorem 5.1. has been proven.

REFERENCES

- 1. Bamberger, A., Chavent, G. & Lailly, P. (1979). About the stability of the inverse problem in 1-D wave equations-application to the interpretation of seismic profiles. *Applied Mathematics and Optimization*, 5, 1-47.
- 2. Isakov, V. (1991). An inverse hyperbolic problem with many boundary measurements, Comm. *Partial Differential Equations, 16,* 1183-1195.
- Gugat, M., Leugering, G. & Sklyar, G. (2005). L^p Optimal Boundary control for the wave equation, SIAM J. Control Optim., 44(1), 49-74.
- 4. Hasanov, A. (2009). Simultaneous determination of the source terms in a linear hyperbolic problem from the final overdetermination: weak solution approach. *IMA Journal of Applied Mathematics*, 74, 1-19.
- 5. Mordukhovich, B. S. & Raymond, J. P. (2004). Dirichlet boundary control of hyperbolic equations in the presence of state constraints. *Appl. Math. Optim.*, 49, 145-157.
- 6. Yamamoto, M. (1995). Stability, reconstruction formula and regularization for an inverse source hyperbolic problem by a control method. *Inverse Problems*, *11*, 481-496.
- 7. Ladyzhenskaya, O. A. (1985). Boundary Value Problems in Mathematical Physics. New York, Springer-Verlag.
- 8. Subaşı, M. (2002). An Optimal Control Problem Governed by the Potential of A Linear Schrödinger Equation. *Applied Mathematics and Computation*, *131*(1), 95-106.
- 9. Subaşı, M. (2004). A Variational method of optimal control problems for nonlinear Schrödinger Equation, Numerical Methods for Partial Differential Equations, 20(1), 82-89.
- Zeidler, E. (1985). Nonlinear Functional Analysis and Its Applications. Variational Methods and Optimization. New York, Springer-Verlag.
- 11. Ekeland, I. & Temam, R. (1976). *Convex Analysis and Variational Problems*. Stud. Math. Appl., *vol. 1*, New York, North-Holland Publishing Company.

Iranian Journal of Science & Technology, Trans. A, Volume 34, Number A1