Antimicrobial activity of Iranian propolis and its chemical composition

^{*1}Yaghoubi M.J., ¹Ghorbani Gh., ²Soleimanian Zad S., ²Satari R.

¹Department of Animal Sciences, ² Department of Food Sciences and Technology, College of Agriculture, Isfahan University of Technology, Isfahan, Iran. Received 25 March 2006; Revised 4 Jul 2006; Accepted 30 Jul 2006

ABSTRACT

The objective of this study was to investigate the antimicrobial activity of ethanol extract of Iranian propolis on some microorganisms using disc diffusion method. The chemical composition of the propolis was also investigated using thin layer chromatography and spectrophotometric methods. Ethanol extract of propolis showed activity only against Gram-positives and fungi, whereas no activity was observed against Gram-negatives. Thin layer chromatography screening revealed the presence of pinocembrine, caffeic acid, kaempferol, phenethyl caffeate, chrysin, and galangin in Iranian propolis. The total flavonoid and phenolic contents were 7.3% and 36%, respectively, which suggests that the strong antimicrobial activity of Iranian propolis may be due to high levels of phenolic and flavonoid compounds.

Keywords: Iranian propolis, Antimicrobial activity, Phenolic compounds, Flavonoid.

INTRODUCTION

Propolis is a mixture of bees wax and resins collected by the honeybee from plant buds, leaves, and exudates (1). Bees use propolis not only as a building material but also as a means of maintaining low levels of bacterial and fungal in the hive (2).

Propolis has long been used in oriental folk medicine for curing infections (3) and in European ethno-pharmacology as an antiseptic and anti-inflammatory agent for healing wounds and burns (1).

Many pharmaceutical properties including antibacterial (4), antifungal (5), antiviral (6), antiprotozoan (7), anti-inflammatory (8), antioxidant (9), hepatoprotective (10), immunostimulating (11), antitumor (12), and cytostatic (13) activities have been reported for propolis; hence, its wide recognition as a useful substance in medicine (14).

More than 150 components such as polyphenols, phenolic aldehydes, sesquiterpene quinines, coumarins, amino acids, steroids and inorganic components have been identified in propolis samples (15). The properties and chemical composition of propolis vary with geographical origin (5) and the differences in chemical composition are basically due to differences in the bearing plants (16). Although many active components have been identified in propolis (14, 15) and its antimicrobial activity has been demonstrated for different microorganisms (1),

Correspondence: mj_yaghoubi@yahoo.com

there is no report that we know on antimicrobial activity of Iranian propolis. The objective of the present study was to investigate the antimicrobial activity of ethanol extract of Iranian propolis and to analyze its chemical compositions.

MATERIALS AND METHODS

Extraction of propolis

Propolis sample was collected from an experimental apiary located in the Lavark area, Isfahan, central Iran. Hand collected propolis was kept in a dry place and stored at 4 °C until its processing. The sample was cut into small pieces, grounded and extracted with 80% ethanol (1:10 w/v) in a shaker (300 rpm) at room temperature for 48 h. The ethanol extract solution was then filtered through a Whatman # 41 filter paper. Based on the dry weight of the solution, the ethanol extract of propolis (EEP) solution was further adjusted with appropriate amounts of 80% ethanol to obtain solutions containing various amounts of EEP.

Microorganisms

The following microorganisms were used in this study to test antimicrobial activity of propolis. *Staphylococcus aureus* PTCC 1189, *Staphylococcus epidermidis* PTCC1114, *Bacillus subtilis* PTCC 1023, *Bacillus cereus* PTCC 1015, *Bacillus liqueiniformis* PTCC 1331, *Candida albicans* PTCC 5027, *Salmonella enteritidis*

PTCC 1091, *Escherichia coli* PTCC 1038, *Klebsiella pneumoniae* PTCC 5027, and *Proteus vulgaris* PTCC 1182. All microorganisms were provided by Biotechnology Institute, Iranian Research Organization for Sciences and Technology, Tehran.

Preparation of inoculums

All bacteria were cultured for 8 h at 37° C in a liquid medium (brain heart infusion) and used as inoculums. The yeast was sub-cultured in yeast glucose broth at 30° C for 8 h and then used for the test. The turbidity of the suspension was adjusted to the McFarland 0.5 turbidity standard.

Antimicrobial activity

Antimicrobial activity of propolis ethanol extract was investigated by the disc diffusion method (17). The antimicrobial screening was performed using brain heart infusion broth for bacteria and sabouraud dextrose agar for yeasts.

Sterile paper discs (Whatman # 4 paper, 6 mm diameter) were loaded with 5 µl of propolis extract dilutions (67, 16.7, 8.3, 4.1, 2.0, 1.0, 0.2, 0.1, 0.06, and 0.01 mg/ml). Discs were dried for 5 hrs at 37 °C in a sterile incubator and then placed on seeded agar plates. Six discs were applied to each petri dish. The 80% ethanol and commercial disc of ampicilline (1µg- oxoid) were used as negative and positive controls, respectively. Plates were incubated at 37° C and 30° C for 24 h for bacteria and C. albicans, respectively. Inhibitory zone diameters were measured with Scion Image software (Scion Corporation, Frederick, MD). All experiments were performed in triplicates. Values are expressed as means. The data were submitted to analysis of variance using general liner model procedure of statistatical analyzer software (18). The chosen level of significance for all statistical tests was P< 0.01.

Thin layer chromatography analysis

Thin layer chromatography (TLC) analysis of the propolis ethanol extract was performed on silica gel (Alufolien Kieselgel Merck F_{254}) with mobile phase petroleum ether/ethyl acetate 7:3. Spots were visualized by U.V. light (366nm); spraying with 60% sulfuric acid in ethanol and heating at 100 °C. Standard substances were used as described in the literature (2).

Estimation of total flavonoids and phenolics content

Total flavonoid content was determined using a colorimetric method by a previously described method. (19). The Folin-Ciocalteu method was used to determine the quantities of total phenolics (20).

RESULTS AND DISCUSSION

The mean diameters of microbial growth inhibited by different concentrations of EEP are shown in Table 1. Among the bacteria and fungi, B. cereus was the most sensitive to the highest concentration of EEP (67 mg/ml). The sensitivity of the tested microorganisms followed the order: B. cereus > B. liqueniformis > S. epidermidis > S. aureus and C. albicans > B. subtilis. The least active concentration against the tested microorganisms were 4.1 mg/ml for C. albicans and S. epidermidis, 2.0 mg/ml for S. aureus and B. cereus, and 8.3 mg/ml for B. subtilis and B. liqueniformis. The 80% ethanol (negative control) did not show any inhibitory effects on the tested microorganisms. The results also showed that, at a concentration of 67 mg/ml, EEP was more effective than standard ampicilline on S. aureus, epidermidis and B. cereus strains, with S. significant differences (P <0.01), but less active on B. subtilis.

Ethanol extract of propolis showed activity only against Gram-positive bacteria and fungi, whereas, no activity was observed against Gramnegative bacteria. Similar results have been reported in other studies (21, 8, 4, 22), which support our findings that propolis is mainly active against Gram-positives. However it has been reported that EEP is effective on Gram-negative bacteria at higher concentrations (23).

The studies carried out on the antimicrobial activity of propolis show conflicting results (14). The variation in the antimicrobial activity of propolis has been attributed to the differences in its chemical components (14). For example, correlation has been shown between the flavonoid content and antimicrobial activity of propolis against B. Subtilis (24). It has also been reported that the flavonoid content varies considerably in the 38 samples collected from different parts of Croatia with different climates and vegetation (24). The TLC screening showed eleven spots upon exposure to U.V. light (Figure 1). Compared to standard substances, only six out of eleven spots were recognized, showing the presence of flavonoid aglycones such as pinocembrin, chrysin, galangin, and kaempferol, phenolic acids like caffeic acid and esters such as phenethyl caffeate. It is well documented that among the most potent compounds in propolis antimicrobial are flavonone pinocembrin and flavonol galangin (25). Caffeic acid and its esters, volatile fractions with phenols, terpenoids and chrysin have also exhibited notable antimicrobial activity as well (26). No spot was observed in the TLC pattern sample after obtaining a retention factor (R_f) of 0.64, indicating that the pinostrobin compound $(R_f = 0.82)$ either is not present or has

Microorganism ·	EEP ² concentrations										Amnicilling	сE
	1	2	3	4	5	6	7	8	9	10	Ampicinine	5.E .
S. epidermidis	4.8 ^a	2.6 ^c	1.7 ^d	1.0 ^e	-	-	-	-	-	-	3.0 ^b	0.10
S. aureus	4.1 ^a	2.0 ^b	1.5 ^c	1.0 ^d	1.0 ^d	-	-	-	-	-	2.2 ^b	0.13
B. cereus	6.1 ^a	3.8 ^b	3.1 ^c	2.1 ^d	1.0 ^e	-	-	-	-	-	4.1 ^b	0.20
B. subtilis	2.6 ^b	1.3 ^c	0.5 ^d	-	-	-	-	-	-	-	9.6 ^a	0.20
B. liqueniformis	5.1 ^a	2.1 ^b	1.0 ^c	-	-	-	-	-	-	-	-	0.13
S. enteritidis	-	-	-	-	-	-	-	-			-	-
E. coli	-	-	-	-	-	-	-	-	-	-	-	-
P. vulgaris	-	-	-	-	-	-	,	-	-	-	-	-
K. pneumoniae	-	-	-	-	-	-	•	-		-	-	-
C. albicans	4.1 ^a	2.6 ^b	1.7 ^c	1.0 ^d	-	-	-	,	-	-	-	0.22

Table 1. The mean of the diameters¹ (mm) of microbial growth inhibited by different concentrations of ethanol extract of Iranian propolis and Ampicilline.

¹Values expressed are averages of three replicates. Values within each row followed by different letters are significantly different at P<0.01.

²1: 67mg/ml, 2: 16.7 mg/ml, 3: 8.3 μg/ml, 4: 4.1 mg/ml, 5: 2.0 mg/ml, 6: 1.0 mg/ml, 7: 0.2 mg/ml, 8: 0.1 mg/ml, 9: 0.06 mg/ml, 10: 0.01 mg/ml.

a too low concentration to be detected. Furthermore, the sample used was free from benzyl ferulate ($R_f = 0.57$).



Figure 1. Charting Thin layer chromatography (visible at daylight) of ethanol extract of Iranian propolis. Colors are reported as seen at U.V. light (366 nm).

The sample was also analyzed for total flavonoid and total phenolic contents, using spectrophotometric procedure. The total flavonoid and phenolic components were 7.3% and 36%, respectively. The concentrations of these compounds compared to Turkish propolis (2) were considerably higher.

CONCLUSION

The results of this study indicate the antimicrobial activity of ethanolic extract of Iranian propolis. From these results it may be concluded that Gram-positives bacteria are more susceptible to EEP antibacterial activity than Gram-negatives bacteria. The results of TLC analysis confirmed the presence of pinocembrin, caffeic acid, kaempferol, phenethyl caffeate, chrysin, and galangin in Iranian propolis. The strong antimicrobial activity of Iranian propolis may be due to high total phenolic and flavonoid contents. There are numerous questions yet to be answered concerning chemical compositions and antibacterial properties of Iranian propolis and further research is required for clarification.

ACKNOWLEDGEMENT

The authors would like to thank Mr. S. H. Khoshoee for technical assistance. This research was supported by Isfahan University of Technology, Iran.

REFERENCES

- 1. Ghisalberti EL. Propolis: a review. Bee World 1979; 60: 59-84.
- 2. Popova M, Silici S, Kaftanoglu O, Bankova V. Antibacterial activity of Turkish propolis and its qualitative and quantitative chemical composition. Phytomedicine 2005; 12: 221-228.
- 3. Cheng PC, Wong G. Honey bee propolis: prospects in medicine. Bee World 1996; 77: 8-15.
- 4. Lu LC, Chen YW, Chou CC. Antibacterial and DPPH free radical-scavenging activities of the ethanol extract of propolis collected in Taiwan. J Food Drug Analysis 2003; 11: 277-282.
- 5. Kujumgiev A, Tsvetkova I, Serkedjieva Y, Bankova V, Christov R, Popov S. Antibacterial, antifungal and antiviral activity of propolis of different geographic origin. J Ethnopharmacol 1999; 64: 235-240.
- Amoros M, Lurton E, Boustie J, Girre L, Sauvager F, Cormier M. Comparison of the anti-Herpes simplex virus activities of propolis and 3-methyl-butyl-2-enyl caffeate. J Nat Prod 1994; 57: 644-647.
- 7. Scheller S, Szaflarski J, Tustanowski J, Nolewajka E, Stojko A. Biological properities and clinical application of propolis I. Arzeneimittel-Forschung Drug Res 1977; 27: 889-890.
- 8. Dobrowolski JW, Vohora SB, Sharma K, Shah SA, Naqvi SAH, Dandiya PC. Antibacterial, antifungal, antiamoebic, anti inflammatory and antipyretic studies on propolis bee products. J Ethnopharmacol 1991; 35: 77-82.
- 9. Isla MI, Moreno MIN, Sampietro AR, Vattuone MA. Antioxidant activity of Argentina propolis extracts. J Ethnopharmacol 2001; 76: 165-170.
- 10. Gonzales R, Corcho I, Remirez D, Rodriguez S, Ancheta O, Merino N, Gonzales A, Pascual C. Hepatoprotective effects of propolis extract on carbon tetrachloride-induced liver injury in rats. Phytother Res 1995; 9: 114-117.
- Dimov V, Ivanovska N, Bankova V, Nikolov N, Popov S. Immunomodulatory action of propolis: IV. Prophylatic activity against Gram-negative infections and adjuvant effect of water soluble derivative. Vaccine 1992; 10: 817-823.
- 12. Orsolic N, Sver L, Terzic S, Tadic Z, Basic I. Inhibibitory effect of water-soluble derivative of propolis and its polyphenolic compounds on tumor growth and metastasizing ability: a possible mode of antitumor action. Nutr Cancer 2003; 47(2): 156-163.
- Frenkel K, Wei H, Bhimani R, Ye J, Zadunaisky JA, Huang MT, Ferraro T, Conney AH, Grunberger D. Inhibition of tumor promoter-mediated processes in mouse skin and bovine lens by caffeic acid phenethyl ester. Cancer Res 1993; 53: 1255-1261.
- 14. Burdock GA. Review of the biological properties and toxicity of bee propolis (propolis). Food Chem Toxicol 1998; 36: 347-363.
- 15. Marcucci MC. Propolis: chemical composition, biological properties and the therapeutic activity. Apidologie 1995; 26: 83-99.
- 16. Markham KE, Mitchel KA, Wilkins AL, Daldy JA, Lu Y. HPLC and GC-MS identification of the major organic constituents in New Zealand propolis. Phytochemistry 1996; 42: 205-211.
- 17. Bauer AW, Kirby MDK, Sherries JC, Truck M. Antibiotic susceptibilities testing by standard single disc diffusion method. Am J Clin Pathol 1966; 45: 493-496.
- 18. SAS Institute Inc. SAS/STAT User's Guide, release 8.02 edition. SAS Institute Inc., Cary, NC. 1999.
- 19. Eberhardt MV, Lee CY, Liu RH. Antioxidant activity of fresh apples. Nature 2000; 405(6789): 903-904.
- 20. Woisky RG, Salatino A. Analysis of propolis: some parameters and procedures for chemical quality control. J Apicult Res 1998; 37: 99-105.
- 21. Davey RW, Grange JM. Antibacterial properties of propolis (bee glue). J R Soc Med 1990; 83(3): 159-160.
- 22. Nieva Moreno MI, Isla MI, Cudmani NG, Vattuone MA, Sampietro AR. Screening of antibacterial activity of Amaicha del Valle (Tucumán, Argentina) propolis. J Ethnopharmacol 1999; 68: 97-102.
- 23. Sforcin JM, Fernandes JA, Lopes CAM, Bankova V, Funari SRC. Seasonal effect on Brazilian propolis antibacterial activity. J Ethnopharmacol 2000; 73: 243-249.
- 24. Pepeljnjak S, Jalsenjak I, Maysinger D. Flavonoid content in propolis extracts and growth inhibition of Bacillus subtilis. Pharmazie 1985; 40: 122-123.
- 25. Koo H, Rosalen PL, Cury JA, Ambrosano GMB, Murata RM, Yatsuda R, Ikegaki M, Alencar SM, Park YK. Effect of a new variety of Apis mellifera propolis on mutans streptococci. Curr Microbiol 2000; 41: 192-196.
- Kosalec I, Bakmaz M, Pepeljnjak S. Analysis of propolis from the continental and Adriatic regions of Croatia. Acta Pharm 2003; 53: 275-285.