

An Efficient Method for the Nitration of Phenols with NaNO_2 in the Presence of 3-Methyl-1-Sulfonic Acid Imidazolium Chloride

A. Khazaei^{1,*}, M.A. Zolfigol^{1,*}, A.R. Moosavi-Zare¹ and A. Zare²

Abstract. Nitrophenols can be obtained in high yields via nitrosation-oxidation of phenols by 3-methyl-1-sulfonic acid imidazolium chloride {[Msim]Cl} as a new Brønsted acidic ionic liquid and NaNO_2 at room temperature. In situ generation of HNO_2 and a radical-cation mechanism via the nitrous acid catalyzed pathway appear to be applicable to phenol nitration using this reagent system.

Keywords: 3-Methyl-1-sulfonic acid imidazolium chloride {[Msim]Cl}; Brønsted acidic ionic liquid; Nitrosation-oxidation; Nitration; Nitrophenols; Sodium nitrite (NaNO_2).

INTRODUCTION

Ambient temperature ionic liquids, especially those based on 1,3-dialkylimidazolium cations, have gained considerable interest as promising alternative green solvents in organic synthesis [1]. These ionic liquids have several interesting properties, such as excellent chemical and thermal stability, non-volatility, non-coordinating natures, safety, good solvating capability, a wide liquid range, and ease of recycling [1]. Furthermore, their hydrophobicity/hydrophilicity can be tuned by appropriate modification of the cations or anions [1]. Therefore, wide applications in catalytic and non-catalytic reactions have been found for room temperature ionic liquids [2-6]. In addition, the synthesis of task-specific ionic liquids which have a functional group in their framework may expand the applications of ionic liquids in organic chemistry [7,8]. Among ionic liquids, acidic ones have been successfully used in many organic transformations [7-10]. In continuation of our previous studies on the applications of acidic reagents and catalysts in organic synthesis [11,12], more recently, we have synthesized ionic liquid 3-methyl-1-

sulfonic acid imidazolium chloride {[Msim]Cl} as a new Brønsted acidic reagent from the simple reaction of 1-methylimidazole with chlorosulfonic acid at room temperature (Figure 1) [13]. We believe that this new ionic liquid can be applied as a reagent, as well as catalyst, for different organic transformations. Herein, we found that the acidity of this Brønsted acidic reagent was suitable for the in situ generation of HNO_2 , when used in conjunction with NaNO_2 , for the nitration of phenols.

The nitration of organic compounds is a very active and rewarding area of research, being also the subject of a large body of literature [14]. Moreover, nitro-aromatic compounds have been extensively utilized as chemical feedstock for a wide range of useful materials, such as dyes, pharmaceuticals, perfumes and plastics [15]. The nitration of benzene and toluene is one of the most important routes to substituted aromatics in the production of chemical intermediates [16]. Nitration normally requires the use of a corrosive liquid phase nitric acid-sulfuric acid mixture, leading to an excessive waste stream. Furthermore, the nitration reactions are not usually regioselective and some over-nitration or competitive oxidation may occur [17]. The nitration of phenol, taken as a special case, has been achieved by several nitrating agents [18-27]. For example, some nitrating agents including sodium nitrite [18,19], concentrated nitric acid, acid anhydrides or triflates [20-22], peroxy nitrite [23], metal nitrates [24-26] and nitrogen oxides [27] have been utilized for this transformation. Although several

1. Faculty of Chemistry, Bu-Ali Sina University, Hamedan, P.O. Box 6517838683, Iran.

2. Department of Chemistry, Payame Noor University (PNU), Iran.

*. Corresponding authors. E-mail: Khazaei_1326@yahoo.com, mzolfigol@yahoo.com and zolfi@basu.ac.ir

Received 21 December 2009; received in revised form 27 February 2010; accepted 6 April 2010




Figure 1. The synthesis of ionic liquid [Msim]Cl.

methods for the nitration of phenols are known, newer methods continue to attract attention for their experimental simplicity and effectiveness. In this work, we wish to use [Msim]Cl/NaNO₂ as a new and efficient reagent for the nitration of phenolic compounds to give the corresponding nitrated phenols at high yields and short reaction times.

RESULTS AND DISCUSSION

During the course of our studies on the utilization of NO⁺ in functional group transformations [28], we thought that phenol (**1**) could be converted to *p*-nitrosophenol (**2**), selectively, by a 3-methyl-1-sulfonic acid imidazolium chloride (3 eq)/NaNO₂ (1 eq) system (1 equivalent of [Msim]Cl is needed for the oxidation step). Moreover, phenol nitrosation is rapid and yields almost entirely the *para*-isomer which can be readily converted to *p*-nitrophenol via a mild oxidation with HNO₃ [29] etc. (Figure 2, path a). Therefore, we decided to produce *p*-nitrophenol via the nitrosation-oxidation strategy in a one-pot reaction under mild conditions [30]. We selected [Msim]Cl as the acid source (a proton source for the in situ generation of HNO₂ and NO⁺ under mild conditions)/NaNO₂ system for the nitrosation of phenols (Figure 2, path b). In most reported methods for the nitration of phenols, NaNO₃ (1 eq) was needed for the second step; in situ generation of HNO₃ for the oxidation of *p*-nitrosophenol (Figure 2, path a). In contrast to the reported procedures in aqueous media [31], we observed that nitrosation and oxidation were occurred simultaneously using the [Msim]Cl/NaNO₂ system. Thus, in this reaction, mono-nitration was occurred,




Figure 2. The mono-nitration of phenol.

and *p*-nitrophenol (**3**) as well as *o*-nitrophenol (**4**) were obtained at 39% and 50%, respectively, after 25 min (Figure 2, path b).

A good range of 4-substituted phenols (**5**) was also subjected to the nitration reaction in the presence of 3-methyl-1-sulfonic acid imidazolium chloride and NaNO₂ at room temperature (Figure 3). The nitration reactions proceeded under mild conditions at room temperature and afforded the products at high yields and short reaction times (Figure 3 and Table 1). The presented nitration reactions were readily carried out by grinding the nitrating agents and phenols (**1** or **5**) in a mortar. The mono-nitrophenols were obtained by a simple workup.

A competitive reaction was also performed between phenols and anisole. It was observed that exclusive phenol nitration proceeded; anisole remaining intact in the reaction mixtures after 24 hours (Figure 4). Selective mono-nitration of 4,4'-dihydroxydiphenyl (**5 k**) was also achieved by controlling the stoichiometric of the reagents (Table 1, entry 12).

The 3-Methyl-1-sulfonic acid imidazolium chloride/sodium nitrite system is similar to N₂O₄ (N₂O₄ ⇌ NO⁺NO₃⁻) in the case of generation of HNO₂ (eq 1) and NO⁺ that work as a nitrosating agent (Figure 5, eq 2). Thus, on the basis of our observations, the previously reported results concerning the applications of N₂O₄ [32] and metal nitrate dinitrogen tetroxide complexes [M(NO₃)_m.nN₂O₄] [27], and the reported mechanism for the nitration of phenols [33], the following reported nitrous acid catalyzed mechanism may be approved (Figure 5) [19,34].

In another study, the [Msim]Cl/NaNO₂ system was examined for di-nitration of phenols. For this purpose, 6.2 eq of [Msim]Cl and 2 eq of NaNO₂ were reacted with 1 eq of *p*-methoxyphenol. In this reaction,




Figure 3. The mono-nitration of phenols using [Msim]Cl/NaNO₂ system.





Figure 4. The competitive nitration reaction between phenols and anisole.

Table 1. Mono-nitration of phenols using [Msim]Cl/NaNO₂ system at room temperature.

5	a	b	c	d	e	f	g	h	i	j	k
X	F	Cl	Br	CN	Ph	CHO	OCH ₃	COCH ₃	CH ₂ Ph	NHOAc	4-HOC ₆ H ₄ -
Entry	Substrate	Product	Time (min)	Yield ^a (%)	M.p.(°C)						
					Found	Reported					
1	1	3	25	39	111-114	115 [35]					
		4		50	44-46	44 [35]					
2	5a	6a	15	94	72-73	73-74 [36]					
3	5b	6b	15	96	89-90	91 [37]					
4	5c	6c	15	95	87-89	84 [37]					
5	5d	6d	20	93	139-142	145 [36]					
6	5e	6e	25	87	61-64	61-64 [38]					
7	5f	6f	25	91	139-141	140-142 [19]					
8	5g	6g	20	93	52-54	54-56 [38]					
9	5h	6h	35	90	121-123	123 [37]					
10	5i	6i	25	91	60-62	66 [19]					
11	5j	6j	30	89	177-180	180 [38]					
12	5k	6k	40	78	181-183	183-184 [19]					
13	1	3,4	24 h	— ^b	—	—					

a: Isolated yields.


b: The reaction was not carried out in the absence of [Msim]Cl.


Figure 5. The proposed mechanism for the nitration of phenols.

mono- and di-nitrated products were obtained at 8% and 74% yields, respectively, within 50 min (Figure 6).

We also tried to use the [Msim]Cl/NaNO₂ system for the nitration of aniline; however, our attempt was not successful.

To compare the efficiency of the [Msim]Cl/NaNO₂ system for the nitration of phenols with the other systems, we have tabulated the results of the mono-nitration of *p*-bromophenol under different reaction


Figure 6. The di-nitration of *p*-methoxyphenol.

conditions in Table 2. As shown in Table 2, [Msim]Cl/NaNO₂ is superior to the previously reported reagent system.

CONCLUSION

In summary, we have introduced 3-methyl-1-sulfonic acid imidazolium chloride/NaNO₂ as an efficient and novel reagent for the efficient nitration of phenols. The promising points of this method are the simple experimental procedure, mild reaction conditions, safety, high yields, short reaction times and the minimization of chemical waste, as compared to their liquid phase counterparts.

EXPERIMENTAL

All Chemicals were purchased from the Merck or Fluka Chemical Company. The nitration products were characterized by comparison of their ¹H NMR, TLC and melting points with those in the authentic samples.

Table 2. Comparison of the results of nitration of *p*-boromophenol using different reagents.

Entry	Reagent System	Time (min)	Yield (%)	Ref.
1	Cs _{2.5} H _{0.5} PW ₁₂ O ₄₀ /Fe(NO ₃) ₃ /CH ₂ Cl ₂	360	98	[26]
2	H ₃ PW ₁₂ O ₄₀ /Fe(NO ₃) ₃ /CH ₂ Cl ₂	390	97	[26]
3	NaHSO ₄ .H ₂ O/NaNO ₃ /wet SiO ₂ /CH ₂ Cl ₂	130	70	[39]
4	Mg(HSO ₄) ₂ /NaNO ₃ /wet SiO ₂ /CH ₂ Cl ₂	130	78	[39]
5	Silica sulfuric acid/NaNO ₂ /wet SiO ₂ /CH ₂ Cl ₂	120	90	[38]
6	Tribromoisocyanuric acid/NaNO ₂ /wet SiO ₂ /CH ₂ Cl ₂	60	92	[19]
7	[Msim]Cl/NaNO ₂	15	95	—

Melting points were recorded on a Stuart Scientific Apparatus SMP3 (UK) in open capillary tubes.

Procedure for the Preparation of Ionic Liquid [Msim]Cl

A round-bottomed flask (100 mL) was charged with 1-methylimidazole (0.410 g, 5 mmol) in dry CH₂Cl₂ (50 mL) and then chlorosulfonic acid (0.605 g, 5.2 mmol) was added dropwise over a period of 5 min at room temperature. After the addition was completed, the reaction mixture was stirred for 20 min, stood for 5 min, and the CH₂Cl₂ was decanted. The residue was washed with dry CH₂Cl₂ (3 × 50 mL) and dried under vacuum to give [Msim]Cl as a viscous colorless oil at 92% yield, 0.912 g [13].

Typical Procedure for the Mono-Nitration of Phenol (1) Using [Msim]Cl/NaNO₂ System

A mixture of compounds consisting of phenol (0.094 g, 1 mmol), [Msim]Cl (0.616 g, 3.1 mmol) and NaNO₂ (0.069 g, 1 mmol) was ground vigorously in a mortar at room temperature. After completion of the reaction, as monitored by TLC, the reaction mixture was extracted with EtOAc (2 × 15 mL) and the organic extracts were combined. Anhydrous Na₂SO₄ (0.5 g) was added to the extracts. After 3 min, the resulting mixture was filtered, and the solvent was evaporated. The residue was a mixture of *o*- and *p*-nitrophenol. Only *o*-nitrophenol (**4**) is soluble in *n*-pentane. *n*-Pentane was added to the mixture, stirred and filtered; the residue was *p*-nitrophenol (**3**, 39% yield). The solvent of the filtrate was evaporated to give *o*-nitrophenol in 50%.

General Procedure for the Mono-Nitration of 4-Substituted Phenols Using [Msim]Cl/NaNO₂ System

A mixture of 4-substituted phenol (1 mmol), [Msim]Cl (0.616 g, 3.1 mmol) and NaNO₂ (0.069 g, 1 mmol) was ground vigorously in a mortar at room temperature. After completion of the reaction, as monitored by

TLC, the reaction mixture was extracted with EtOAc (2 × 15 mL) and the organic extracts were combined. Anhydrous Na₂SO₄ (0.5 g) was added to the extracts. After 3 min, the resulting mixture was filtered, and the solvent was evaporated to give the pure product.

Notice: The reactions displayed in Figure 4, as well as entry 13 of Table 1, were performed in a 10 mL round-bottomed flask with magnetic stirring.

Spectral Data of Ionic Liquid [Msim]Cl

Viscous colorless oil; ¹H NMR (DMSO-d₆, 300 MHz): δ (ppm) 3.77 (s, 3H, CH₃), 7.46 (s, 1H), 7.51 (s, 1H), 8.84 (s, 1H), 13.96 (s, 1H); ¹³C NMR (DMSO-d₆, 75 MHz): δ (ppm) 36.5, 120.6, 124.2, 138.6; MS (m/z): 199 (M⁺+1), 198 (M⁺), 183 (M⁺-CH₃), 163 (M⁺-Cl), 82 (M⁺-ClSO₃H), 67 (M⁺-CH₄ClSO₃); Anal. Calcd. for C₄H₇ClN₂O₃S: C, 24.19; H, 3.55; N, 14.10. Found: C, 24.41; H, 3.69; N, 13.92 .

ACKNOWLEDGMENTS

The authors gratefully acknowledge partial support of this work by the Research Affairs Office of Bu-Ali Sina University (Grant number 32-1716 entitled “development of chemical methods, reagent and molecules”), and the Center of Excellence for the Development of Chemical Methods (CEDCM), Hamedan, I.R. Iran.

REFERENCES

1. Rogers, R.D., *Ionic Liquids as Green Solvents: Progress and Prospects*, American Chemical Society, Washington, DC (2005).
2. Zare, A., Moosavi-Zare, A.R., Hasaninejad, A., Parhami, A., Khalafi-Nezhad, A. and Beyzavi, M.H. “Green, catalyst-free protocol for the efficient synthesis of *N*-sulfonyl aldimines and ketimines in ionic liquid [bmim]Br”, *Synth. Commun.*, **39**, pp. 3156-3165 (2009).
3. Zare, A., Parhami, A., Moosavi-Zare, A.R., Hasaninejad, A., Khalafi-Nezhad, A. and Beyzavi, M.H. “A

- catalyst-free protocol for the green and efficient condensation of indoles with aldehydes in ionic liquids”, *Can. J. Chem.*, **87**, pp. 416-421 (2009).
- Zare, A., Hasaninejad, A., Khalafi-Nezhad, A., Moosavi Zare, A.R. and Parhami, A. “Organic reactions in ionic liquids: MgO as efficient and reusable catalyst for the Michael addition of sulfonamides to α , β -unsaturated esters under microwave irradiation”, *ARKIVOC*, **13**, pp. 105-115 (2007).
 - Zare, A., Hasaninejad, A., Moosavi Zare, A.R., Parhami, A., Sharghi, H. and Khalafi-Nezhad, A. “Zinc oxide as a new, highly efficient, green, and reusable catalyst for microwave-assisted Michael addition of sulfonamides to α , β -unsaturated esters in ionic liquids”, *Can. J. Chem.*, **85**, pp. 438-444 (2007).
 - Zare, A., Hasaninejad, A., Safinejad, R., Moosavi-Zare, A.R., Beyzavi, M.H., Khalafi-Nezhad, A., Miralai-Moreddi, M., Dehghani, E. and Kazerooni-Mojarrad, P. “Ionic liquid-accelerated Michael addition of pyrimidine and purine nucleobases to α , β -unsaturated esters: a rapid approach to carbocyclic nucleosides synthesis”, *ARKIVOC*, **16**, pp. 51-64 (2008).
 - Miao, C.-X., He, L.-N., Wang, G.-Q. and Wang, J.-L. “TEMPO and carboxylic acid functionalized imidazolium salts/sodium nitrite: An efficient, reusable, transition metal-free catalytic system for aerobic oxidation of alcohols”, *Adv. Synth. Catal.*, **351**, pp. 2209-2216 (2009).
 - Gu, D.-G., Ji, S.-J., Jiang, Z.-Q., Zhou, M.-F. and Loh, T.-P. “An efficient synthesis of bis(indolyl)methanes catalyzed by recycled acidic ionic liquid”, *Synlett*, pp. 959-962 (2005).
 - Sonu, N.G., Kad, G.L. and Singh, J. “Acidic ionic liquid [bmim]HSO₄: An efficient catalyst for acetalization and thioacetalization of carbonyl compounds and their subsequent deprotection”, *Catal. Commun.*, **8**, pp. 1323-1328 (2007).
 - Dabiri, M., Salehi, P., Baghbanzadeh, M., Shakouri, M., Otokesh, S., Ekrami, T. and Doosti, R. “Efficient and eco-friendly synthesis of dihydropyrimidinones, bis(indolyl)methanes, and *N*-alkyl and *N*-arylimides in ionic liquids”, *J. Iran. Chem. Soc.*, **4**, pp. 393-401 (2007).
 - Salehi, P., Zolfigol, M.A., Shirini, F. and Baghbanzadeh, M. “Silica sulfuric acid and silica chloride as efficient reagents for organic reactions”, *Curr. Org. Chem.*, **10**, pp. 2171-2189 (2006).
 - Shirini, F., Zolfigol, M.A., Salehi, P. and Abedini, M. “Applications of some metal hydrogen sulfates in organic transformations”, *Curr. Org. Chem.*, **12**, pp. 183-202 (2008).
 - Zolfigol, M.A., Khazaei, A., Moosavi-Zare, A.R. and Zare, A. “Ionic liquid 3-methyl-1-sulfonic acid imidazolium chloride as a novel and highly efficient catalyst for the very rapid synthesis of bis(indolyl)methanes under solvent-free conditions”, *Org. Prep. Proced. Int.*, **42**, pp. 95-102 (2010).
 - Min, S. and Shi-Cong, C. “Perfluorinated rare earth metals catalyzed nitration of aromatic compounds”, *J. Fluor. Chem.*, **113**, pp. 207-209 (2002).
 - Esakkidurai, T. and Pitchumani, K. “Zeolite-mediated regioselective nitration of phenol in solid state”, *J. Mol. Catal. A: Chem.*, **185**, pp. 305-309 (2002).
 - Dagade, S.P., Waghmode, S.B., Kadam, V.S. and Dongare, M.K. “Vapor phase nitration of toluene using dilute nitric acid and molecular modeling studies over beta zeolite”, *Appl. Catal. A: Gen.*, **226**, pp. 49-61 (2002).
 - Iranpoor, N., Firouzabadi, H. and Heydari, R. “Ionic complex of N₂O₄ with 18-crown-6: A highly efficient and selective reagent for nitration of phenols”, *Synth. Commun.*, **29**, pp. 3295-3302 (1999).
 - Zolfigol, M.A., Ghaemi, E. and Madrakian, E. “Trichloroisocyanuric acid/NaNO₂ as a novel heterogeneous system for the selective mononitration of phenols under mild conditions”, *Synlett*, pp. 191-194 (2003).
 - Niknam, K., Zolfigol, M.A., Madrakian, E. and Ghaemi, E. “Tribromoisocyanuric acid/NaNO₂: A new reagent for mononitration of phenols under mild and heterogeneous conditions”, *South Afr. J. Chem.*, **60**, pp. 109-112 (2007).
 - Tasneem, A.M.M., Rajanna, K.C. and Saiparakash, P.K. “Ammonium nickel acid sulphate mediated nitration of aromatic compounds with nitric”, *Synth. Commun.*, **31**, pp. 1123-1127 (2001).
 - Heravi, M., Benmorad, T., Bakhtiari, K., Bamoharram, F. and Oskooie, H. “H_{3+x}PMO_{12-x}V_xO₄₀ (heteropolyacids)-catalyzed regioselective nitration of phenol to *o*-nitrophenol in heterogeneous system”, *J. Mol. Catal. A: Chem.*, **264**, pp. 318-321 (2007).
 - Hajipour, A.R. and Ruoho, A. “Nitric acid in the presence of P₂O₅ supported on silica gel - a useful reagent for nitration of aromatic compounds under solvent-free conditions”, *Tetrahedron Lett.*, **46**, pp. 8307-8310 (2005).
 - Tajik, H., Zolfigol, M.A., Albadi, J. and Eslami, R. “Nitration of some aromatic compounds by sodium nitrate in the presence of benzyltriphenylphosphonium peroxodisulfate”, *Synth. Commun.*, **37**, pp. 2771-2776 (2007).
 - Anuradha, V., Srinivas, P.V., Aparna, P. and Madhusudana Rao, J. “*p*-Toluenesulfonic acid catalyzed regiospecific nitration of phenols with metal nitrates”, *Tetrahedron Lett.*, **47**, pp. 4933-4935 (2006).
 - Tajik, H., Niknam, K. and Parsa, F. “Using acidic ionic liquid 1-butyl-3-methylimidazolium hydrogen sulfate in selective nitration of phenols under mild conditions”, *J. Iran. Chem. Soc.*, **6**, pp. 159-164 (2009).
 - Amani, K. and Maleki, F. “Catalytic effects of some Keggin-type heteropoly acids and polyoxometalates on selective nitration of phenols”, *J. Iran. Chem. Soc.*, **4**, pp. 238-243 (2007).

27. Firouzabadi, H., Iranpoor, N. and Zolfigol, M.A. "Dinitrogen tetroxide complexes of iron and copper nitrates as new reagents for selective mono- and dinitration of phenolic compounds", *Synth. Commun.*, **27**, pp. 3301-3311 (1997).
28. Zolfigol, M.A. "Silica sulfuric acid/ NaNO_2 As a novel heterogeneous system for production of thionitrites and disulfides under mild conditions", *Tetrahedron*, **57**, pp. 9509-95011 (2001).
29. Maleski, R.J., Kluge, M. and Sicker, D. "A facile access to substituted 2-nitrosophenols and 2-nitrophenols via regioselective nitrosation of resorcinol monoethers", *Synth. Commun.*, **25**, pp. 2327-2335 (1995).
30. Zolfigol, M.A., Kiany-Borazjani, M., Sadeghi, M.M., Mohammadpoor-Baltork, I. and Memarian, H.R. "An efficient method for the oxidation of Hantzsch 1,4-dihydropyridine to their corresponding pyridine derivatives under mild and heterogeneous conditions", *Synth. Commun.*, **30**, pp. 551-558 (2000).
31. Ishikawa, T., Watanabe, T., Tanigawa, H., Saito, T., Kotake, K.I., Ohashi, Y. and Ishii, H. "Nitrosation of phenolic substrates under mildly basic conditions: Selective preparation of *p*-quinone monooximes and their antiviral activities", *J. Org. Chem.*, **61**, pp. 2774-2779 (1996).
32. Zolfigol, M.A., Zebarjadian, M.H., Chehardoli, G.A., Keypour, H., Salehzadeh, S. and Shamsipur, M. "*N*-Nitrosation of secondary amines with $[\text{NO}^+ \cdot \text{crown.H}(\text{NO}_3)_2]^-$ ", *J. Org. Chem.*, **66**, pp. 3619-3620 (2001).
33. Lehnig, M. "The mechanism of the nitration of donor-activated benzenes with nitric and nitrous acid as studied by ^{15}N CIDNP", *Tetrahedron Lett.*, **40**, pp. 2299-2302 (1999).
34. Zolfigol, M.A., Madrakian, E. and Ghaemi, E. "Trichloroisocyanuric acid/ NaNO_2 /wet SiO_2 as an efficient system for the selective dinitration of phenols under solvent-free conditions", *Synlett*, pp. 2222-2224 (2003).
35. Furniss, B.S., Hannaford, A.J., Smith, P.W.G. and Tatchell, A.R., *Vogel's Text Book of Practical Organic Chemistry*, 4th Edn., Longman: London & New York, USA (1986).
36. Cornelis, A., Laszlo, P. and Pennetreau, P. "Nitration of phenols by clay-supported ferric nitrate", *Bull. Soc. Chim. Belg.*, **93**, pp. 961-972 (1984).
37. Poirier, J.M. and Vottero, C. "Mononitration de phenols par des nitrates metalliques", *Tetrahedron*, **45**, pp. 1415-1422 (1989).
38. Zolfigol, M.A., Madrakian, E. and Ghaemi, E. "Silica sulfuric acid/ NaNO_2 as a novel heterogeneous system for the nitration of phenols under mild conditions", *Molecules*, **7**, pp. 734-742 (2002).
39. Zolfigol, M.A., Madrakian, E. and Ghaemi, E. "Nitration of phenols under mild and heterogeneous conditions", *Molecules*, **6**, pp. 614-620 (2001).

BIOGRAPHIES

Ardeshir Khazaei was born in 1948 in Nahavand, Iran. He obtained BS and MS degrees from Pittsburgh University, USA, in 1975, and with Professor John Crendel, in 1977, respectively. He received his PhD from Manchester University, UK, with Professor Ronald Erick Bank, in 1990. He became a faculty member of Bu-Ali Sina University, Iran, a Professor in 2003, and national outstanding Professor in 2006. His research includes: Discovery and Development of Polymer Supported Reagent, Synthesis and Application of New N-Halo Compounds such as N-Halo Sulfonamides.

Mohammad Ali Zolfigol was born in 1966 in Salehabad (Ashtian), Iran. He obtained his BS from Arak University, Iran, his MS from Isfahan University of Technology, with Professor Shadpour Mallakpour, and his PhD from Shiraz University, with Professor Nasser Iranpoor. He became a faculty member of Bu-Ali Sina University in 1997 and Professor in 2005. In 2003, he was selected as distinguished researcher by the Ministry of Science, Research and Technology of Iran. He was also awarded at the Khwarizmi International Festival and at COMSTEC, in 2008. His research interests include: the Discovery and Development of New Synthetic Methods by the Synthesis and Application of New Solid-Supported Reagents, especially Silica-Based Resins.

Ahmad Reza Moosavi-Zare was born in Shiraz, Iran, in 1974. He received his BS degree in Chemistry from Yazd University, Iran, in 1998, and his MS degree in Organic Chemistry from Mohaghegh Ardebili University (Iran) in 2006. He is currently working on his PhD in Organic Chemistry under the supervision of Professor Mohammad Ali Zolfigol and Professor Ardeshir Khazaei in the Organic Chemistry Department at Bu-Ali Sina University, Hamadan, Iran. His research interests include: Applications of Ionic Liquids, Solvent-Free Conditions and Microwave Irradiation in Organic Synthesis, and Green Chemistry.

Abdolkarim Zare was born in Bushehr, Iran, in 1977. He received his BS degree in Chemistry from Yazd University, Iran, in 1998, and his MS and PhD degrees in Organic Chemistry from Shiraz University, Iran, with Professor Hashem Sharghi, in 2002, and with Professor Ali Khalafi-Nezhad, in 2006, respectively. He is faculty member of Payame Noor University (PNU). His research field includes: Applications of Solvent-Free Conditions, Ionic Liquids and Microwave Irradiation in Organic Synthesis, and Study of Methodology in Organic Chemistry.