

FUZZY IDEALS OF NEAR-RINGS WITH INTERVAL VALUED MEMBERSHIP FUNCTIONS

B. Davvaz*

Department of Mathematics, Yazd University, Yazd, Islamic Republic of Iran

Abstract

In this paper, for a complete lattice \mathcal{L} , we introduce interval-valued \mathcal{L} -fuzzy ideal (prime ideal) of a near-ring which is an extended notion of fuzzy ideal (prime ideal) of a near-ring. Some characterization and properties are discussed.

1. Introduction

Zadeh in [19] introduced the concept of a fuzzy subset of a non-empty set X as a function from X to $[0,1]$. Goguen in [10] generalized the fuzzy subset of X , to \mathcal{L} -fuzzy subset, as a function from X to a lattice \mathcal{L} .

Since Rosenfeld [18] in 1971 introduced the concept of fuzzy subgroups following Zadeh, fuzzy algebra theory has been developed by many researchers. Liu [12] defined the fuzzy ideals of a ring and discussed the operations on fuzzy ideals. Mukherjee and Sen [16], Malik and Mordeson [16], Mashinchi and Zahedi [14], Zahedi [21], shown the meaning of the fuzzy prime ideals and its nature. The notion of fuzzy ideals and its properties were applied to various areas: distributive lattice [2], BCK-algebra [17], hyperrings [6,8], near-rings [1,11], hypernear-rings [7].

In 1975, Zadeh [20] introduced the concept of interval-valued fuzzy subsets (in short written by i-v fuzzy sets), where the values of the membership functions are intervals of numbers instead of the numbers. In [4], Biswas defined interval-valued fuzzy subgroups of the same nature of Rosenfeld's fuzzy subgroups.

In this paper, for a complete lattice \mathcal{L} , we define Interval-valued \mathcal{L} -fuzzy ideals (prime ideals) of a near-

ring, and we obtain an exact analogue of fuzzy ideals. In particular, we show there exists a one-to-one correspondence between the set of all f -invariant i-v \mathcal{L} -fuzzy prime ideals of R and the set of all i-v \mathcal{L} -fuzzy prime ideals of R' , where R and R' are near-rings and f is a homomorphism from R onto R' .

2. Basic Definitions

From now on this paper \mathcal{L} is a complete lattice [3], i.e. there is a partial order \leq on \mathcal{L} such that, for any $S \subseteq \mathcal{L}$, infimum of S and supremum of S exist and these will be denoted by $\bigwedge_{s \in S} \{s\}$ and $\bigvee_{s \in S} \{s\}$, respectively. In

particular for any elements $a, b \in \mathcal{L}$, in $f\{a, b\}$ and $\sup\{a, b\}$ will be denoted by $a \wedge b$ and $a \vee b$, respectively. Also, \mathcal{L} is a ditributive lattice with a least element 0 and a greatest element 1. If $a, b \in \mathcal{L}$; we write $a \geq b$ if $b \leq a$, and $a > b$ if $a \geq b$ and $a \neq b$.

Definition 2.1. Given two elements $a, b \in \mathcal{L}$ with $a \leq b$, we define the following closed interval set:

$$[a, b] = \{c \in \mathcal{L} | a \leq c \leq b\}.$$

Suppose $\mathcal{D}(\mathcal{L})$ denotes the family of all closed intervals of \mathcal{L} .

Keywords: Fuzzy set; Near-ring; Fuzzy ideal; Level set

* E-mail: davvaz@yazduni.net

Definition 2.2. Let $I_1 = [a_1, b_1]$, $I_2 = [a_2, b_2]$ and $I_i = [a_i, b_i]$ be elements of $\mathcal{D}(\mathcal{L})$ then we define

$$I_1 \wedge I_2 = [a_1 \wedge a_2, b_1 \wedge b_2],$$

$$I_1 \vee I_2 = [a_1 \vee a_2, b_1 \vee b_2],$$

$$\bigwedge_i \{I_i\} = [\bigwedge_i \{a_i\}, \bigwedge_i \{b_i\}],$$

$$\bigvee_i \{I_i\} = [\bigvee_i \{a_i\}, \bigvee_i \{b_i\}].$$

We call $I_2 \leq I_1$ if and only if $a_2 \leq a_1$ and $b_2 \leq b_1$.

Definition 2.3. Let X be a non-empty set. An \mathcal{L} -fuzzy subset F defined on X is given by

$$F = \{(x, \mu_F(x)) \mid x \in X\}, \text{ where } \mu_F : X \rightarrow \mathcal{L}.$$

Definition 2.4. Let X be a non-empty set. An interval-valued \mathcal{L} -fuzzy subset F defined on X is given by

$$F = \{(x, [\mu_F^L(x), \mu_F^U(x)]) \mid x \in X\},$$

where μ_F^L and μ_F^U are two \mathcal{L} -fuzzy subsets of X such that $\mu_F^L(x) \leq \mu_F^U(x)$ for all $x \in X$.

Suppose $\hat{\mu}_F(x) = [\mu_F^L(x), \mu_F^U(x)]$. If $\mu_F^L(x) = \mu_F^U(x) = c$ where $0 \leq c \leq 1$, then we have $\hat{\mu}_F(x) = [c, c]$ which we also assume, for the sake of convenience, to belong to $\mathcal{D}(\mathcal{L})$. Thus $\hat{\mu}_F(x) \in \mathcal{D}(\mathcal{L})$ for all $x \in X$. Therefore the i-v fuzzy subset F is given by

$$F = \{(x, \hat{\mu}_F(x)) \mid x \in X\}, \text{ where } \hat{\mu}_F : X \rightarrow \mathcal{D}(\mathcal{L}).$$

Definition 2.5. Let f be a mapping from a set X into a set Y . Let A be an i-v \mathcal{L} -fuzzy subset of X . then the image of A , i.e., $f[A]$ is the i-v fuzzy subset of Y with the membership function defined by

$$\hat{\mu}_{f[A]}(y) = \begin{cases} \bigvee_{z \in f^{-1}(y)} \{\hat{\mu}_A(z)\} & \text{if } f^{-1}(y) \neq \emptyset \\ [0, 0] & \text{otherwise} \end{cases} \quad \text{for all } y \in Y$$

Let B be an i-v \mathcal{L} -fuzzy subset of Y . Then the inverse image of B , i.e., $f^{-1}[B]$ is the i-v \mathcal{L} -fuzzy subset of X with the membership function given by

$$\hat{\mu}_{f^{-1}[B]} = \hat{\mu}_B(f(x)) \quad \text{for all } x \in X.$$

Definition 2.6. Let X and Y be any two non-empty sets and $f : X \rightarrow Y$ be any function. An i-v \mathcal{L} -fuzzy subset of F of X is called f -invariant if

$$f(x) = f(y) \Rightarrow \hat{\mu}_F(x) = \hat{\mu}_F(y), \quad \text{where } x, y \in X.$$

Definition 2.7. A non-empty set R with two binary operations $+$ and \cdot is called a near-ring [5,15] if

- 1) $(R, +)$ is a group,
- 2) (R, \cdot) is a semigroup,
- 3) $x \cdot (y + z) = x \cdot y + x \cdot z$ for all $x, y, z \in R$.

To be more precise, they are left near-rings because the left distributive law is satisfied. We will use the word near-ring to mean left near-ring. We denote xy instead of $x \cdot y$. Note that $x0 = 0$ and $x(-y) = -xy$ but in general $0x \neq 0$ for all $x \in R$ [15, Lemma 1.10]. A near-ring R is called a zero symmetric if $0x = 0$ for all $x \in R$.

Definition 2.8. Let $(R, +, \cdot)$ be a near-ring. An ideal of R is a subset I of R such that

- 1) $(I, +)$ is a normal subgroup of $(R, +)$,
- 2) $RI \subseteq I$,
- 3) $(r + i)s - rs \in I$ for all $i \in I$ and $r, s \in R$.

Note that if I satisfies (1) and (2) then it is called a left ideal of R . If I satisfies (1) and (3) then it is called a right ideal of R . Let P be an ideal of R . We call P a prime ideal if for any ideal $I, J \subseteq R$, $IJ \subseteq P$ then $I \subseteq P$ or $J \subseteq P$.

i-v \mathcal{L} -Fuzzy Ideals in a Near-Ring

In this section first we define interval-valued \mathcal{L} -fuzzy subnear-rings and ideals and then we explain some results in this connection.

Definition 3.1. Let $(R, +, \cdot)$ be a near-ring. An i-v \mathcal{L} -fuzzy subset F of R is called an i-v \mathcal{L} -fuzzy subnear-ring, if the following hold:

- 1) $\hat{\mu}_F(x) \wedge \hat{\mu}_F(y) \leq \hat{\mu}_F(x - y)$ for all $x, y \in R$,

$$2) \hat{\mu}_F(x) \wedge \hat{\mu}_F(y) \leq \hat{\mu}_F(x \cdot y) \text{ for all } x, y \in R.$$

Furthermore F is called an i-v \mathcal{L} -fuzzy ideal of R , if F is an i-v \mathcal{L} -fuzzy subnear-ring of R and

$$3) \hat{\mu}_F(x) = \hat{\mu}_F(y + x - y) \text{ for all } x, y \in R,$$

$$4) \hat{\mu}_F(x) \leq \hat{\mu}_F(xy) \text{ for all } x, y \in R,$$

$$5) \hat{\mu}_F(i) \leq \hat{\mu}_F((x+i)y - xy) \text{ for all } x, y, i \in R.$$

Note that F is an i-v \mathcal{L} -fuzzy left ideal of R if it satisfies (1), (3) and (4), and F is an i-v \mathcal{L} -fuzzy right ideal of R if it satisfies (1), (2), (3) and (5).

Now, we give an example of an i-v \mathcal{L} -fuzzy ideal of a near-ring.

Example 3.2. Let $R = \{0, a, b, c\}$ be a set with two binary operations as follows:

$+$	0	a	b	c
0	0	a	b	c
a	a	0	c	b
b	b	c	a	0
c	c	b	0	a

\cdot	0	a	b	c
0	0	0	0	0
a	0	0	0	0
b	0	0	0	0
c	0	0	a	a

Then $(R, +, \cdot)$ is a near-ring. Define an i-v \mathcal{L} -fuzzy subset F by membership function $\hat{\mu}_F: R \rightarrow \mathcal{D}(\mathcal{L})$ by $\hat{\mu}_F(b) = \hat{\mu}_F(c) < \hat{\mu}_F(a) < \hat{\mu}_F(0)$. Then F is an i-v \mathcal{L} -fuzzy ideal of R .

Lemma 3.3. For an i-v \mathcal{L} -fuzzy ideal F of a near-ring R , we have

$$\hat{\mu}_F(x) = \hat{\mu}_F(-x) \leq \hat{\mu}_F(0) \text{ for all } x \in R.$$

Proposition 3.4. Let F be an i-v \mathcal{L} -fuzzy ideal of R . If $\hat{\mu}_F(x - y) = \hat{\mu}_F(0)$ then $\hat{\mu}_F(x) = \hat{\mu}_F(y)$.

Proof. Assume that $\hat{\mu}_F(x - y) = \hat{\mu}_F(0)$. Then

$$\begin{aligned} \hat{\mu}_F(x) &= \hat{\mu}_F(x - y + y) \\ &\geq \hat{\mu}_F(x - y) \wedge \hat{\mu}_F(y) \\ &= \hat{\mu}_F(0) \wedge \hat{\mu}_F(y) \\ &= \hat{\mu}_F(y). \end{aligned}$$

Similarly, using $\hat{\mu}_F(y - x) = \hat{\mu}_F(x - y) = \hat{\mu}_F(0)$, we get

$$\hat{\mu}_F(y) \geq \hat{\mu}_F(x).$$

Corollary 3.5. $[\mu_F^L, \mu_F^U]$ is an i-v \mathcal{L} -fuzzy ideal of a near-ring R if and only if μ_F^L, μ_F^U are \mathcal{L} -fuzzy ideals of R . Now, we define

$$F_t^L = \{x \in X \mid \mu_F^L(x) \geq t\} \quad \text{and} \\ F_s^U = \{x \in X \mid \mu_F^U(x) \geq s\}.$$

Then $\hat{\mu}_F$ is an i-v \mathcal{L} -fuzzy ideal of R if and only if for every t, s where $0 \leq t \leq s \leq 1$, $F_t^L, F_s^U \neq \emptyset$ are ideals of R .

Definition 3.6. Let F_1 and F_2 be two i-v \mathcal{L} -fuzzy subsets of a near-ring R . Then $F_1 \cap F_2$ and $F_1 \circ F_2$ are defined as follows:

$$\hat{\mu}_{F_1 \cap F_2} = \hat{\mu}_{F_1}(x) \wedge \hat{\mu}_{F_2}(x),$$

$$\hat{\mu}_{F_1 \circ F_2}(x) = \begin{cases} \bigvee_{x=yz} \{\hat{\mu}_{F_1}(y) \wedge \hat{\mu}_{F_2}(z)\} \\ [0, 0] & \text{if } x \text{ is not expressible as } x = yz. \end{cases}$$

Lemma 3.7. Let R be a near-ring, we have

- 1) If F_1, F_2 are two i-v \mathcal{L} -fuzzy ideals of R (right or left) then $F_1 \cap F_2$ is an i-v \mathcal{L} -fuzzy ideal of R (right or left), respectively;
- 2) If R is a zero-symmetric and if F_1 is an i-v \mathcal{L} -fuzzy right ideal and F_2 is an i-v \mathcal{L} -fuzzy left ideal, then $F_1 \circ F_2 \subseteq F_1 \cap F_2$.

Proof. (1) It is an immediate consequence of Corollary 3.5 and Definition 3.6.

(2) We assume R is a zero symmetric near-ring. If $\hat{\mu}_{F_1 \circ F_2}(x) = 0$, there is nothing to prove. Otherwise

$$\hat{\mu}_{F_1 \circ F_2}(x) = \bigvee_{x=yz} \{\hat{\mu}_{F_1}(y) \wedge \hat{\mu}_{F_2}(z)\}.$$

Since F_1 is an i-v \mathcal{L} -fuzzy left ideal, we have

$$\hat{\mu}_{F_1}(z) \leq \hat{\mu}_{F_1}(yz) = \hat{\mu}_{F_1}(x),$$

and since F_1 is an i-v \mathcal{L} -fuzzy right ideal, we have

$$\hat{\mu}_{F_1}(x) = \hat{\mu}_{F_1}(yz) = \hat{\mu}_{F_1}((0+y)z - 0z) \geq \hat{\mu}_{F_1}(y).$$

Therefore

$$\hat{\mu}_{F_1 \circ F_2}(x) \leq \hat{\mu}_{F_1}(x) \wedge \hat{\mu}_{F_2}(x) = \hat{\mu}_{F_1 \cap F_2}(x).$$

Definition 3.8. Let X be a non-empty set and F be an i-v \mathcal{L} -fuzzy subset of X . Then we define

$$F_{[t,s]} = \{x \in X \mid \hat{\mu}_F(x) \geq [t, s]\}.$$

The set $F_{[t,s]}$ is called the “level set” of F .

It is easy to see that $F_{[t,s]} = F_t^L \cap F_s^U$.

Now, we obtain the relation between an i-v \mathcal{L} -fuzzy ideal and level ideals. This relation is expressed in terms of a necessary and sufficient condition.

Theorem 3.9. Let R be a near-ring and F be an i-v \mathcal{L} -fuzzy subset of R . Then F is an i-v \mathcal{L} -fuzzy ideal of R if and only if for every t, s where $0 \leq t \leq s \leq 1$, $F_{[t,s]} \neq \emptyset$ is an ideal of R .

Proof. The proof is similar to the proof of Theorem 3.4 of [7], by considering the suitable modification with using Definitions 2.4 and 3.1.

Definition 3.10. An i-v \mathcal{L} -fuzzy ideal P of a near-ring R is said to be prime if P is not constant function and for any i-v \mathcal{L} -fuzzy ideals F_1, F_2 in R , $F_1 \circ F_2 \subseteq P$ implies $F_1 \subseteq P$ or $F_2 \subseteq P$.

Proposition 3.11. Let P be an i-v \mathcal{L} -fuzzy prime ideal of a near-ring R . Define

$$\pi = \{x \in R \mid \hat{\mu}_P(x) = \hat{\mu}_P(0)\},$$

then π is a prime ideal in R .

Proof. The proof is similar to the proof of Theorem 3.7 in [1].

Proposition 3.12. Let R be a near-ring and F_1, F_2 are i-v \mathcal{L} -fuzzy prime ideals of R , then $F_1 \cap F_2$ is an i-v \mathcal{L} -fuzzy prime if and only if $F_1 \subseteq F_2$ or $F_2 \subseteq F_1$.

Proof. The proof is straightforward, in view of the fact that $F_1 \circ F_2 \subseteq F_1 \cap F_2$.

We have the following corollary which plays an important role in the determination of i-v \mathcal{L} -fuzzy prime ideals.

Corollary 3.13. Let R be a near-ring. Then every ideal of R is a level ideal of an i-v \mathcal{L} -fuzzy ideal of R .

Proof. Let I be any ideal of a near-ring R and let $[\alpha_1, \alpha_2] \leq [\beta_1, \beta_2] \neq [0, 0]$ be elements in $\mathcal{D}(\mathcal{L})$. Then the fuzzy subset F is defined as follows:

$$\hat{\mu}_F(x) = \begin{cases} [\beta_1, \beta_2] & \text{if } x \in I \\ [\alpha_1, \alpha_2] & \text{otherwise.} \end{cases}$$

We have $I = F_{[\beta_1, \beta_2]}$ and by Theorem 3.9, it is enough to prove that F is an i-v \mathcal{L} -fuzzy ideal.

An element $[\alpha_1, \alpha_2] \neq [1, 1]$ in $\mathcal{D}(\mathcal{L})$ is called “prime” if for any $[a_1, a_2], [b_1, b_2] \in \mathcal{D}(\mathcal{L})$, $[a_1, a_2] \wedge [b_1, b_2] \leq [\alpha_1, \alpha_2]$ implies either $[a_1, a_2] \leq [\alpha_1, \alpha_2]$ or $[b_1, b_2] \leq [\alpha_1, \alpha_2]$.

Theorem 3.14. Let I be a prime ideal of a near-ring R and let $[\alpha_1, \alpha_2]$ a prime element in $\mathcal{D}(\mathcal{L})$. Let P be the fuzzy subset of R defined by

$$\hat{\mu}_P(x) = \begin{cases} [1, 1] & \text{if } x \in I \\ [\alpha_1, \alpha_2] & \text{otherwise.} \end{cases}$$

Then P is an i-v \mathcal{L} -fuzzy prime ideal.

Proof. By Corollary 3.13, P is clearly a non-constant i-v \mathcal{L} -fuzzy ideal. Let F_1 and F_2 be any i-v \mathcal{L} -fuzzy ideals and let $F_1 \not\subseteq P, F_2 \not\subseteq P$. Then there exist x, y in R , such that $\hat{\mu}_{F_1}(x) \not\leq \hat{\mu}_P(x)$ and $\hat{\mu}_{F_2}(y) \not\leq \hat{\mu}_P(y)$. This implies that $\hat{\mu}_P(x) = \hat{\mu}_P(y) = [\alpha_1, \alpha_2]$ and hence $x \notin I$ and $y \notin I$. Since I is prime, there exists $r \in R$ such that $xry \notin I$. Now, we have $\hat{\mu}_{F_1}(x) \not\leq [\alpha_1, \alpha_2]$ and $\hat{\mu}_{F_2}(ry) \not\leq [\alpha_1, \alpha_2]$ (otherwise $\hat{\mu}_{F_2}(y) \leq [\alpha_1, \alpha_2]$ and since $[\alpha_1, \alpha_2]$ is prime, $\hat{\mu}_{F_1}(x) \wedge \hat{\mu}_{F_2}(ry) \leq [\alpha_1, \alpha_2]$ and hence $(F_1 \circ F_2)(xry) \leq [\alpha_1, \alpha_2] = \hat{\mu}_P(xry)$ so that $F_1 \circ F_2 \subseteq P$. Hence P is an i-v \mathcal{L} -fuzzy prime.

Lemma 3.15. Let f be a mapping from a non-empty set X into a non-empty set Y , and let A, B are i-v \mathcal{L} -fuzzy subsets of X, Y , respectively, such that

$$\hat{\mu}_A = [\mu_A^L, \mu_A^U]: X \rightarrow \mathcal{D}(\mathcal{L}) \text{ and}$$

$$\hat{\mu}_B = [\mu_B^L, \mu_B^U]: Y \rightarrow \mathcal{D}(\mathcal{L}).$$

Then

$$\hat{\mu}_{f[A]} = [f(\mu_A^L), f(\mu_A^U)] \text{ and}$$

$$\hat{\mu}_{f^{-1}[B]} = [f^{-1}(\mu_B^L), f^{-1}(\mu_B^U)].$$

Using Lemma 3.15, the following propositions are obvious.

Proposition 3.16. Let f be a homomorphism from a near ring R onto a near-ring R' , and A be any f -invariant i-v \mathcal{L} -fuzzy prime ideal of R . Then $f[A]$ is an i-v \mathcal{L} -fuzzy prime ideal of R' .

Proposition 3.17. Let f be a homomorphism from a near ring R onto a near-ring R' , and B be any f -invariant i-v \mathcal{L} -fuzzy prime ideal of R' . Then $f^{-1}[B]$ is an i-v \mathcal{L} -fuzzy prime ideal of R .

Theorem 3.18. Let f be a homomorphism from a near ring R onto a near-ring R' , then the mapping $A \rightarrow f[A]$ defines a one-to-one correspondence between the set of all f -invariant i-v \mathcal{L} -fuzzy prime ideals of R and the set of all i-v \mathcal{L} -fuzzy prime ideals of R' .

References

1. Abou-Zaid, S. On fuzzy subnear-rings and ideals, *Fuzzy Sets and Systems*, **44**:139-46, (1991).
2. Yuan Bo and Wu Wangming, Fuzzy ideals on a distributive lattice, *Ibid.*, **35**: 231-40, (1990).
3. Birkhoff, G. *Lattice Theory*, Amer. Math. Soc. Colloq. Publ. Vol. 25, Amer. Math. Soc., Providence, RI, (1984).
4. Biswas, R. Rosenfeld's fuzzy subgroups with interval valued membership functions, *Fuzzy Sets and Systems*, **63**: 87-90, (1994).
5. Clay, J. R. *Near-Rings; Geneses and Applications*, Oxford, New York, (1992).
6. Davvaz, B. On H_v -rings and fuzzy H_v -ideals, *J. Fuzzy Math.*, **6**: 33-42, (1998).
7. Davvaz, B. On Hypernear-rings and Fuzzy Hyperideals, *Ibid.*, **7**: 745-53, (1999).
8. Davvaz, B. Some results on L -fuzzy H_v -ideals, *Pure Math. Appl.*, **10**: 31-40, (1990).
9. Dickson, L. E. Definitions of a group and a field by independent postulates, *Trans. Am. Math. Soc.*, **6**: 198-204, (1905).
10. Goguen, J. A. L -fuzzy sets, *J. Math. Anal. Appl.*, **18**: 145-74, (1967).
11. Seung Dong Kim and Hee Sik Kim, On fuzzy ideals of near-rings, *Bull. Korean Math. Soc.*, **33**: 593-601, (1996).
12. Liu, W. J. Fuzzy invariant subgroups and fuzzy ideals, *Fuzzy Sets and Systems*, **8**: 133-39, (1982).
13. Malik, D. S. and Mordeeson, J. N. Fuzzy prime ideals of a ring, *Ibid.*, **53**: 237-50, (1991).
14. Mashinchi, M. and Zahedi, M. M. On fuzzy ideals of a ring, *J. Sci. I. R. Iran*, **1**(3): 208-10, (1990).
15. Meldrum, J. D. P. *Near-Rings and Their Links with Groups*, Pitman, London, (1985).
16. Mukhrejee, T. K. and Sen, M. K. On fuzzy ideals in rings I, *Fuzzy Sets and Systems*, **21**: 99-104, (1987).
17. Ougen, X. Fuzzy BCK-algebra, *Math. Japonica*, **36**: 935-42, (1991).
18. Rosenfeld, A. Fuzzy groups, *J. Math. Anal.*, **35**: 512-17, (1971).
19. Zadeh, L. A. Fuzzy sets, *Inf. And Control.*, **8**: 338-53, (1965).
20. Zadeh, L. A. The concept of a linguistic variable and its application to approximate reasoning-1, *Ibid.*, **8**: 199-249, (1975).
21. Zahedi, M. M. A characterization of L -fuzzy prime ideals, *Fuzzy Sets and Systems*, **44**: 147-60, (1991).