

# Parasitic Contamination of Raw Vegetables in Amol, North of Iran

Abolghasem Siyadatpanah<sup>1</sup>; Fatemeh Tabatabaei<sup>2</sup>; Amir Emami Zeydi<sup>3,4</sup>; Adel Spotin<sup>5,6</sup>; Vahid Fallah-Omrani<sup>5,\*</sup>; Mehrdad Assadi<sup>7</sup>; Sepideh Moradi<sup>2</sup>; Ali Rostami<sup>5</sup>; Fatemeh Memari<sup>5</sup>; Fateme Hajialiani<sup>5</sup>

<sup>1</sup>Faculty of Paramedical Science, Mazandaran University of Medical Sciences, Sari, IR Iran

<sup>2</sup>Department of Parasitology and Mycology, Tehran University of Medical Sciences, Tehran, IR Iran

<sup>3</sup>Department of Nursing, Faculty of Nursing and Midwifery, Mazandaran University of Medical Sciences, Sari, IR Iran

<sup>4</sup>Faculty of Nursing and Midwifery, Mashhad University of Medical Sciences, Mashhad, IR Iran

<sup>5</sup>Department of Parasitology and Mycology, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

<sup>6</sup>Student Research Committee, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran

<sup>7</sup>Department of Microbiology, Faculty of Medicine, Tabriz Branch, Islamic Azad University, Tabriz, IR Iran

\*Corresponding author: Vahid Fallah-Omrani, Department of Parasitology and Mycology, Shahid Beheshti University of Medical Sciences, Tehran, IR Iran. Tel: +98-9117333391, Fax: +98-2172019014, E-mail: vomrani@gmail.com

Received: November 7, 2012; Revised: December 25, 2012; Accepted: February 2, 2013

**Background:** One of the most important ways of pathologic and non-pathologic parasites transmission is through eating raw vegetables. Iranian people believe that eating raw vegetables is an important way to get some essential vitamins and minerals.

**Objectives:** According to the high prevalence of pathologic parasites and also exporting different vegetables to the other provinces such as Tehran, we conducted this study to evaluate the parasitic pollution of raw vegetables in Amol city.

**Materials and Methods:** This is a descriptive and analytic study compromised of 200 samples from 10 different vegetables including spinach, garden cress, parsley, lettuce, peppermint, radish, green onion, basil, coriander and scallion. The samples were evaluated using timely-sediment method and then centrifuging concentrating. The obtained results were analyzed by SPSS software and  $\chi^2$ /Fisher's Exact Test.

**Results:** Results showed that 93 (46.5%) of all vegetables (collected from 20 regions) were contaminated with pathologic and non-pathologic parasites. Among these, spinach and coriander were the highest (17.2%) and the lowest (3.2%) contaminated samples, respectively. The isolated parasites were nematode larva 25.8%, *Giardia* cyst 22.5%, *Entamoeba coli* cyst 19.3%. However, the obtained results from 100 restaurants and kebab samples did not show any parasitic contamination.

**Conclusions:** Given hygiene issues and health status of people, the prevalence of some parasitic contaminations is the biggest problems among vegetable consumers. Using organized and new methods for irrigating the agricultural fields, preventing animals to enter the vegetable's farmlands and also improving people's knowledge on proper washing of vegetables can be useful in decreasing parasites prevalence.

**Keywords:** Parasites; Vegetables; Iran

## 1. Background

In many countries, especially in developing countries, parasitic infections are one of the most important health and economic problems, health extension, development and struggle against these diseases are important parts of mass programming in these countries. Some of the parasitic infections that are transmitted through food, soil or water are still the most important health concerns even in developed countries (1). Eating raw vegetables is a way of parasite transmission to human and the determination of vegetables contaminants and disinfecting were the main concern of health and therapeutic organiza-

tions, for many years (2). During the past years, by using man-made fertilizer and now because of using animal fertilizer, vegetables are infected by pathologic organisms and if we do not wash properly, they can transmit the parasites and other microorganisms to the human, and they can be a potential source of expanding infection in parasites life cycle (3). Different studies in all over the world has been performed in this regard and the results showed that there are noticeable parasitic contamination in different areas (4-10). Parasitic contaminations were studied in different researches in our country and

### Implication for health policy/practice/research/medical education:

Consuming vegetables among Iranian people could be taken into consideration from the viewpoint of parasitic contaminations which have impact on public health and socioeconomic.

Copyright © 2013, Infectious Diseases and Tropical Medicine Research Center; Published by Kowsar Corp. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

the results of such studies showed that parasitic prevalence is different in each region. In this respect, parasitic contamination in some areas of our country was about 13% and in some others was about 56% (11, 12).

## 2. Objectives

We performed this study to evaluate the parasitic pollution in Amol city, because there is different prevalence of vegetable contamination in different areas of our country, there is not any study about vegetables contamination in north area of our country, almost all of the people living in north of Iran are farming and many of them use animal fertilizer for their fields which can maintain the parasitic life cycle, there is high intake of vegetables among the people living in this area of the country.

## 3. Materials and Methods

This study was performed in 2012 in Amol city, North Iran. For the first step, the space of farming fields was evaluated and then all were divided into twenty different geographical areas. Twenty samples (each contains 500 g) containing ten vegetables including spinach, garden cress, parsley, lettuce, peppermint, radish, green onion, basil, coriander and scallion were collected. Hence, overall 200 samples were collected. Each sample kept in container containing four liter water and eight grams of anionic detergent for eight hours to separating larva, ova, and the cysts of the parasites. After eight hours, vegetables have been removed from the bucket and washed by high-pressure water for three times to complete separating parasites and adding them to the content of each bucket. Each bucket were positioned in quiet place and maintained for 14 hours in laboratory to get complete sedimentation of parasitic materials. Then, top layer of the content sent out and the sedimentation content of each bucket were collected in one liter bottles. During investigation, sediments were divided in several centrifuging tubes and then they were centrifuged for five minutes at 1500 rpm. Four different slides were made by using physiological serum and logol solution and then evaluated microscopically at 10X. and 40X. For the final step, the resulting data were evaluated and analyzed by using descriptive statistical methods and SPSS software and  $\chi^2$ /Fisher's Exact Test.

## 4. Results

There were pathologic and non-pathologic parasitic contaminations, in 93 cases (46.5%). The highest contamination was observed in spinach (16 cases, 17.2%) and the lowest amount of contamination was in coriander (3 cases, 3.2%). The contaminating agent in 47 cases was a uni-cellular and in 46 cases was a multi-cellular agent. In uni-cellular pathologic group, the highest prevalence was for *Giardia* cyst which had seen in 21 samples.

In uni-cellular non-pathologic group, the highest prevalence was for *Entamoeba coli* which had seen in 18 samples. *Entamoeba histolytica/dispar* was an important pathologic *Amoeba* which was reported. The highest level of contamination was observed in fields which were near the small and large lakes. There were five cases of entritic coccidian's oocytes which more of them were near villages and residency areas. Among multi-cellular group, nematode larva was the highest infective agent. Six *Trichostrongylus* eggs were the highest and two *Hymenolepis nana* eggs were the lowest contaminants among worm eggs. Other parasites in this study are shown in Table 1 and Figure 1.


## 5. Discussion

The results of the present study show that 46.5% of vegetables collected from Amol city lands were contaminated with parasites. The highest number was for *Giardia* cysts and soil-nematode larva. Moreover, there were different values of vegetable contaminations reported in other studies in other cities of Iran. Nazemi et al. showed that 38% of their samples, which were collected in Shahrood in 1390, were infected by parasites. Again, in that study, the highest prevalence was for *Giardia* cyst which is in accordance to the present study (13). Homayoni mentioned *Giardia* infection as a highest contaminant agent (14). According to the fields' irrigation, it is expected to see high contamination with these uni-cellular parasites. The second parasite, detected in this study, was *Entamoeba coli* compromised 19.3% of all parasites. These results are the same as the results of Nazemi et al. (13) and also Al-megrin et al. in Saudi Arabia (15). The isolation of this parasite does not have any pathologic importance, but it shows that vegetables are contaminated by human feces and it has a great importance (16).

*Entamoeba histolytica/dispar* compromised 3.2% of all parasitic contaminations of this study and also Jiroft studies (17). In this study, most of the strains were non-pathologic *dispar* ones. But, in respect to limited parasitological methods to isolate *Entamoeba dispar* from *Entamoeba histolytica* and due to the probability of pathologic strain pollutions, contamination of eatable vegetables in Amol city is of great importance. Rabditoide larva compromised 12% of multi-cellular pollutions which is in accordance to the results of Dehghani et al. from Yazd (18). According to the clinical experiences in laboratories of Amol city, after *Giardia*, the high frequent parasitic agent is *Strongyloides stercoralis*. Although these parasites are in plants and free living, these results is of importance due to the probability of contamination with *Strongyloides stercoralis* and some pathologic nematodes. 6.4% of all contaminations were belonged to *Trichostrongylus* egg which is in accordance to the other studies in Esfahan and Yazd (11, 18).

**Table 1.** Contaminants of the Applied Vegetables

Vegetables	Basil	Radish	Scallion	Spinach	Coriander	Green Onion	Green Mint	Parsley	Garden Cress	Lettuce	Total, No. (%)
<b>Parasite</b>											
Cyst of <i>Giardia</i>	1	0	5	4	2	1	1	1	1	5	21 (22.5)
Cyst of <i>Entamoeba coli</i>	2	3	2	1	0	0	3	2	4	1	18 (19.3)
Cyst of <i>Entamoeba histolytica/dispar</i>	0	0	1	1	0	0	0	1	0	0	3 (3.2)
Oocyst of <i>Coccidia</i>	0	0	1	2	0	0	0	1	0	1	5 (5.3)
Larva of Nematodes	2	3	2	3	1	2	3	3	2	3	24 (25.8)
Egg of <i>Taenia</i>	0	0	0	1	0	0	0	1	0	1	3 (3.2)
Egg of <i>Hymenolepis nana</i>	0	0	0	0	0	0	0	1	0	1	2 (2.1)
Egg of <i>Dicrocoelium</i>	0	0	0	1	0	0	0	1	1	1	4 (4.3)
Egg of <i>Trichostrongylus</i>	1	0	1	2	0	1	0	1	0	0	6 (6.4)
Egg of <i>Fasciola hepatica</i>	0	1	0	1	0	1	0	1	0	0	4 (4.3)
Egg of <i>Toxocara</i>	0	0	1	0	0	0	1	0	0	1	3 (3.2)
Total, No. (%)	6 (6.4)	7 (7.5)	13 (13.9)	16 (17.2)	3 (3.2)	5 (5.3)	8 (8.6)	13 (13.9)	8 (8.6)	14 (15)	93 (100)

**Figure 1.** Frequency of Contamination in Vegetables

All of the farmers in north area of Iran are using animal fertilizer to improve the soil properties and their products. Field animals such as sheep (which are the main hosts of these parasites) live in this environment and this can result in human infections. For these reasons, it seems necessary to prevent animal entrance into the vegetables farm lands. In the current study, *Dicrocoelium* eggs compromised 4.3% of all worm eggs. This amount of contamination is reported in other cities of Iran such as Qazvin, Esfahan and Yazd (11, 16, 18). According to the high

prevalence of this Termatoda in animals vs. humans, we can conclude this sort of contamination is due to animal fertilizers. This parasite is a zoonotic agent and in desired circumstances, it can cause problems for human health. While according to the transmission way of this parasite, it seems that vegetables are not good transmitting vectors of infections but they have an important role in parasitic life cycle. *Taenia* eggs were seen in about 3.2% cases of this study which is also reported in other studies at different prevalence (13, 16). Because of vegetable contamination by *Taenia* eggs, the human infection and Hydatidosis in this way is probable. Therefore, vegetable wash before intake is very important to prevent human infections.

*Fasciola hepatica* eggs were seen in about 4.3% cases which is also reported by Malakoutian et al. in Kerman (17). Although this is an animal disease and is mostly reported in animals such as sheep, but it can cause pseudo-diseases (spurious infection) and complications in humans. The presence of these parasites and other organisms in animals is due to not using fences to prevent entrance of animals. Another parasite is *Toxocara* egg which is detected in 3.2% of cases. In a study by Uga et al. the presence of this egg was confirmed in Vietnam (8). *Toxocara* infection in humans, especially children resulted in ophthalmologic complications including retinal inflammation and ulcer and finally blindness. In our study, the lowest prevalence was for coriander contamination which is corresponded with Shahnazi et al. (16). In addi-

tion, in a study in Qazvin, the same results were obtained. One reason for this low contamination rate of coriander is the shape of this plant which has not too many layers and also has thin leaves. According to the high amount of parasitic contamination in vegetables, we can prevent human infections by educating people to wash and disinfect vegetables properly.

## Acknowledgements

There is no acknowledgement.

## Authors' Contribution

Concept(a), design(b), definition of intellectual content (c), literature search (d), microscopic observation (e), sample collection (f), data analysis (g), statistical analysis (h), manuscript preparation (i), manuscript edition (j). Abolghasem Siyadatpanah: b, d, e, i; Fatemeh Tabatabaei: c; Amir Emami Zeydi: b, f, i; Adel Spotin: h, j; Mehrdad Assadi: e, d; Sepideh Moradi: d; Ali Rostami: d; Vahid Fallah Omrani: a, h, i, j; Fatemeh Memari: d; Fateme Hajjaliani: d.

## Financial Disclosure

The authors have no proprietary interest in any aspect.

## Funding/Support

This research was supported by a grant from Department of Nursing, Faculty of Nursing and Midwifery, Mazandaran University of Medical Sciences, Amol, Iran.

## References

1. Markel EK, Voge M, John DJ. *Medical parasitology*. 9 ed. Philadelphia: WB. Saunders; 2006. p. 240-50.
2. Damen JG, Banwat EB, Egah DZ, Allanana JA. Parasitic contamination of vegetables in Jos, Nigeria. *Ann Afr Med*. 2007;6(3):115-8.
3. Gharavi MJ, Jahani MR, Rokni MB. Parasitic contamination of vegetables from farms and markets in Tehran. *Iran J Public Health*. 2002;31(3-4):83-6.
4. Choi DW, Lee S. Incidence of parasites found on vegetables collected from markets and vegetable gardens in Taegu area. *Korean J Parasitol*. 1972;10(1):44-51.
5. Bojar H, Klapac T. Contamination of soil with eggs of geohelminths in recreational areas in the Lublin region of Poland. *Ann Agric Environ Med*. 2012;19(2):267-70.
6. Klapac T. [Contamination of soil with geohelminth eggs on vegetable organic farms in the Lublin voivodeship, Poland]. *Wiad Parazytol*. 2009;55(4):405-9.
7. Coelho LM, Oliveira SM, Milman MH, Karasawa KA, Santos R. [Detection of transmissible forms of enteroparasites in water and vegetables consumed at schools in Sorocaba, Sao Paulo state, Brazil]. *Rev Soc Bras Med Trop*. 2001;34(5):479-82.
8. Uga S, Hoa NT, Noda S, Moji K, Cong L, Aoki Y, et al. Parasite egg contamination of vegetables from a suburban market in Hanoi, Vietnam. *Nepal Med Coll J*. 2009;11(2):75-8.
9. Erdog˘rul ˆ, Şener H. The contamination of various fruit and vegetable with *Enterobius vermicularis*, *Ascaris* eggs, *Entamoeba histolyca* cysts and *Giardia* cysts. *Food Control*. 2005;16(6):557-60.
10. Kozan E, Gonenc B, Sarimehmetoglu O, Aycicek H. Prevalence of helminth eggs on raw vegetables used for salads. *Food Control*. 2005;16(3):239-42.
11. Izadi S, Abedi S, Ahmadian S, Mahmoodi M. Study of the current parasitic contamination of the edible vegetables in Isfahan in order to identify preventive measures. *Sci J Kurdistan U Med Sci*. 2006;11(2):51-8.
12. Ranjbar Bahadˆri S, Estiri H, Kashefnejad M. Comparing the vegetable infection before and after washing, disinfecting and packaging vegetable processing farms in Sabzevar, Iran. 2. 2010;16(4):234-9.
13. Nazemi S, Raei M, Amiri M, Chaman R. Parasitic contamination of raw vegetables in Shahrour, Semnan. *Zahedan J Res Med Sci*. 2012;14(8):84-6.
14. Homayouni MM, Khalagi N. Parasitic infestation of consumed vegetables in Tehran in 1383. *J Army Univ Med Sci*. 2007;4(4):1053-6.
15. Al-Megrin WAI. Prevalence intestinal parasites in leafy vegetables in Riyadh, Saudi Arabia. *Int J Trop Med*. 2010;1(5):20-3.
16. Shahnazi M, Sharifi M, Kalantari Z, Allipour Heidari M, Agamirkarimi N. The study of consumed vegetable parasitic infections in Qazvin. *J Qazvin U Med Sci*. 2009;12:83.
17. Malakoutian M, Hosseini M, Bahrami H. Parasitic contamination of consuming vegetables in Kerman city Iran. *Hormozgan Med J*. 2009;13(1):55-62.
18. Dehgani Firozabadi A, Azizi M, Anvari MH. [A study of the contaminated vegetables in distribution centers of Yazd city]. *Toloo-EBehdasht J Health School*. 2004;2(1):11-5.