

CONFORMAL VECTOR FIELDS ON TANGENT BUNDLE WITH A SPECIAL LIFT FINSLER METRIC*

E. PEYGHAN^{1**}, A. RAZAVI² AND A. HEYDARI³

¹Department of Mathematics, Faculty of Science, University of Arak, Arak, I. R. of Iran

²Department of Mathematics and Computer Science, Amirkabir University, Tehran, I. R. of Iran

³Faculty of Science, Tarbiatmodares University, Tehran, I. R. of Iran

Emails: e-peyghan@araku.ac.ir, arazavi@aut.ac.ir, aheydari@modares.ac.ir

Abstract – On a Finsler manifold, we define conformal vector fields and their complete lifts and prove that in certain conditions they are homothetic.

Keywords – Conformal vector field, complete lift, finsler manifold, lift metric

1. PRELIMINARIES

Let (M, g) be a Riemannian manifold, a vector field V on M is called a *conformal vector field* if its local 1-parameter group of transformations is a local conformal transformation. It is well known that V is a conformal vector field on M if and only if there is a scalar function λ on M such that $L_V g = 2\lambda g$. When λ is a constant, V is called *homothetic*, especially when $\lambda = 0$, V is a *killing vector field* or an *infinitesimal isometry* [1].

On a Finsler manifold (M, F) , let V be a vector field with the complete lift V^c , then V is called conformal vector field if there is a scalar function ρ on TM such that $L_{V^c} g = 2\rho g$, where $g = (g_{ij})$ is the corresponding fundamental Finsler tensor defined by $g_{ij}(x, y) = (\frac{1}{2}F^2)_{y^i y^j}(x, y)$.

Let TM be the tangent space with a canonical coordinate system (x^i, y^i) , then the vertical tangent bundle of $TM_0 = TM \setminus \{0\}$ is defined by

$$VTM = span\{\frac{\partial}{\partial y^1}, \dots, \frac{\partial}{\partial y^n}\}.$$

A non-linear connection on TM_0 is a complementary distribution HTM defined by

$$HTM = span\{\frac{\delta}{\delta x^1}, \dots, \frac{\delta}{\delta x^n}\},$$

where $\frac{\delta}{\delta x^i} = \frac{\partial}{\partial x^i} - N_i^j \frac{\partial}{\partial y^j}$, and N_i^j are the connection coefficients. HTM is a vector bundle completely determined by the smooth functions $N_i^j(x, y)$ on TM [2, 3]. Moreover, we have

$$TTM_0 = VTM \oplus HTM \quad (1)$$

*Received by the editor April 14, 2007 and in final revised form August 28, 2007

**Corresponding author

Let ∇ be a linear connection on VTM , then (HTM, ∇) is called a *Finsler connection* on M . Indeed, a Finsler connection is a triad (N, F, C) where $N(N_j^i)$ is a nonlinear, $F(F_j^i)$ is the horizontal part and $C(C_j^i)$ is the vertical part of this connection. Now let (M, F) be Finsler manifold then a Finsler connection is called a *metric Finsler connection* if g is parallel with respect to ∇ . According to the Miron framework this means g is both horizontally and vertically a metric [4, 5, 6]. The *Cartan connection* is a metric Finsler connection for which the deflection, horizontal, and vertical torsion tensor fields vanish.

The *curvature tensor* of a metric Finsler connection is defined by

$$R(X, Y) = [\nabla_X, \nabla_Y] - \nabla_{[X, Y]}$$

where $X, Y \in \mathcal{X}(TM)$.

They are called horizontal or vertical according to the choice of X and Y in HTM or VTM . Then we have [5]

$$R_{kj}^h = \delta_i F_{kj}^h - \delta_j F_{ki}^h + F_{kj}^m F_{mi}^h - F_{ki}^m F_{mj}^h + C_{km}^h R_{ji}^m,$$

$R_{ij}^h = \delta_j N_i^h - \delta_i N_j^h$, where we have put $\partial_i = \frac{\partial}{\partial x_i}$, $\partial_{\bar{i}} = \frac{\partial}{\partial y_i}$, $\delta_i = \partial_i - N_i^m \partial_{\bar{m}}$. When ∇ is a Cartan connection then $N_i^h = y^m F_{mi}^h$.

Proposition 1. [4] Let M be an n -dimensional Finsler manifold with a Cartan connection, then we have the following equations:

- (1) $F_{ij}^h = \frac{1}{2} g^{hm} (\delta_i g_{mj} + \delta_j g_{im} - \delta_m g_{ij})$;
- (2) $C_{ijk} = \frac{1}{2} \dot{\partial}_k g_{ij}$ where $C_{ijk} = C_{ik}^m g_{jm}$;
- (3) $y^m C_{mij} = 0$;
- (4) $R_{ji}^h = y^m R_{mij}^h$.

The Cartan horizontal and vertical covariant derivative of a tensor field of type (1,2) are locally as follows:

$$\begin{aligned} \nabla_j T_{ki}^h &:= T_{k|j}^h = \delta_j T_{ki}^h + F_{mj}^h T_{ki}^m - F_{kj}^m T_{mi}^h - F_{ij}^m T_{km}^h; \\ \nabla_{\bar{j}} T_{ki}^h &:= T_{k|\bar{j}}^h = \partial_{\bar{j}} T_{ki}^h + C_{mj}^h T_{ki}^m - C_{kj}^m T_{mi}^h - C_{ij}^m T_{km}^h. \end{aligned} \quad (2)$$

2. LIFT METRICS AND CONFORMAL VECTOR FIELDS

a) Complete Lift Vector Fields and Lie Derivative

Let $V = v^i \partial_i$ be a vector field on M . Then V induces an infinitesimal point transformation on M . This is naturally extended to a point transformation of the tangent bundle TM which is called *extended point transformation*. Let V be a vector field on M and $\{\varphi_t\}$ the local 1-parameter group of M generated by V . Let $\tilde{\varphi}_t$ be the extended point transformation of φ_t , then $\{\tilde{\varphi}_t\}$ induces a vector field V^c on TM which is called the complete lift of V [7, 8].

It can be shown that the extended point transformation is a transformation induced by the complete lift vector field of V , $V^c = v^i \delta_i + y^j \nabla_j v^i \partial_{\bar{i}}$ with respect to the decomposition (1), where ∇ is a linear connection.

The Lie derivation of an arbitrary tensor, T_i^k , is given locally by [9]:

$$L_V T_i^k = v^a \nabla_a T_i^k + v^a \nabla_a v^b \nabla_b T_i^k - T_i^a \nabla_a v^k + T_a^k \nabla_i v^a$$

or equivalently,

$$L_V T_i^k = v^a \partial_a T_i^k + y^a \partial_a v^b \partial_b T_i^k - T_i^a \partial_a v^k + T_a^k \partial_i v^a.$$

So we have

$$L_V y^i = v^a \partial_a y^i + y^a \partial_a v^b \partial_b y^i - y^a \partial_a v^i = y^a \partial_a v^i - y^a \partial_a v^i = 0, \quad (3)$$

$$L_V g_{ij} = v^a \partial_a g_{ij} + y^a \partial_a v^b \partial_b g_{ij} + g_{aj} \partial_i v^a + g_{ia} \partial_j v^a. \quad (4)$$

where ∇ is a linear connection.

In Finsler geometry, L_V is replaced by $L_{\tilde{V}}$, where \tilde{V} is the lift of V . We also have this interchanging formula between Cartan covariant derivatives and Lie derivatives.

$$\nabla_k L_V g_{ij} - L_V \nabla_k g_{ij} = g_{aj} L_V F_{ik}^a + g_{ai} L_V F_{jk}^a. \quad (5)$$

b) A Lift Metric on Tangent Bundle

V. Oproiu introduced a family of Riemannian metrics on the tangent space of Riemannian manifolds and considered locally symmetric, Kählerian and anti-Hermitian conditions with these metrics [10-12]. Then Abbassi-Sarih proved in [13] that the Oproiu metrics form a particular subclass of the so-called *g-natural metrics* on the tangent space [14, 15]. Also in [16], Boeckx-Vanhecke obtained an almost contact metric on the unit tangent space.

In this section we consider a new Riemannian metric on the tangent space, and in the next section obtain some conditions which reduce the conformal vector fields to be homothetic.

Let (M, F) be a Finsler manifold, define a tensor field G on TM by

$$G(x, y) = \alpha h_{ij}(x, y) dx^i dx^j + 2\beta h_{ij}(x, y) dx^i \delta y^j + \gamma h_{ij}(x, y) \delta y^i \delta y^j$$

where α, β and γ are real numbers and $h_{ij}(x, y)$ are components of a generalized Lagrange metric [6, 17]. It is clear that G is nonsingular if $\alpha\gamma - \beta^2 \neq 0$ and positive definite if $\alpha\gamma - \beta^2 > 0$, defining, respectively, a *pseudo-Riemannian* or *Riemannian lift metrics* on $T(M)$.

We are going to consider the metric G with $h_{ij}(x, y)$ of the following special deformation of $g_{ij}(x)$

$$h_{ij}(x, y) = a(F^2) g_{ij}(x, y),$$

where $y_i = g_{ij}(x, y) y^j$ and $a: Im(F^2) \subseteq R_+ \rightarrow R_+$ with $a > 0$. For shortness we set $g_1 = h_{ij} dx^i dx^j$, $g_2 = 2h_{ij} dx^i \delta y^j$ and $g_3 = h_{ij} \delta y^i \delta y^j$, therefore $G = \alpha g_1 + \beta g_2 + \gamma g_3$.

3. MAIN RESULTS

Analogous to the Riemannian geometry, by straightforward calculation we have the following results in Finsler geometry [18, 19].

Lemma 1. Let (M, F) be a Finsler manifold with Cartan connection, then we have

- (1) $[\delta_i, \delta_j] = R_{ij}^h \partial_h$;
- (2) $[\delta_i, \partial_j] = \partial_j N_i^h \partial_h$;

Winter 2008

Iranian Journal of Science & Technology, Trans. A, Volume 32, Number A1

$$(3) [\partial_{\bar{i}}, \partial_{\bar{j}}] = 0.$$

Lemma 2. Let (M, F) be a Finsler manifold with Cartan connection, then we have

- (1) $L_{V^c} \delta_i = -\partial_i v^h \delta_h - L_v N_i^h \partial_{\bar{h}};$
- (2) $L_{V^c} \partial_{\bar{i}} = -\partial_i v^h \partial_{\bar{h}};$
- (3) $L_{V^c} dx^h = \partial_m v^h dx^m;$
- (4) $L_{V^c} \delta y^h = L_v N_m^h dx^m + \partial_m v^h \delta y^m.$

Proof: First we give the proof of part (2). By a simple calculation, we have:

$$\begin{aligned} L_{V^c} \partial_{\bar{i}} &= [v^c, \partial_{\bar{i}}] \\ &= [v^h \delta_h + y^m v^h|_m \partial_{\bar{h}}, \partial_{\bar{i}}] \\ &= v^h [\delta_h, \partial_{\bar{i}}] - \partial_{\bar{i}} (v^h) \delta_h + y^m v^h|_m [\partial_{\bar{h}}, \partial_{\bar{i}}] - \partial_{\bar{i}} (y^m v^h|_m) \partial_{\bar{h}} \\ &= \partial_{\bar{i}} (v^h N_h^r - y^m v^r|_m) \partial_{\bar{r}} \\ &= -\partial_i v^r \partial_{\bar{r}}; \end{aligned}$$

The proof of part (1) is similar to (2).

Since $(dx^h, \delta y^h)$ is the dual basis of $(\delta_h, \partial_{\bar{h}})$, if we put

$$L_{V^c} \delta y^h = \alpha_m^h dx^m + \beta_m^h \delta y^m,$$

then we have

$$0 = L_{V^c} (\delta y^h (\delta_i)) = (L_{V^c} \delta y^h) \delta_i + \delta y^h (L_{V^c} \delta_i) = \alpha_i^h - L_{V^c} N_i^h,$$

and

$$0 = L_{V^c} (\delta y^h (\partial_{\bar{i}})) = (L_{V^c} \delta y^h) \partial_{\bar{i}} + \delta y^h (L_{V^c} \partial_{\bar{i}}) = \beta_i^h - \partial_i v^h.$$

Thus we get (4). In the same way as the proof of part (4), we can prove (3).

Lemma 3. Let (M, g) be a Finsler manifold with Cartan connection, then we have

- (1) $L_{V^c} g_1 = a(F^2)(2\varphi g_{ij} + L_v g_{ij}) dx^i dx^j;$
- (2) $L_{V^c} g_2 = 2a(F^2) g_{mi} (L_v N_j^m) dx^i dx^j + 2a(F^2)(2\varphi g_{ij} + L_v g_{ij}) \delta y^i \delta y^j;$
- (3) $L_{V^c} g_3 = 2a(F^2) g_{mi} L_v N_j^m dx^i \delta y^j + a(F^2)(2\varphi g_{ij} + L_v g_{ij}) \delta y^i \delta y^j.$

$$\text{where } \varphi = y^m v^h|_m y^h \frac{a'(F^2)}{a(F^2)}.$$

Proof: From the above lemma, we get

$$\begin{aligned} L_{V^c} g_1 &= L_{V^c} (h_{ij} dx^i dx^j) = V^c (a(F^2) g_{ij}) dx^i dx^j + 2a(F^2) g_{ij} (L_{V^c} dx^i) dx^j \\ &= ((v^h \delta_h + y^m v^h|_m \partial_{\bar{h}}) a(F^2)) g_{ij} + ((v^h \delta_h + y^m v^h|_m \partial_{\bar{h}}) g_{ij}) a(F^2) \\ &\quad + 2a(F^2) g_{ij} (\partial_r v^i dx^r) dx^j \\ &= 2a(F^2) \varphi g_{ij} dx^i dx^j + a(F^2) L_v g_{ij} dx^i dx^j. \end{aligned}$$

Thus we have (1). (2) and (3) are easily proof in the same way as the proof of (1).

Definition 1. Let X be a conformal vector field on TM with the associated function ρ . X is called quasi-inessential vector field if $\rho - \varphi$ is a function of (x^h) , namely there exists a function Ω of (x^h) such that $\rho = \Omega + \varphi$. If Ω is constant, then X is called quasi-homothetic vector field. Moreover, if $\Omega = 0$ then X is called quasi-isometry vector field on TM .

Remark: These classes of vector fields contain the classes of inessential, homothetic and isometry vector fields as special cases, respectively (for $\varphi = 0$). Hence, the forthcoming results hold for inessential, homothetic and isometry vector fields.

Theorem 1. Let (M, F) be a C^∞ connected Finsler manifold, TM its tangent bundle and G the Riemannian (or pseudo-Riemannian) metric on TM derived from g . Then every complete lift conformal vector field on TM is quasi-homothetic.

Proof: Let V be a vector field on M , V^c the complete lift vector field of V which is conformal, and let G be a pseudo-Riemannian metric on TM derived from g . We have by definition $L_{V^c}G = 2\rho G$. The Lie derivative of G gives

$$\begin{aligned} L_{V^c}G &= \alpha a(F^2)(2\varphi g_{ij} + L_V g_{ij})dx^i dx^j + 2\beta a(F^2)(2\varphi g_{ij} + L_V g_{ij})dx^i \delta y^j \\ &\quad + 2\beta a(F^2)g_{ai}L_V N^a_j dx^i dx^j + \gamma a(F^2)(2\varphi g_{ij} + L_V g_{ij})\delta y^i \delta y^j \\ &\quad + 2\gamma a(F^2)g_{aj}L_V N^a_i dx^i \delta y^j. \end{aligned} \quad (6)$$

So we have

$$\begin{aligned} L_{V^c}G &= a(F^2)[\alpha(2\varphi g_{ij} + L_V g_{ij}) + 2\beta g_{ai}(L_V N^a_j)]dx^i dx^j \\ &\quad + a(F^2)[2\beta(2\varphi g_{ij} + L_V g_{ij}) + 2\gamma g_{aj}(L_V N^a_i)]dx^i \delta y^j \\ &\quad + \gamma a(F^2)(2\varphi g_{ij} + L_V g_{ij})\delta y^i \delta y^j = 2\rho G. \end{aligned}$$

Comparing with the definition of G , we find

$$\alpha L_V g_{ij} + \beta(g_{ai}L_V N^a_j + g_{aj}L_V N^a_i) = 2\alpha \Omega g_{ij}; \quad (7)$$

$$\beta L_V g_{ij} + \gamma g_{aj}L_V N^a_i = 2\beta \Omega g_{ij}; \quad (8)$$

$$\gamma L_V g_{ij} = 2\gamma \Omega g_{ij}. \quad (9)$$

Where $\Omega = \rho - \varphi$.

I) If $\gamma \neq 0$, then from (9) we have

$$L_V g_{ij} = 2\Omega g_{ij}$$

and from (8) we have

$$L_V N^a_i = 0.$$

Using this and $N^h_i = y^m F^h_{mi}$ we get

$$0 = L_V N_i^h = L_V (y^m F_{mi}^h) = y^m L_V F_{mi}^h, \quad (10)$$

where the last equality follows from equation (3).

II) If $\gamma = 0$, since $\alpha\gamma - \beta^2 \neq 0$ we have $\beta \neq 0$. From (8) we get

$$L_V g_{ij} = 2\Omega g_{ij}$$

and from (7) we have

$$g_{ai} L_V N_j^a + g_{aj} L_V N_i^a = 0.$$

Using this, equation (3) and $N_i^h = y^m F_{mi}^h$, we have

$$y^m (g_{ai} L_V F_{mj}^a + g_{aj} L_V F_{mi}^a) = 0. \quad (11)$$

In each case I and II we have

$$L_V g_{ij} = 2\Omega g_{ij} \quad (12)$$

or from equation (4)

$$v^a \partial_a g_{ij} + g_{aj} \partial_i v^a + g_{ia} \partial_j v^a + y^a \partial_a v^b \partial_b g_{ij} = 2\Omega g_{ij}.$$

Applying $\partial_{\bar{k}}$ to both sides of the above equation, we find that

$$2v^a \partial_a C_{ijk} + 2C_{ajk} \partial_i v^a + 2C_{iak} \partial_j v^a + 2\partial_k v^a C_{ija} + 2y^a \partial_a v^b \partial_b C_{ijb} = 2g_{ij} \partial_{\bar{k}} \Omega + 4\Omega C_{ijk}.$$

By using $y^i C_{ijk} = 0$, we obtain $\partial_{\bar{k}} \Omega = 0$. Therefore Ω is a function of x alone. From (5) we have

$$y^k (\nabla_k L_V g_{ij} - L_V \nabla_k g_{ij}) = y^k (g_{aj} L_V F_{ik}^a + g_{ai} L_V F_{jk}^a).$$

By using (10), (11) and (12) in each case I and II we find that

$$y^k \nabla_k \Omega = 0.$$

Since Ω is a function of x alone, we obtain $\partial_i \Omega = 0$. This, together with the connectedness of M , shows that Ω is constant.

Note: In a special case when $a'(F^2) = 0$ e.g. $a(t) = (t - F^2)^2 + 1$ follows from lemma 3, that $\varphi = 0$ and hence $L_{V^c} G = 2\rho G$, where ρ depends on x only. Therefore we have:

Corollary 1. Let (M, F) be a C^∞ connected Finsler manifold, TM its tangent bundle and G the Riemannian (or pseudo-Riemannian) metric on TM derived from g with $a'(F^2) = 0$. Then every complete lift conformal vector field on TM is homothetic.

REFERENCES

1. Bejancu, A. (1990). *Finsler geometry and applications*. England: Ellis Horwood Limited Publication.
2. Yano, K. & Ishihara, S. (1973). *Tangent and Cotangent Bundles*. New York, Marcel Dekker.
3. Yano, K. & Kobayashi, S. (1996). Prolongations of tensor fields and connection to tangent bundle I, *General theory. J. Math. Soc. Japan.*, 18, 194-210.
4. Miron, R. (1981). Introduction to the theory of Finsler spaces. *Proc. Nat. Sem. Brasov, On Finsler spaces*.

5. Miron, R. & Anastasiei, M. (1981). *Vector bundles and Lagrange spaces with application to Relativity*. Romania: Geometry Balkan Press.
6. Miron, R. & Anastasiei, M. (1994). *The geometry of Lagrange spaces: Theory and Applications*. Kluwer Academic Publishers. FTPH.
7. Akbar-Zadeh, H. (1979). Transformations infinitesimals conformes des varietes finsleriennes compactes. *Ann. Polon. Math.*, 36, 213-229.
8. Bidabad, B. (2006). Conformal vector fields on tangent bundle of a Finsler manifolds. *Balkan Journal of Geometry and Its Applications*, 2, 28-35.
9. Yano, K. (1957). *The theory of Lie Derivatives and Its Applications*. North Holland.
10. Oproiu, V. (1999). A locally symmetric Kähler Einstein structure on the tangent bundle of a space form. *Beiträge Zur Algebra und Geometrie/Contributions to Algebra and Geometry*, 40, 363-372.
11. Oproiu, V. (2001). A Kähler Einstein structure on the tangent bundle of a space form. *Int. J. Math. Sci.*, 25, 183-195.
12. Oproiu, V. & Papaghiuc, N. (2004). Some classes of almost anti-Hermitian structures on the tangent bundle. *Mediterr. J. Math.*, 1, 269-282.
13. Abbassi, M. T. K. & Sarih, M. (2005). On some Hereditary properties of Riemannian g-natural metrics on tangent bundles of Riemannian manifolds. *Differential Geometry and Its Applications*, 22, 19-47.
14. Abbassi, M. T. K. (2004). Note on the classification theorems of g-natural metrics on the tangent bundle of a Riemannian manifold (M, g) . *Commnet. Math. Univ. Carolinae*, 45(4), 591-596.
15. Kowalski, O. & Sekizawa, M. (1988). Natural transformations of Riemannian metrics on manifolds to metrics on tangent bundle- a classification-. *Bull. Tokyo Gakuei Univ.*, 40(4), 1-29.
16. Boeckx, E. & Vanhecke, L. (2000). Harmonic and minimal vector fields on tangent and unit tangent bundles. *Differential Geometry and Its Applications*, 13, 77-93.
17. Peyghan, E. & Heydari, A. (2008). Conformal vector fields on tangent bundle of a Riemannian manifold. *Journal of Mathematical Analysis and Applications*, 347(1), 136-142.
18. Yamauchi, K. (1996). On infinitesimal conformal transformations of the tangent bundles over Riemannian manifolds. *Ann. Rep. Asahikawa. Med. Coll.*, 17, 1-7.
19. Kobayashi, S. (1995). A theorem on the affine transformation group of a Riemannian manifold. *Nagoya Math. J.*, 9, 39-41.