

THE STRUCTURE OF DERIVATIONS FROM A FULL MATRIX ALGEBRA INTO ITS DUAL*

R. ALIZADEH¹ AND G. H. ESSLAMZADEH^{2**}

¹Department of Mathematics, Shahed University, P. O. Box 18151-159, Tehran, I. R. of Iran
 Email: alizadeh@shahed.ac.ir

²Department of Mathematics, Shiraz University, Shiraz 71454, I. R. of Iran
 Email: esslamz@shirazu.ac.ir

Abstract – Let A be a unital algebra over a field of characteristic zero. We show that every derivation from $M_n(A)$ into its dual $M_n(A)^*$ is the sum of an inner derivation and a derivation induced by a derivation from A into A^* .

Keywords – Derivation, full matrix algebra, dual space

1. INTRODUCTION

Throughout A is a unital algebra over a field of characteristic zero and M is an A -bimodule. We denote the full matrix algebra of $n \times n$ matrices over A with the usual operations by $M_n(A)$. E_{ij} s, $1 \leq i, j \leq n$ are also usual matrix units in $M_n(A)$. For all $x \in A$ we display matrix whose (i, j) th entry is x and zero elsewhere, by $x \otimes E_{ij}$. The dual A^* of A is the set of all linear maps from A into its field. We denote the action of an element $g \in A^*$ on an element $a \in A$ with $\langle g, a \rangle$. Also, A^* is an A -bimodule with the following module operations:

$$\langle a.f, b \rangle = \langle f, ba \rangle \text{ and } \langle f.a, b \rangle = \langle f, ab \rangle, \forall a, b \in A, f \in A^*.$$

A derivation $D: A \rightarrow M$ is a linear map which satisfies the identity $D(ab) = D(a)b + aD(b)$ $a, b \in A$. We say that D is inner if there exists $m \in M$ such that $D(a) = am - ma$ for all $a \in A$. Every derivation $D: A \rightarrow M$ induces a derivation $\bar{D}: M_n(A) \rightarrow M_n(M)$ by $\bar{D}((a_{ij})) = (D(a_{ij}))$.

Benkart and Osborn [1] characterized derivations of $M_n(A)$, where A is a unital nonassociative algebra with $\text{char}(A) \neq 2, 3$ and $n > 2$. They showed that every derivation of $M_n(A)$ is a sum of an inner derivation generated by a matrix with entries in the nucleus N of A , and a derivation induced by a derivation of A . A similar result for full matrix rings was proved in [2]. The case of the centers of upper triangular matrix rings over simple algebras which are finite dimensional modulo, was discussed in [3]. Coelho and Milies in [4] proved a similar result for upper triangular matrices over an arbitrary ring with identity. Jondrup gave a new proof of the latter result in [5].

In this article we prove an analog of the above mentioned result for derivations from $M_n(A)$ into its dual $M_n(A)^*$. In other words, we prove the following theorem.

Theorem: Every derivation from $M_n(A)$ into its dual $M_n(A)^*$ is the sum of an inner derivation and a derivation induced by a derivation from A into A^* .

*Received by the editor May 27, 2007 and in final revised form August 28, 2007

**Corresponding author

2. MAIN RESULT

Let $g \in M_n(A)^*$ and $1 \leq i, j \leq n$. Define $g_{ij} \in A^*$ by $\langle g_{ij}, a \rangle = \langle g, a \otimes E_{ij} \rangle$. We can identify $M_n(A)^*$ with $M_n(A^*)$ via the map

$$\phi: M_n(A)^* \rightarrow M_n(A^*), \quad g \mapsto (g_{ij}).$$

From now on we display $g \in M_n(A)^*$ by (g_{ij}) . For all $(a_{ij}) \in M_n(A)$ we have

$$\langle g, (a_{ij}) \rangle = \langle (g_{ij}), (a_{ij}) \rangle = \sum_{i,j=1}^n \langle g_{ij}, a_{ij} \rangle.$$

Theorem: Every derivation from $M_n(A)$ into its dual $M_n(A)^*$ is the sum of an inner derivation and a derivation induced by a derivation from A into A^* .

Proof: Let $(f_{ij}) \in M_n(A)^*$, $(a_{ij}) \in M_n(A)$ and $a \in A$. Then we have

$$\begin{aligned} \langle (f_{ij})(a_{ij}), a \otimes E_{kl} \rangle &= \langle (f_{ij}), (a_{ij})[a \otimes E_{kl}] \rangle = \langle (f_{ij}), \sum_{s=1}^n [a_{sk} a \otimes E_{sl}] \rangle \\ &= \sum_{s=1}^n \langle f_{sl}, a_{sk} a \rangle = \langle \sum_{s=1}^n f_{sl} a_{sk}, a \rangle. \end{aligned}$$

Thus

$$[(f_{ij})(a_{ij})]_{kl} = \sum_{s=1}^n f_{sl} a_{sk}. \tag{1}$$

Similarly, we have:

$$[(a_{ij})(f_{ij})]_{kl} = \sum_{s=1}^n a_{is} f_{ks}. \tag{2}$$

Suppose $D : M_n(A) \rightarrow M_n(A)^*$ is a derivation. Define

$$D_{ij}^{kl} : A \rightarrow A^*, \quad D_{ij}^{kl}(a) = [D(a \otimes E_{ij})]_{kl}, \quad 1 \leq i, j, k, l \leq n. \tag{3}$$

From (1) and (2) we conclude that for every $a, b \in A$ and every positive integer $m \leq n$ the following equality holds:

$$[D(a \otimes E_{im})][b \otimes E_{mj}]_{kl} = \sum_{r=1}^n D_{im}^{rl}(a) b \delta_{mr} \delta_{jk} = D_{im}^{ml}(a) b \delta_{jk} \tag{4}$$

Where δ is the Kronecker's delta. Similarly, we have

$$[[a \otimes E_{im}]D(b \otimes E_{mj})]_{kl} = \sum_{r=1}^n a \delta_{il} \delta_{mr} D_{mj}^{kr}(b) = a \delta_{il} D_{mj}^{km}(b). \tag{5}$$

From (4) and (5) we conclude that

$$D_{ij}^{kl}(ab) = D_{im}^{ml}(a) b \delta_{jk} + a \delta_{il} D_{mj}^{km}(b). \tag{6}$$

Therefore D_{ii}^{ii} is a derivation from A to A^* , for $1 \leq i \leq n$. Using (6) we have

$$D_{ij}^{jl}(a) = D_{ii}^{il}(1)a, D_{ij}^{ki}(a) = aD_{jj}^{kj}(1) \quad a \in A, i \neq l, k \neq j. \quad (7)$$

Again, using (6) for every $0 \leq i, j, l \leq n$, the following equalities hold.

$$D_{jj}^{jj}(a) = D_{ji}^{ij}(1)a + D_{ij}^{ji}(a). \quad (8)$$

$$D_{ij}^{ji}(a) = D_{il}^{li}(1)a + D_{lj}^{jl}(a). \quad (9)$$

$$D_{lj}^{jl}(a) = D_{ll}^{ll}(a) + aD_{lj}^{jl}(1). \quad (10)$$

$$D_{ji}^{ij}(a) = aD_{ji}^{ij}(1) + D_{jj}^{jj}(a). \quad (11)$$

From (8) we have

$$D_{ji}^{ij}(1) = -D_{ij}^{ji}(1). \quad (12)$$

Also, from (9), (10) and (12) we have

$$D_{ij}^{ji}(a) = D_{il}^{li}(1)a - aD_{lj}^{jl}(1) + D_{ll}^{ll}(a). \quad (13)$$

Using (8) and (11) we conclude that

$$D_{ij}^{ji}(a) = D_{ji}^{ij}(a) - D_{ji}^{ij}(1)a - aD_{jj}^{jj}(1) \quad (14)$$

In addition, using (6), (7) and (14) we obtain

$$\begin{aligned} [D((a_{rs}))]_{ij} &= \sum_{k,l=1}^n D_{kl}^{ij}(a_{kl}) \\ &= \sum_{k=1}^n D_{ki}^{ij}(1)a_{ki} + \sum_{l=1}^n a_{jl}D_{jl}^{ij}(1) - D_{ji}^{ij}(1)a_{ji} - a_{ji}D_{ji}^{ij}(1) + D_{ji}^{ij}(a_{ji}) \\ &= \sum_{k=1}^n D_{kk}^{kj}(1)a_{ki} + \sum_{k=1}^n a_{jk}D_{kk}^{ik}(1) + D_{ij}^{ji}(a_{ji}). \end{aligned} \quad (15)$$

On the other hand, we have

$$0 = [D(E_{kk}E_{ii})]_{ik} = \sum_{s=1}^n D_{kk}^{sk}(1)\delta_{si} + \sum_{s=1}^n \delta_{ks}D_{ii}^{is}(1) = D_{kk}^{ik}(1) + D_{ii}^{ik}(1). \quad (16)$$

Therefore,

$$D_{kk}^{ik}(1) = -D_{ii}^{ik}(1). \quad (17)$$

Now, for $1 \leq k, j \leq n$ define $D_{kj} = D_{kk}^{kj}$. From (15) and (17) we conclude that

$$\begin{aligned} [(D(a_{rs}))]_{ij} &= \sum_{k=1}^n D_{kj}(1)a_{ki} - \sum_{k=1}^n a_{jk}D_{ik}(1) + D_{ij}^{ji}(a_{ji}) \\ &= [(D_{rs}(1))(a_{rs}) - (a_{rs})(D_{rs}(1))]_{ij} + D_{ij}^{ji}(a_{ji}). \end{aligned} \quad (18)$$

Using (13) and (18) for every $1 \leq l \leq n$ we obtain

$$D((a_{ij})) = [(D_{ij}(1) + \text{diag}(D_{11}^{l1}(1), \dots, D_{nl}^{ln}(1)))(a_{ij}) \\ - (a_{ij})[(D_{ij}(1) + \text{diag}(D_{11}^{l1}(1), \dots, D_{nl}^{ln}(1)))] + \overline{D_{il}^{ll}}((a_{ij}))].$$

REFERENCES

1. Benkart, G. M. & Osborn, J. M. (1981). Derivations and automorphisms of non associative matrix algebras. *Trans. Amer. Math. Soc.* 263, 411-430.
2. Mathis, D. (1982). Differential polynomial rings and Morita equivalence. *Comm. Alg.* 10, 2001-2017.
3. Jöndrup, S. (1987). Automorphisms of upper triangular matrix rings. *Arch. Math.* 49, 497-502.
4. Coelho, S. P. & Milies, C. P. (1993). Derivations of upper triangular matrix rings. *Linear Alg. Appl.* 187, 263-267.
5. Jöndrup, S. (1995). Automorphisms and derivations of upper triangular matrices. *Linear Alg. Appl.* 22, 205-215.