

"Research Note"

ON THE SOLVABILITY OF SOME OPERATOR-DIFFERENTIAL EQUATIONS IN COMPLEX DOMAIN*

A. M. AKHMEDOV** AND Y. Y. MUSTAFAYEVA

Baku State University, Department of Mechanics and Mathematics,
 Z. Khalilov st., 23, P. O. Box Az-1148, Baku, Azerbaijan
 Emails: ali_akhmedov@hotmail.com; helenmust@rambler.ru

Abstract – In the present paper an operator-differential equation of second order in complex domain is considered when the coefficients have singularity of pole type at the point $z=0$. A theorem of existence of the solution of the equation is proved and the spectral property of the solution is separately investigated when the coefficients are spectral operators.

Keywords – Banach algebra, operator-differential equation, spectral operators, Boolean algebra

1. SOLVABILITY OF AN OPERATOR-DIFFERENTIAL EQUATION OF SECOND ORDER

Let $L(H)$ be a Banach algebra of linear bounded operators, acting in H , where H is a Hilbert space. Consider the equation

$$\frac{d^2U}{dz^2} = \frac{1}{z} \left(\sum_{k=0}^{\infty} B_k z^k \right) \frac{dU}{dz} + \frac{1}{z^2} \left(\sum_{k=0}^{\infty} A_k z^k \right) U, \quad (1)$$

where z is complex variable, $A_k, B_k \in L(H)$ ($k = 0, 1, 2, \dots$) and the series $\sum_{k=0}^{\infty} A_k z^k$ and $\sum_{k=0}^{\infty} B_k z^k$ are absolutely convergent in the circles $|z| < \rho_1$ and $|z| < \rho_2$, respectively. Let $\rho = \min(\rho_1, \rho_2)$ and later we will consider the problem in the circle $|z| < \rho$.

We seek the solution of (1) in the form:

$$U(z) = \left(\sum_{m=0}^{\infty} U_m z^m \right) z^R, \quad (2)$$

where the operators U_m and R will be determined later

Having calculated derivatives $\frac{dU}{dz}$ and $\frac{d^2U}{dz^2}$, putting them in (1) and applying the abstract analogy of the Frobenius method, we can write out formulas for coefficients U_m :

$$U_0 (R^2 - R) - B_0 U_0 R - A_0 U_0 = 0, \quad (3)$$

$$U_m [R^2 + 2mR + m(m-1)I] - mB_0 U_m - mB_0 U_m - B_0 U_m R - A_0 U_m = F_m, \quad m = 1, 2, \dots, \quad (4)$$

*Received by the editor February 20, 2006 and in final revised form October 21, 2008

**Corresponding author

where

$$F_m = \sum_{\substack{k+p=m-1 \\ p \neq m-1}} (p+1)B_k U_{p+1} + \sum_{\substack{k+p=m \\ p \neq m}} B_k U_p R + \sum_{\substack{k+p=m \\ p \neq m}} A_k U_p. \quad (5)$$

Let us choose U_0 bounded and such that U_0^{-1} exists and is bounded too. Let the operators A_0 , B_0 and U_0 be commutative. If the operator $B_0^2 + 2B_0 + I + A_0$ is a spectral operator of scalar type, then from equation (4) for the desired operator R we obtain

$$R^2 - (B_0 + I)R - A_0 = 0, \quad (6)$$

and therefore

$$R = f(B_0) = \frac{B_0 + I + (B_0^2 + 2B_0 + I + A_0)^{\frac{1}{2}}}{2}, \quad (7)$$

$$\text{or } R = g(B_0) = \frac{B_0 + I - (B_0^2 + 2B_0 + I + A_0)^{\frac{1}{2}}}{2}. \quad (8)$$

Let $A_0 = B_0^k$, where k is some nonnegative integer numbers, then by theorem 3.1 from ([1], p.37) we obtain that for the solvability of equation (4) there must hold the next condition:

$$P(\lambda, \mu) = \mu^2 + (2m-1)\mu - \lambda\mu - \lambda^k - m\lambda + (m-1)m \neq 0 \quad (9)$$

for $\forall (\lambda, \mu) \in \sigma(B_0) \times \sigma(R)$ where $\sigma(B_0)$ and $\sigma(R)$ are spectrums of operators B_0 and R respectively.

Then the solution of equation (4) is determined by the formula

$$U_m = \frac{1}{4\pi^2} \int_{\Gamma_{B_0}} \int_{\Gamma_R} \frac{(B_0 - \lambda I)^{-1} F_m (R - \mu I)^{-1}}{P(\lambda, \mu)} d\mu d\lambda, \quad (10)$$

and Γ_{B_0}, Γ_R are piece-smooth contours, surrounding the spectrums of operators B_0 and R , respectively.

If R is defined by (7) then we have

$$(R - \mu I)^{-1} = (f(B_0) - \mu I)^{-1} = \frac{1}{2\pi} \int_{\Gamma_{B_0}} \frac{(B_0 - \nu I)^{-1} d\nu}{(f(\nu) - \mu)}. \quad (11)$$

Putting (11) into (10) we obtain:

$$U_m = \frac{1}{4\pi^2} \int_{\Gamma_{B_0}} \int_{\Gamma_R} \frac{(B_0 - \lambda I)^{-1} F_m (R - \mu I)^{-1}}{P(\lambda, \mu)} d\mu d\lambda = \frac{1}{4\pi^2} \int_{\Gamma_0} \int_{\Gamma_0} \frac{(B_0 - \lambda I)^{-1} F_m (B_0 - \nu I)^{-1}}{P(\lambda, f(\nu))} d\nu d\lambda. \quad (12)$$

Therefore, the solution of equation (4) is defined by formula (12), and the condition (9) now looks so:

$$P(\lambda, f(\nu)) = f(\nu)^2 + (2m-1)f(\nu) - \lambda f(\nu) - \lambda^k - m\lambda + (m-1)m \neq 0 \quad (13)$$

for arbitrary $(\lambda, \nu) \in \sigma(B_0) \times \sigma(B_0)$.

It is clear that $P(\lambda, f(\nu)) = O(m^2)$. Using this we obtain: $\|U_m\| \leq \frac{c}{m^2} \|F_m\|$. It is not difficult to prove that for any ρ_1 , such that $0 < \rho_1 < \rho$, and for any $m \geq 0$

$$\|U_m\|\rho_1^m \leq const.$$

Then for any $\rho_2: \rho_1 < \rho_2 < \rho$ we have: $\sum_{n=1}^{\infty} \|U_n\|\rho_1^n = \sum_{n=1}^{\infty} \|U_n\|\rho_2^n \left(\frac{\rho_1}{\rho_2}\right)^n \leq const \sum_{n=0}^{\infty} \left(\frac{\rho_1}{\rho_2}\right)^n < \infty$. Hence $\sum_{n=1}^{\infty} U_n \rho_1^n$ is convergent for $\forall \rho_1: 0 < \rho_1 < \rho$, whence follows the existence of solution (2) of equation (1).

For the existence of solution (2) of equation (1) by the theorem 3.1 ([1], p.37), in the case of (8) the following condition must be satisfied.

$$P(\lambda, g(v)) = g(v)^2 + (2m - 1)g(v) - \lambda g(v) - \lambda^k - m\lambda + (m - 1)m \neq 0. \quad (14)$$

Let us point out that according to condition $A_0 = B_0^k$, for the commutative property of operators A_0, B_0 and U_0 it suffices to require the commutative property of operators B_0 and U_0 . So we come to the theorem of existence.

Theorem 1. Let the next conditions be satisfied:

- a) operator $B_0^2 + 2B_0 + I + A_0$ is an operator of scalar type; b) $A_0 = B_0^k$, where k is nonnegative integer;
- c) operators B_0 and U_0 are commutative and operator U_0^{-1} exists and is bounded; d) for $f(v)$, defined in (7), condition (13) holds (or for $g(v)$, defined in (8), condition (14) holds). Then there exists the solution of equation (1) in the form $U(z) = \left(\sum_{n=0}^{\infty} U_n z^n\right) z^R$, where operator R is defined by (7) (by (8)); at that series $\sum_{n=0}^{\infty} U_n z^n$ is absolutely convergent in the circle $|z| < \rho$, $\rho = \min(\rho_1, \rho_2)$.

2. THE CASE OF SPECTRAL COEFFICIENTS

Suppose that $A_i, B_j, i, j = 0, 1, 2, \dots$, are mutually commutative spectral operators.

The next relation is known for the resolvent of a spectral operator B_0 ([2], XV.5.2):

$$(B_0 - \lambda I)^{-1} = \sum_{n_1=0}^{\infty} N^{n_1} \int_{\sigma(B_0)} \frac{E(d\theta)}{(\lambda - \theta)^{n_1+1}}, \quad (15)$$

where N is a quasinilpotent part, E is a resolution of the identity operator B_0 .

Let's put (15) in (12):

$$U_m = \sum_{n_1+n_2=0}^{\infty} N^{n_1+n_2} \int_{\Gamma_{B_0}} \int_{\Gamma_{B_0}} \frac{1}{P(\lambda, f(v))} \int_{\sigma(B_0)} \frac{E(d\theta)}{(\lambda - \theta)^{n_1+1}} \int_{\sigma(B_0)} \frac{E(d\eta)}{(v - \eta)^{n_2+1}} d\lambda d\nu F_m. \quad (16)$$

Denoting $P_1(\lambda, v) = P(\lambda, f(v))$, by Fubini's theorem we obtain

$$U_m = \left(\sum_{n_1+n_2=0}^{\infty} \frac{N^{n_1+n_2}}{n_1!n_2!} \int_{\sigma(B_0)} \int_{\sigma(B_0)} \frac{\partial^{n_1+n_2}}{\partial \theta^{n_1} \partial \eta^{n_2}} \left(\frac{1}{P_1(\theta, \eta)} \right) E(d\theta) E(d\eta) \right) F_m. \quad (17)$$

Let's denote the first factor of the product in (17) by V and consider separately its first addend:

$$V = \int_{\sigma(B_0)} \int_{\sigma(B_0)} \frac{E(d\theta) E(d\eta)}{P_1(\theta, \eta)} + \sum_{n_1+n_2=1}^{\infty} \frac{N^{n_1+n_2}}{n_1!n_2!} \int_{\sigma(B_0)} \int_{\sigma(B_0)} \frac{\partial^{n_1+n_2}}{\partial \theta^{n_1} \partial \eta^{n_2}} \left(\frac{1}{P_1(\theta, \eta)} \right) E(d\theta) E(d\eta). \quad (18)$$

Similar to work [3], we can prove that except for the first term in (18) the rest of the sum represents a quasinilpotent operator, and $\int_{\sigma(B_0)} \int_{\sigma(B_0)} \frac{E(d\theta)E(d\eta)}{P_1(\theta, \eta)}$ is an operator of scalar type.

From the general formula (4) for F_m it can be easily proved by induction that if U_0 is spectral and commutative with $A_i, B_j, i, j = 0, 1, \dots$, then all the operators F_m and consequently U_m are spectral. So, the following theorem is proved:

Theorem 2. Let all the conditions of theorem 1 be satisfied. If operators $A_i, B_j, i, j = 0, 1, \dots$, and U_0 are spectral and mutually commutative, then besides the statement of theorem 1, it is also true that operator coefficients $U_m, m = 1, 2, \dots$, in (2) are spectral too.

3. ON A SPECTRAL SOLUTION

Let's consider the conditions under which an operator-differential equation (1) in Hilbert space has a solution being a spectral operator.

Let \mathcal{Q} be a complete algebra in N . Danford sense ([2], XVII.1), generated by the family of commutative spectral operators $\tau = \{U_0, A_i, B_j, j = 0, 1, \dots\}$ and their resolutions of the identity operator, and closed in a uniform operator topology. It is clear that operators $U_m \in \mathcal{Q}$ and, therefore, the finite sums $\sum_{m=0}^n U_m z^m \in \mathcal{Q}$. As in paragraph 1 the convergence of series $\sum_{m=0}^{\infty} U_m z^m$ in a uniform operator topology was proved. Then, taking into account the closedness of algebra \mathcal{Q} in a uniform operator topology, the sum of series $\sum_{m=0}^{\infty} U_m z^m$ belongs to \mathcal{Q} , too. Suppose that the Boolean algebra generated by the resolutions of the identity of the operators of family $\tau = \{U_0, A_i, B_j, i, j = 0, 1, \dots\}$ is bounded. Then by theorem XVII.2.14 from [2] any operator from \mathcal{Q} is spectral and, therefore, so is the sum $\sum_{m=0}^{\infty} U_m z^m$.

Since R is a spectral operator, then by the known theorem on an analytic function of spectral operator ([2], XV.5.6) the operator $e^{R \ln z}$ is spectral, too.

As a function of B_0 operator z^R commutes with all $U_m, m = 0, 1, \dots$, and, therefore, with $\sum_{m=0}^{\infty} U_m z^m$. The product of two commutative spectral operators in Hilbert space is a spectral operator, so $U(z) = (\sum_{m=0}^{\infty} U_m z^m) z^R$ is a spectral operator too. So we proved the following

Theorem 3. If Boolean algebra, generated by the resolutions of the identity operator of spectral commutative operators of family $\tau = \{U_0, A_i, B_j, j = 0, 1, \dots\}$ is bounded and the conditions of theorem 1 are satisfied, then equation (1) has a solution being a spectral operator.

It should be noted that the question of solvability of equation (1) was investigated in the partial case in the papers [3-6].

REFERENCES

1. Daletski, T. L. & Kreyn, M. G. (1970). Stability of solutions to the differential equations in the Banach space. *Moscow, 534, (in Russian)*.
2. Dunford, N. & Schwartz, J. T. (1974). Linear operators, Part III, Spectral Operators. *Moscow*.
3. Akhmedov, A. M. & Gadjiyev, A. A. (1996). On the solvability of some classes of operator equations, *Proc. of*

Azer. Math. Soc., 2, Baku (3-13).

4. Akhmedov, A. M. & Gadjiyev, A. A. (1997). On the solvability of operator-differential equations of the second order. *J. News of Baku State Univ., ser. of phys. and math. sci.*, 2 (in Russian).
5. Akhmedov, A. M. (1998). On the solvability of operator-differential equations in complex domain. *Works of the Inst of Math. & Mech.*, VII (XVI), Baku, 29-33.
6. Gadjiyev, A. A. (1997). Solvability of some operator-differential equations in Banach space. *Cand. dissert. Baku*, 111.

Archive of SID