Archive of SID

Iranian Journal of Science & Technology, Transaction A, Vol. 34, No. A1 Printed in the Islamic Republic of Iran, 2010 © Shiraz University

## HIGH SPIN STATES IN DEEP-INELASTIC AND COMPOUND-NUCLEUS REACTIONS<sup>\*</sup>

## S. MOHAMMADI

Physics Department, Payame Noor University, Mashad, 91735, I. R. of Iran Emails: Mohammadi@pnu.ac.ir & smohammadi1958@yahoo.co.uk

Abstract – Compound-nucleus reactions provide the standard mechanism to populate states with high angular momentum in neutron deficient nuclei. Neutron-rich nuclei with mass A < 150 can be studied in spontaneous and induced fission. Projectile fragmentation has proven to be an efficient method of populating nuclei far from the valley of stability. However, in the case of heavy nuclei this method is still limited to species with isomeric states. Deep-inelastic reactions are another reaction mechanism which can be used to study neutron-rich nuclei and are able to populate relatively high-spin states. In this article we compare the advantages and disadvantages of each method.

Keywords - Compound-nucleus, deep-inelastic, reaction mechanisms, neutron-rich nuclei, high spin states

### **1. INTRODUCTION**

The gamma rays de-exciting the high spin states in neutron deficient as well as neutron rich nuclei from fusion-evaporation and deep-inelastic reactions and other emission particles have been recorded in three heavy ion reaction experiments.

In order to study rare reaction channels, some method of channel selection must be employed to coincide with the detection of gamma-rays. This usually takes one of three forms (a) an array of charged particle and neutron detectors; (b) a mass separator to detect the recoiling nuclei and measuring their atomic mass number; and (c) an inner BGO ball which is acting as a multiplicity filter and a total-energy spectrometer.

The first two are usually used for fusion-evaporation reactions, and the third one for deep-inelastic reactions. The first one has the disadvantage that for very weak channels, target contaminants can dominate the gamma spectra. The second one can, in principle, give spectra with lower background and insensitivity to states below isomers, but is restricted by the transmission efficiency for recoiling nuclei of the mass separator. For the last case, the multiplicity of gamma rays is very important.

For a comparison of the different channel selections, some data from three different heavy ion reactions in different labs are used, which will be explained in some detail.

## a) Compound-Nucleus Reaction $^{19}Ne + {}^{40}Ca$

For this experiment we used the reaction  ${}^{19}\text{Ne} + {}^{40}\text{Ca}$  with a beam energy of 70 MeV. The radioactive  ${}^{19}\text{Ne}$  beam was produced at Louvain-la-Neuve accelerator laboratory in a two stage process using the isotope separator on line (ISOL) method, which uses two cyclotrons. The first one produced 30 MeV protons which bombard a thick Lithium Fluoride target to produce the radioactive atoms via the  ${}^{19}\text{F}$  (p, n)  ${}^{19}\text{Ne}$  reaction. The radioactive  ${}^{19}\text{Ne}$  atoms as well as a large number of stable  ${}^{19}\text{F}$  isobaric contaminants

<sup>\*</sup>Received by the editor October 26, 2008 and in final revised form June 23, 2010

#### S. Mohammadi

were then injected into a second cyclotron, which was tuned as a mass spectrometer so that the intensity of the <sup>19</sup>F contaminants was reduced far below the radioactive beam intensity after acceleration. Finally, the beam was incident on a thick, 1.6 mgcm<sup>-2</sup>, <sup>40</sup>Ca target.

Gamma rays were identified from residual nuclei using an array of 8 TESSA [1] (Total Energy Suppression Shield Array) germanium detectors. Charge particles evaporated in the reaction were detected by an array of 128 silicon strip segments with a thickness 300 µm arranged in an octagonal shape and placed in the forward direction. The arrangement of the silicon array in the forward hemisphere and the gamma detectors in the backward hemisphere were convenient experimentally.

Figure 1 shows channel selections gated on charge particle multiplicities of 1, 2 and 3. Some results of this experiment about the production of nuclei were published before [2, 3].



Fig. 1. Random subtracted spectra gated on protons/  $\alpha$ -particle multiplicities of 1, 2 and 3

## b) Compound-Nucleus Reaction $^{24}Mg + {}^{40}Ca$

In this experiment, the fusion-evaporation reaction  $^{24}Mg + {}^{40}Ca$  with a beam energy of 65 MeV provided by ATLAS accelerator was performed at Argonne National Laboratory. Gamma rays were detected using the AYEBALL detector array, consisting of 18 germanium detectors of both 20% efficient TESSA type detectors and 70% efficient EUROGAM [4] detectors.

Isobaric identification of subsequent decay gamma rays was achieved by detecting the recoiling nuclei through the Argonne fragment mass analyzer (FMA) [5]. For a given charge state, the FMA disperses the residual nuclei according to their mass over charge (A/Q) ratio in the X direction at the focal plane, where they are detected by a position sensitive parallel grid avalanche counter (PGAC) [5]. Some results of this experiment regarding the production of nuclei have been previously published [6, 7]. Elemental separation was provided by monitoring the recoil energy loss in a split anode ionization chamber placed behind the focal plane of the FMA. A ring of eleven NE213 scintillation detectors was placed in front of the AYEBALL array, at the entrance of the FMA. These detectors were used to detect

neutrons and subsequently select evaporation channels in the analysis, which involved one or more neutrons.

Gating on the recoils reduces the amount of scattered beam in the subsequent spectra. In Fig. 2, we have shown the gamma ray spectra of the separated nuclei.

Finally, Fig. 3 shows a comparison of the effectiveness of using the ion chamber and neutron detectors to separate different nuclei from each other. As is shown in the figure, using the ion chamber gating is a more efficient method of obtaining isotopically pure spectra than neutron gating.



Fig. 2. Z separation for obtained nuclei by gating on different parts of the total ion chamber signal



Fig. 3. Comparison of the effectiveness of using ion chamber and neutron detectors to separate the 3pn channel from the 4p channel in the A=86 gated data

#### S. Mohammadi

# c) Deep-Inelastic Reaction $^{82}Se + ^{192}Os$

In order to obtain information on the ground state bands of neutron rich nuclei around A~190, the  ${}^{82}$ Se +  ${}^{192}$ Os deep inelastic reaction at 460 MeV bombarding beam energy we used to populate the nuclei around  ${}^{192}$ Os. A Thick  ${}^{192}$ Os target (>50 mg/cm<sup>2</sup>) with 0.2 mm Ta backing was used to stop all of the recoils in the target, minimizing the broadening of the lines due to Doppler shift.

The bombarding energy was obtained from the ALPI linear accelerator and was chosen to be 20% above the Coulomb barrier of the colliding nuclei. Gamma rays were detected using Gamma ray spectrometer (GASP) [8] array at Legnaro, Italy, which consist of 40 Compton suppressed hyper pure high efficiency n-type germanium detectors and a  $4\pi$  calorimeter composed of 80 BGO crystals. The geometry of the GASP array is based on a polyhedron with 122 faces. 40 faces are used by the germanium detectors and the remaining 80 to the inner BGO ball. The BGO detector thickness (65 mm) is sufficient to absorb 95% of gamma rays of 1 MeV. The resulting total efficiency is 70%. In the case of high multiplicity events, like in standard fusion reactions, the total inner ball efficiency is very close to 100%. The read-out of the crystal is made with standard PMT `s and the electronic treatment of the signals are such that the energy and time information of each individual BGO detector can be recorded. The BGO ball adds a background reduction factor that is reaction dependent, but can be conservatively estimated to be about 2.

Two and three dimensional gamma ray matrices were used to construct level schemes of the nuclei of interest. Typical coincidence spectra are shown in Figs. 4 and 5. To determine the spine and parity of states in a cascade of radiations, it is necessary to measure the angular distribution of one radiation relative to the direction of another radiation. Some results of this experiment were also published before [9-12].



Fig. 4. Gamma ray coincidence spectrum for <sup>188</sup>Os nucleus gated on 574 keV transition



Fig. 5. Gamma ray coincidence spectrum for <sup>190</sup>Os nucleus gated on 589 keV transition

### 2. RESULTS AND CONCLUSIONS

Here, the results for two compound nucleus reactions and one deep inelastic reaction are compared. As can be seen from Fig. 1, charged particle detection proved to be a very useful on-line tool for identifying gamma rays from a compound nucleus reaction. An ideal detector would be compact, but the design would need to address the problems arising from beam particles that elastically scattered from the target. The granularity of the silicon detector and the choice of thickness were each beneficial. The moderate thickness of the silicon allowed protons to be effectively distinguished from alpha particles, which can lead to confident channel identification. However, by going to weak channels like  $2\alpha$ , 3p or  $3\alpha$ , the statistics becomes very poor.

In the second compound nucleus experiment, the mass and charge of the recoiling nuclei can be identified using the Fragment Mass Analyzer (FMA). For certain recoils, a charge state anomaly can cause an isobaric contamination in the gated spectra. This contamination can later be removed with Z separation, using an energy loss signal from a split anode ion chamber at the focal plane of the instrument. In this experiment, neutron detectors were used to detect neutrons and subsequently select evaporation channels which involve one or more neutrons. However, as Fig. 3 shows, the neutron evaporation channels could be clearly resolved by deconvoluting the spectra with and without the neutron condition. In general, the ion chamber gating method was preferred to separate the isobarically gated spectra by an individual element. Here again the statistics become poor by using additional detector separators.

In the last experiment, deep inelastic reactions are the most general reaction mechanism which can be used to study neutron rich nuclei in any mass region. In this experiment, although we used only gamma ray detectors with increasing numbers, the statistics of spectra are relatively good but not sufficient, which means that we must use additional channel separators.

Thus by using a large number of gamma ray detectors together with other detectors like recoil separators, charge particle detectors and ion chambers, unambiguous channel selections will be possible for both neutron deficient and neutron rich regions.

### S. Mohammadi

#### REFERENCES

- Nolan, P. J., Gifford, D. W. & Twin, P. J. (1985). The Performance of a Bismuth Germanate Escape Suppression Spectrometer. *Nucl. Instr. and Meth.*, A236, 95.
- Catford, W. N., Mohammadi, S., Regan, P. H., Purry, C. S., Gelletly, W., Walker, P. M., Gyapong, G. J., Simpson, J., Warner, D. D., Davinson, T., Neal, R., Page, R. D., Shotter, A. C., Hibbert, I. M., Wadsworth, R., Forbes, S. A., Bruce, A. M., Thwaites, C., Thirolf, P., Van Duppen, P., Galster, W., Ninane, A., Vervier, J., Decrock, P., Huyse, M., Szerypo, J. & Wauter, J. (1997). High resolution gamma-ray spectroscopy with a radioactive beam. *Nucl. Phys.* A616, 303c.
- Catford, W. N., Mohammadi, S., Regan, P. H., Purry, C. S., Gelletly, W., Walker, P. M., Gyapong, G. J., Simpson, J., Warner, D. D., Davinson, T., Neal, R., Page, R. D., Shotter, A. C., Hibbert, I. M., Wadsworth, R., Forbes, S. A., Bruce, A. M., Thwaites, C., Thirolf, P., Van Duppen, P., Galster, W., Ninane, A., Vervier, J., Decrock, P., Huyse, M., Szerypo, J. & Wauter, J. (1996). High resolution gamma-ray spectroscopy with a radioactive beam. *Nucl. Instr. and Meth.*, A371, 449.
- 4. Beausang, C. W., Forbes, S. A., Fallon, P., Nolan, P. J., Twin, P. J., Mo, J. N., Lisle, J. C., Bentley, M. A., Simpson, J. C., Beck, F. A, Curien, D., deFrance, G., Duchêne, G. & Popescu, D., Measurements on prototype Ge and BGO detectors for the Eurogam array. (1992). *Nucl. Instr. and Meth.*, A313, 37.
- Davids, C. N., Back, B. B., Bindra, K., Henderson, D. J., Kutschera, W., Lauritsen, T., Nagame, Y., Sugathan, P., Ramayya, A. V. & Walters, W. B.(1992). In Beam Spectroscopy. *Nucl. Instr. and Meth.*, *B70*, 358.
- Vincent, S. M., Regan, P. H., Mohammadi, S., Gelletly, W., Blumenthal, D., Carpenter, M. P., Davids, C. N., Janssens, R. V. F., Lister, C. J., Pearson, C. J., Simpsong, J., Seweryniak, D., Schwartz, J. & Warner, D. D. (1999). Near yrast study of the *fpg* shell nuclei <sup>58</sup>Ni, <sup>61</sup>Cu and <sup>61</sup>Zn. *Physical Review C*, V60, 064308.
- Mohammadi, S., Vincent, S. M., Regan, P. H., Gelletly, W., Blumenthal, D., Carpenter, M. P., Davids, C. N., Janssens, R. V. F., Lister, C. J., Pearson, C. J., Simpsong, J., Seweryniak, D., Schwartz, J. & Warner, D. D. (1999). High spin states in <sup>58</sup>Ni. *Journal of Physics*, *G*25, 909.
- 8. Bazzaco, D. (1992). Proc. Of the Int. Conf. on Nuclear Structure at High AngularMomentum (376). Ottawa, Report No. AECL 10613.
- Mohammadi, S., Podolyak, Zs., Angeles, G. D. E., Zhang, Y. H., Axiotis, M., Bazzacco, D., Bizzeti, P. G., Brandolini, F., Broda, R., Bucurescu, D., Farnea, E., Gelletly, W., Gadea, A., Ionescu-Bujor, M., Iordachescu, A., Kroll, T., Langdown, S. D., Lunardi, S., Marginean, N., Martinez, T., Medina, N. H., Quintana, B., Regan, P. H., Rubio, B., Ur, C. A., Valiente-Dobon, J. J. & Walker, P. M. (2004). High-Spin States Populated in Deep-Inelastic Reactions. *Brazilian Journal of Physics*, V34(3A), 792.
- Podolyak, Zs., Mohammadi, S., Angeles, G. D. E., Zhang, Y. H., Axiotis, M., Bazzacco, D., Bizzeti, P. G., Brandolini, F., Broda, R., Bucurescu, D., Farnea, E., Gelletly, W., Gadea, A., Ionescu-Bujor, M., Iordachescu, A., Kroll, T., Langdown, S. D., Lunardi, S., Marginean, N., Martinez, T., Medina, N. H., Quintana, B., Regan, P. H., Rubio, B., Ur, C. A., Valiente-Dobon, J. J. & Walker, P. M. (2004). Structure of Neutron-Rich Nuclei from Deep-Inelastic Reactions. *International Journal of Modern Physics E, V13*(1), 123.
- Mohammadi, S., Angels, G. De, Axiotis, M., Bazzacco, D., Bizzeti, P. G., Brandolini, F., Broda, R., Bucurescu, D., Farnea, E., Gelletly, W., Gadea, A., Ionescu\_Bujor, M., Iordachescu, A., Kroll, Th., Longdown, S., Lunardi, S., Marginean, N., Martinez, T., Medina, N., Podolyak, Zs., Quintana, B., Regan, P. H., Rubio, B., Ur, C. A., Valiente, J.-J., Walker, P. M. & Zhang, Y. H. (2006). Yrast States in <sup>188, 190</sup>Os Nuclei. *International Journal of Modern Physics E, V15*(8), 1797.
- 12. Mohammadi, S. & Podolyak, Zs. (2007). High Spin States in <sup>191</sup>Os. IJST, Trans. A, 31(A2), 171.