
Biofacies and palaeoecology of the Jahrum Formation in Lar area, Zagros Basin, (SW Iran)

E. Nafarieh^{1*}, H. Vaziri-Moghaddam¹, A. Taheri² and A. Ghabeshavi³

¹*Faculty of Sciences, Department of Geology, University of Isfahan, Isfahan, Iran*

²*Faculty of Earth Sciences, Department of Geology, Shahrood University of Technology, Shahrood, Iran*

³*Department of Geology, National Iranian South Oil Company, Ahvaz, Iran*

E-mails: el_2090na@yahoo.com, avaziri7304@gmail.com, ali.ataheri@gmail.com & aghabeishavi@yahoo.com

Abstract

Biofacies and palaeoecology of the limestone of the Jahrum Formation in the Lar area in the southwest of Iran (Zagros Basin) is addressed in this paper. Our detailed analysis of biofacies and palaeoecology shows that the Jahrum Formation in the studied area were deposited in a carbonate open shelf dominated by heterozoan and, subordinately, photozoan skeletal assemblages. Based on analysis of larger benthic foraminiferal assemblages and biofacies features, two major depositional environments are identified. These include inner shelf and middle shelf environments. The inner shelf facies is characterized by wackestone-packstone, dominated by various taxa of imperforate foraminifera. The middle shelf is represented by wackestone-packstone with a diverse assemblage of larger foraminifera with perforate wall. The distribution of the larger benthic foraminifera indicates that shallow marine carbonate sediments of the Jahrum Formation at the studied areas have been deposited in the photic zone of tropical to subtropical oceans.

Keywords: Biofacies; Palaeoecology; Jahrum Formation; Eocene; Zagros Basin

1. Introduction

The limestone of Jahrum Formation (Paleocene-Eocene), constitutes one of the reservoir intervals in the Zagros Basin in Iran. The studied area is located in Fars Province (SW Iran), which is part of the Zagros fold-and-thrust belt (Fig. 1) [1, 2]. The Jahrum Formation, the focus of this study, takes its name from the type section at Kuh-e Jahrum in Fars Province. The type section was described by James and Wynd [3]. It is divided into three carbonate units with a total thickness of 467.5 m. This formation progressively overlies the silty marl, dolomites and evaporates of the Sachun Formation (Fig. 2) [4]. Where the latter is absent, it overlies either the Pabdeh or Gurpi formations. The upper contact with the Asmari Formation is unconformable. On the Fars area, there is a transition between the Jahrum and Pabdeh Formation. Studies of the Jahrum Formation have focused mainly on their lithostratigraphy and biostratigraphy, James and Wynd [3]; Rahaghi [5]; Kalantari [6]; Hottinger [7]. A few previous studies have been focused on detailed investigation of

work is still necessary. Shallow marine carbonate sediments of the Jahrum Formation exhibit a great diversity and abundance of larger foraminifera, and they can be easily identified in thin section and in the field. Consequently, larger foraminifera provide a useful tool for reconstructing paleoenvironments in lithologically monotonous Jahrum successions. The most prominent components of the studied sediments are nummulitids and alveolinids.

This paper examines in detail the biofacies of Jahrum Formation in the Lar area and provides palaeoenvironmental interpretations of the sedimentary succession.

2. Methods of study

More than 328 thin sections were analyzed under the Petrographic microscope for biofacies composition. The textural classification of Dunham [11] and Embry and Klovan [12] were used to describe biofacies types. The biofacies and assemblages of benthic hyaline and imperforate foraminifera are used in interpreting palaeoenvironmental conditions of the Jahrum Formation.

*Corresponding author

Received: 30 May 2011 / Accepted: 29 January 2012
sedimentological aspects [8-10], however, detailed palaeoenvironmental interpretations and biofacies



Fig. 1. Location of the studied sections in southwest Iran




Fig. 2. Cenozoic stratigraphic correlation chart of the Iranian Sector of the Zagros Basin, [4]

3. Geological setting

The Iranian plateau extends over a number of continental fragments welded together along suture zones of oceanic character. Each fragment differs in its sedimentary history, age of magmatism and metamorphism, and its structural character and intensity of deformation [13]. These fragments are in the following provinces: (1) Zagros, (2) Sanandaj-Sirjan, (3) Urumieh-Dokhtar, (4) Central Iran, (5) Alborz, (6) Kopeh Dagh, (7) Lut and (8) Makran (Fig. 3) [14]. The Jahrum Formation is part of the Cenozoic deposits (Paleocene-Eocene) of the Zagros Basin in southwest Iran. The Zagros Mountains and adjacent areas are well-known for their vast huge hydrocarbon reservoirs and very young tectonic activities [2].





Fig. 3. (a) General map of Iran showing eight geologic provinces [13]. (b) Subdivisions of the Zagros province

The Zagros Basin was a continental margin attached to the eastern edge of Africa during the Phanerozoic. Throughout the Permian, detachment of the Iran plate including Alborz, Central-East-Iran microcontinent, and Sanandaj-Sirjan from the Arabian plate caused the formation of the Neo-Tethys Ocean [1, 15].

The closure of the Neo-Tethys Ocean, mostly during the Late Cretaceous, was because of the subduction of the Arabian plate toward the northeast, beneath the Iranian subplate [13, 16-18].

The Zagros fold-thrust belt resulted from the continent-continent collision between the Arabian margin and the Eurasian Plate, caused the subsequent closure of the Neo-Tethys Ocean during the Cenozoic [19, 20]. The Zagros fold-and-thrust belt of Iran is a result of the Alpine orogenic events [21] in the Alpine-Himalayan mountain range.

This study is based on two outcrop sections in the Lar area (Kuh-e Gach and Kuh-e Kurdeh) (Fig. 4).

Kuh-e Kurdeh section is located about 40 Km northeast of Lar city. This section was measured at 27° 49' 26" N and 54° 40' 9"E. The Kuh-e Gach section is located about 30 Km southeast of Lar City. It was measured in detail at 27° 38' 55" N and 54° 37' 16" E.

The total thickness of the Jahrum Formation is 404.5 m and 437 m in the Kuh-e Gach and Kuh-e Kurdeh sections, respectively.




Fig. 4. Location and geological map of the study areas, Kuh-e Gach and Kuh-e Kurdeh Anticline, sw of Iran, Lar area

4. Biofacies analysis

Facies analysis of Jahrum, the Formation in the study area has resulted in recognition of 7 biofacies types (Figs. 5, 6), characterizing platform development (Figs. 7, 8). Each biofacies is characterised by typical skeletal components and textures. The general environmental interpretations of the biofacies are discussed in the following paragraphs.

4.1. Biofacies A. Large and flat nummulitids bioclastic wackestone

The predominate fauna are larger benthic foraminifera with perforate walls (nummulitidae). Nummulitidae are represented by *Operculina* and *Nummulites*. This biofacies has a fine grained matrix. Other bioclasts include *Linderina*, *Amphistegina* and echinoid.

The foraminifera assemblage of this facies (Fig. 5a) shows close affinities to that described by Cosovic et al. [22] of the Adriatic carbonate platform (Istrian Peninsula) and Taheri et al. [10] of the Jahrum Formation in the Zagros Basin. Such assemblages are characteristic of lower slope carbonate environments.




Fig. 5. Microfacies of the Jahrum Formation at Kuh-e Gach section in Lar area. (a) Biofacies A, Large and flat nummulitids bioclastic wackestone. (b) Biofacies B, Lens-shaped nummulitids bioclastic wackestone-packstone. (c) Biofacies C, Foraminifera (perforate and imperforate) bioclastic wackestone-packstone. (d) Biofacies D, High diversity imperforate foraminifera packstone-grainstone. (e) Biofacies F, Miliolids packstone. (f) Biofacies G, Stromatolitic boundstone




Fig. 6. Microfacies of the Jahrum Formation at Kuh-e Kurdeh section in Lar area. (a) Biofacies B, Lens-shaped nummulitids bioclastic wackestone-packstone. (b) Biofacies C, Foraminifera (perforate and imperforate) bioclastic wackestone-packstone. (c) Biofacies D, High diversity imperforate foraminifera packstone-grainstone. (d) Biofacies E, *Dictyoconus Coskinolina* wackestone-packstone. (e) Biofacies F, Miliolids packstone. (f); Biofacies G, Stromatolitic boundstone.

The presence of large and flat foraminifera such as nummulitidae, in comparison with analogues in the modern platform [23-27], allowed us to interpret this facies as having been deposited in the lower photic zone.

4.2. Biofacies B. Lens-shaped nummulitids bioclastic wackestone-packstone

This biofacies (Figs. 5b, 6a) has a high diversity of benthic biota including large benthic foraminifera and echinoid. The larger foraminifera consists of small-lens shaped *Nummulites*, *Operculina* and *Amphistegina*. Fragmentation of larger foraminifera is common and they are distributed irregularly among the larger foraminifera. Nummulitids with

robust and small size tests are abundant biogenic components in biofacies B. Depositional textures are represented by wackestone- packstone. Peloids are also present.

The change in shape of test of larger perforate foraminifera with depth has been documented in the Cenozoic carbonate successions [28-32]. Proliferation of perforates benthic foraminifera is indicative of normal marine conditions [28]. The sediments with robust and lens specimens reflect shallower water than those containing larger and flat nummulitids and discocyclinids [29, 31]. The relatively high degree of fragmentation of the larger foraminifera indicate moderate turbulence conditions for this facies.




Fig. 7. Vertical facies distribution and biofacies of the Jahrum Formation at Kuh-e Gach section in Lar area, Zagros Basin




Fig. 8. Vertical facies distribution and biofacies of the Jahrum Formation at Kuh-e Kurdeh section in Lar area, Zagros Basin

4.3. Biofacies C. Foraminifera (perforate and imperforate) bioclastic wackestone-packstone

The main components of this biofacies (Figs. 5c, 6b) are benthic foraminifera, fragments of macrofossils and peloids. Both hyaline and imperforate foraminifera are present. Hyaline foraminifera are represented by small lens shaped *Nummulites*, *Linderina*, *Amphistegina*, *Operculina*, *Orbitolites*, and *Spherogypsina*, whereas among imperforate forms, miliolids, textularids, *Austrorillina*, *Archias*, *Peneroplis* and *Orbitolites*

are common. Echinoderms, gastropods and dasytylidae are also present. The features of biota and stratigraphic position of biofacies C indicate that sedimentation took place in the semirestricted lagoonal area. Co-occurrence of normal marine perforates foraminifera and platform-interior imperforates foraminifera suggest that there was no effective barrier present to separate the platform interior from the open marine [28, 33]. Nebelsick et al. [34], Corda and Brandano [35] and Vaziri-Moghaddam et al. [36] considered the similar facies as representative of a shelf lagoon.

4.4. Biofacies D. High diversity imperforate foraminifera wackestone-packstone grainstone

The abundant components of this biofacies are benthic foraminifera with imperforated walls such as: *Orbitolites*, *Somalina*, *Dictyoconus*, *Rhapydionina*, *Alveolina*, textularids and miliolids (Figs. 5d and 6c). These deposits include different textures ranging from wackestone to packstone and grainstone. Peloids are also present. In some samples, a subordinate amount of dasycladacean are also present.

The occurrence of a large number of imperforate foraminifera tests indicates that the sedimentation took place in a shelf lagoon setting with relatively low to moderate current energy [28, 33].

4.5. Biofacies E. *Dictyoconus Coskinolina* wackestone-packstone

This facies is dominated by benthic foraminifera (mainly *Dictyoconus* and *Coskinolina*) and other bioclasts (Fig. 6d). Small Peloids are also present. Textures reflect poorly sorted wackestone-packstone. Some of the grains have been partially micritized.

Both the fossil content and the sediment texture suggest a low-energy shallow subtidal environment. The features of component and stratigraphic position indicate that sedimentation took place in the lagoonal area.

4.6. Biofacies F. Miliolids packstone-grainstone

This biofacies is dominated by the occurrence of small miliolids and Peloids. Rare echinids fragments are also present (Figs. 5e and 6e).

This facies was deposited in very restricted innermost shelf areas. The abundance of peloids, miliolids and low diversity of fauna support this interpretation.

4.7. Biofacies G. Stromatolitic boundstone

These deposits are represented by a mud-supported texture formed by millimeter thick lamina, generally without fossils, irregularly undulating and laterally continuous (stromatolitic type cryptoalgal laminae) (Figs. 5f and 6f). The cyanobacteria with their filamentous features trapping and binding the sedimentary particles produced a laminated sediment or stromatolite.

This facies was deposited in a tidal flat environment [37-41]. Modern stromatolites are most common in shallow, intertidal and supratidal zones, although they may occur under subtidal conditions [42].

5. Sedimentary model

On the basis of biofacies variation, a sedimentary model can be proposed for the Jahrum Formation, suggesting that it was accumulated in an open shelf carbonate platform (Fig. 9). As a result of the facies interpretations and palaeoecology of larger foraminifera, it can be stated that middle shelf and higher portions of the inner shelf environments are present among the studied area. Planktonic foraminifera are absent in the studied sections, because the setting is located in shallow tropical sea environments, not-suitable for their accumulation.

In the study areas, the inner shelf deposits consist of an open lagoon, protected lagoon and tidal flat. Tidal flat facies is characterized by stromatolite boundstone. The wavy or flat-laminated stromatolite boundstones are formed by trapping and binding fine-grained carbonate sediments by cyanobacteria in the upper intertidal zone.

In the protected lagoon, the most abundant biofacies are medium to coarse grained larger foraminifera with imperforate wall-bioclast wackestone-packstone. The presence of imperforate foraminifera that include *Archaias*, *Peneroplis*, *Dendritina*, *Alveolina*, *Austrotrillina*, *Orbitolites*, *Dictyoconus*, *Coskinolina* and miliolids indicates a low-energy, upper photic, shallow shelf lagoon depositional environment. Generally the upper photic zone is dominated by porcellaneous larger foraminifera, predominantly living in symbiosis with dinophyceans, chlorophyceans or rhodophyceans [33]. Open lagoon shallow subtidal environments are characterized by biofacies types that include mixed open marine bioclasts (such as echinoids and perforate foraminifera) and protected environment fauna (such as imperforate foraminifera). The diversity association of skeletal components represents a shallow subtidal environment, with optimal conditions with regard to salinity and water circulation.

Nummulitids with robust and small size tests are abundant fauna in the upper middle shelf environments. The sediments with robust and lens specimens reflect shallower water than those containing larger and flat nummulitids and discocyclinids [29, 31]. The change in shape of test of the larger perforate foraminifera with depth has been documented in the Cenozoic carbonate successions [28-32]. The relatively high degree of fragmentation of the larger foraminifera points to moderate turbulence conditions for this facies. Lower middle shelf facies are differentiated from upper middle shelf by the greater amount of micritic matrix, an increase in the flatness, and size of the perforate foraminifera.

6. Palaeoecology

Larger foraminifera are important constituents of shallow-water carbonates from the Early Eocene to Early Miocene of SW Iran (Zagros Basin). Larger foraminifera which occupied most niches in the photic zone of tropical to subtropical oceans in the Tethyan realm during the Paleogene, provide a useful tool for reconstructing palaeoenvironments and biostratigraphy.

Scheibner et al. [43] proposed that the larger foraminifera turn over (LFT) during the Palaeocene-Eocene transition closely linked with the Paleocene-Eocene thermal maximum (PETM). Larger foraminifera are extreme K-strategists, flourishing in a constant, typically oligotrophic environment [44].

Carbonate production directly or indirectly depends on photosynthesis and consequently on light penetration into water column. The Jahrum Formation contains both larger benthic imperforate foraminifera and, subordinately hyaline foraminifera. Both groups of larger foraminiferal are often supported by endosymbiotic relationships with unicellular algae.

The paleoenvironmental distribution of foraminiferal assemblages and depositional conditions have been reconstructed, based on the depth range of recent foraminifera, foraminifera-

bearing Early Oligocene carbonates from the Lower Inn Valley of Austria [34], Oligo-Miocene foraminiferal limestones of the Zagros Basin [36, 45, 46], Eocene foraminifera limestones of the Adriatic carbonate platform [22], Paleocene-earliest Eocene larger benthic foraminifera of SW Slovenia [47], benthic carbonate assemblages across the Paleocene-Eocene boundary of the Campo section [48] and Early Eocene foraminiferal limestones of the Pyrenees [49]. The occurrence of a large number of porcelaneous imperforate foraminiferal tests may point to the depositional environment being slightly hypersaline. The biotic assemblage indicate a deposition within the photic zone, in a seagrass-dominated environment, as suggested by the presence of epiphytic porcellaneous foraminifera (*Alveolina*, *Archaias*, *Peneroplis*), such an assemblage has been interpreted as a shelf-lagoon environment [50, 37, 51, 36].

Perforate foraminifera that exist in shallow water are characterized by hyaline walls and they protect themselves from UV light by producing very thick, lamellate test walls to avoid photo inhibition of symbiotic algae within the test in bright sunlight, and/or test damage in turbulent water or they occur in moderately deeper water. Flatter tests and thinner test walls with increasing water depth reflect decreased light levels at greater depths or possibly poor water transparency in shallow water [29].




Fig. 9. Depositional model for the platform carbonates of the Jahrum Formation at the studied sections

7. Skeletal grain association

The differentiation between non-tropical and tropical carbonates is mainly based on analysis of skeletal components [52-54]. Non-tropical carbonates are identified by the absence of certain skeletal and non-skeletal grains (e.g. *Halimeda*, ooids) and framework-forming zooxanthellate corals [55]. Non-tropical carbonates are subdivided into warm-temperate and cool-temperate provinces [56]. Tropical associations are restricted to the tropics, whereas sediments of temperate latitudes also extend into the tropics [52- 54, 57]. Lees and Buller [52], Lees [56], Carannante et al. [53] and Hallock and Schlager [58] have concluded that parameters such as depth, nutrients and salinity additionally influence distribution of skeletal grains.

Consequently, certain skeletal grains may exist in various environments. James [55] introduced two new terms for carbonate grain associations ('Photozoan' and 'Heterozoan') that are applicable to the entire Phanerozoic. The cool water sediments are always heterozoan, however, the heterozoan association does not mean that the carbonates are cool water [55].

The most common skeletal components in limestones of the study areas are large benthic foraminifera, whereas echinoids, bryozoans and bivalve components are less common (Fig. 10). Corals are almost absent. The biotic associations and palaeolatitudinal reconstructions [59] suggest that carbonate sedimentation of the Jahrum Formation took place in tropical waters under oligotrophic conditions and is dominated by a heterozoan skeletal assemblage.

Zooxanthellate corals did not make framework structures in the lower latitudes due to worldwide warm sea-surface temperatures and enhanced CO₂ levels [48]. Unlike zooxanthellate corals, increasing summer sea-surface temperatures do not cause symbiont loss in larger foraminifera [60]. Larger foraminifera and their symbionts algae appear to be less susceptible to high summer temperatures [61]. Therefore, the extension of heterozoan assemblages in the Jahrum Formation related to the palaeoecology of zooxanthellate corals.

8. Conclusions

Biogenic components of the Jahrum Formation are dominated by benthic foraminifera. Based on biogenic components and textures, 8 biofacies have been recognized and grouped into two depositional environments that correspond to the inner and middle shelf environments, and are interpreted as a

carbonate platform developed in an open shelf settings. As a result of the facies interpretations and palaeoecology of larger foraminifera, it can be stated that middle shelf and higher portions of the inner shelf environments are present among the studied areas. The biotic assemblages of the Jahrum Formation suggest that carbonate sedimentation took place in subtropical waters with oligotrophic conditions.




Fig. 10. (a) *Dictyoconus indicus*, (J330). (b) *Alveolina rutimeyeri*, (J303). (c) *Orbitolites* sp., (J374), (d) *Linderina brugesi*, (J278). (e) *Somalina stefaninii*, (J256) (f) *Penarchaias glynnjonesi*(3203) (g) *Pyrgo* sp. (3146) (h) *Nummulites* sp., (J274) (i) *Medocia blayensis* (3199). (j) Corallinacean fragment, (J270). (k) Bryozoan fragment (J275). (l) *Amphistegina* sp. (3202)

Acknowledgments

The authors wish to thank the reviewers for their helpful and constructive comments. We would also like to thank the University of Isfahan for providing financial support and National Iranian South Oil Company for fieldwork logistics.

References

- [1] Alavi, M. (1994). Tectonics of the Zagros Orogenic belt of Iran, new data and interpretation. *Tectonophysics*, 229, 211-238.
- [2] Alavi, M. (2004). Regional stratigraphy of the Zagros Fold-Thrust Belt of Iran its proforeland evolution. *American Journal of Sciences*, 304, 1535-1541.
- [3] James, G. A. & Wynd, J. G. (1965). Stratigraphic nomenclature of Iranian Oil Consortium Agreement Area. *AAPG Bulletin* 49 (12), 2182-2245.
- [4] Ala, M. A. (1982). Chronology of trap formation and migration of hydrocarbons in Zagros sector of southwest Iran. *AAPG Bull.* 66, 1536-1542.

- [5] Rahaghi, A. (1980). *Tertiary faunal assemblage of Qom-Kashan, Sabzewar and Jahrum area*. Tehran, National Iranian Oil Company.
- [6] Kalantari, A. (1981). *Iranian Fossils*. Tehran, National Iranian Oil Company.
- [7] Hottinger, L. (2007). Revision of the foraminiferal genus *Globoreticulina* Rahaghi, 1978, and of its associated fauna of larger foraminifera from the late Middle Eocene of Iran. *Carnets de Géologie / Notebooks on Geology, Brest, Article 2007/06* (CG2007-A06) 51p.
- [8] Vaziri-Moghaddam, H., Syrafian, A. & Taraneh P (2002). Biofacies and sequence stratigraphy of the Eocene succession, at Hamzeh-Ali area, north-central Zagros, Iran. *Carbonates and Evaporites*, 17, 60-67.
- [9] Najafi, M. Mahbubi, A. Moussavi-Harami, R. & Mirzaee, R. (2004). Depositional history and sequence stratigraphy of outcropping Tertiary carbonates in the Jahrum and Asmari formations, Shiraz area (SW Iran). *Journal of Petroleum Geology*, 27, 179-190.
- [10] Taheri, A. Vaziri-Moghaddam, H. & Seyrafian, A. (2008). Relationships between foraminiferal assemblages and depositional sequences in Jahrum Formation, Ardal area (Zagros basin, sw Iran). *Historical Biology: A Journal of Paleobiology*, 20, 191-201
- [11] Dunham, R. J. (1962). Classification of carbonate rocks according to their depositional texture, In Ham, W. E. (ed.), *Classification of Carbonate Rocks. A Symposium AAPG, Memo., 1*, 108-121.
- [12] Embry A. F. & Klovan J. E. (1971). A late Devonian reef tract on northeastern banks island, Northwest Territories. *Bulletin of Canadian Petroleum Geology*, 19, 730-781.
- [13] Berberian, M. & King, G. C. P. (1981). Towards the paleogeography and tectonic evolution of Iran. *Canadian Journal of the Earth Sciences*, 18, 210-265.
- [14] Heydari, E. Hassanzadeh, J. Wade, W. J. & Ghazi, A. M. (2003). Permian-Triassic boundary interval in the Abadeh section of Iran with implications for mass extinction. Part 1. *Sedimentology, Palaeogeography, Palaeoclimatology, Palaeoecology*, 193, 405-423.
- [15] Mohajjel, M. Fergusson, C. L. & Sahandi, M. R. (2003). Cretaceous-Tertiary convergence and continental collision, Sanandaj-Sirjan zone, western Iran. *Journal of Asian Earth Sciences*, 21, 397-412.
- [16] Stoneley, R. (1981). The geology of the Kuh-e-Dalneshin area of southern Iran, and its bearing on the evolution of southern Tethys. *Journal of Geological Society London*, 138, 509-526.
- [17] Beydoun, Z. R. Clarke, M. W. H. & Stoneley, R (1992). *Petroleum in the Zagros basin: A late Tertiary foreland basin overprinted on to the outer edge of a vast hydrocarbon-rich paleozoic-Mesozoic passive margin shelf*. Vol. 55, In: Macqueen R. W. & Leckie, D. A. (eds) *Foreland basins and fold belts*. American Association of Petroleum Geologists Memoir.
- [18] Berberian, M. (1995). Master "Blind" thrust faults hidden under the Zagros folds: Active basement tectonics and surface morphotectonics. *Tectonophysics*, 241, 193-224.
- [19] Stöcklin, J. (1968). Structural history and tectonics of Iran, a review. *American Association of Petroleum Geology Bulletin*, 52, 1229-1258.
- [20] Falcon, N. L. (1974). Southern Iran: Zagros mountains. In: Spencer A (ed) *Mesozoic-Cenozoic orogenic belts. Geological Society of London Special Publication*, 4, 199-211.
- [21] Ricou, L. E. Braud, J. & Brunn, J. H. (1977). Le Zagros Mé'm Hors-se'r . *Soc Ge'ol Fr*, 8, 33-52.
- [22] Cosovic, V. Drobne, K. & Moro, A. (2004). Paleoenvironmental model for Eocene foraminiferal limestones of the adriatic carbonate platform (Istrian Peninsula). *Facies*, 50, 61-75.
- [23] Hottinger, L. (1980). Repartition comparee des grands foraminiferes de la mer Rouge et del, Ocean Indian. *Ann Univ Ferrara*, 6, 35-51.
- [24] Hottinger, L. (1982). Larger Foraminifera, giant cells with a historical background. *Naturwissenschaften* 69, 361-371
- [25] Hohenegger, J. (1996). Remarks on the distribution of larger foraminifera (Protozoa) from Palau (western Carolines). In: Aoyama T(ed) *The progress report of the 1995 survey of the research project, man and the environment in Micronesia*. Kagoshima University Research Center for the Pacific Islands, *Occasional Papers*, 32, 19-45.
- [26] Hallock, P. (1999). Symbiont-bearing foraminifera, in: Gupta BKS (ed) *Modern foraminifera. Kluwer Academic, Dordrecht*, 123-139.
- [27] Reiss, Z. & Hottinger, L. (1984). The Gulf of Aqaba. *Ecological Micropaleontology*. 50, 354.
- [28] Geel, T. (2000). Recognition of stratigraphic sequences in carbonate platform and slope deposits: Empirical models based on microfacies analysis of Paleogene deposits in southeastern Spain. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 155, 211-238.
- [29] Beavington-Penney, S. J. & Racey, A. (2004). Ecology of extant nummulitids and other larger benthic foraminifera: Applications in palaeoenvironmental analysis. *Earth-Science Reviews*, 67(3-4), 219-265.
- [30] Nebelsick, J. H. Rasser, M. W. & Bassi, D. (2005). Facies dynamics in Eocene to Oligocene circumalpine carbonates. *Facies*, 51, 197-216.
- [31] Barattolo, F. Bassi, D. & Romano, R. (2007). Upper Eocene larger foraminiferal-coralline algal facies from the Klokova mountain (southern continental Greece). *Facies*, 53(3), 361-375.
- [32] Bassi, D. Hottinger, L. & Nebelsick, J. H. (2007). Larger foraminifera from the Upper Oligocene of the Venetian area, north-east Italy. *Palaeontology*, 50(4), 845-868.
- [33] Romero, J. Caus, E. & Rosell, J. (2002). A model for the palaeoenvironmental distribution of larger foraminifera based on late Middle Eocene deposits on the margin of the South Pyrenean basin (ne Spain). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 179(1-2), 43-56.
- [34] Nebelsick, J. H. Stingl, T. V. & Rasser, M. (2001). Autochthonous facies and allochthonous debris flows compared: Early Oligocene carbonate facies patterns of the lower Inn Valley (Tyrol, Austria). *Facies*, 44, 31-46.
- [35] Corda, L. & Brandano, M. (2003). Aphotic zone carbonate production on a Miocene ramp, central Apennines, Italy. *Sedimentary Geology*, 161(1-2), 55-70.
- [36] Vaziri-Moghaddam, H. Kimiagari, M. & Taheri, A. (2006). Depositional environment and sequence stratigraphy of the Oligo-Miocene Asmari Formation in sw Iran. *Facies*, 52(1), 41-51.

- [37] Flügel, E. (1982). *Microfacies Analysis of Limestones*. Berlin, Springer.
- [38] Hardie, L. A. & Lasemi, Y. (1995). Platform carbonates of the Upper Jurassic Mozduran Formation in the Kopet Dagh Basin, N. E. Iran-facies, palaeoenvironments and sequences. *Sedimentary Geology*, 99, 151-164.
- [39] Hardie, L. A. (1986). Ancient carbonate tidal flat deposits. *Q J Colorado Sch Mine* 81, 37-57S.
- [40] Steinhauß, D. M. & Walker, K. R. (1996). Sequence stratigraphy of an apparently non-cyclic carbonate succession: recognizing subaerial exposure in a largely subtidal, Middle Ordovician stratigraphic sequence in eastern Tennessee. In: Witzke G. A., Ludvigson, J. E., Day, B. J. (eds) Paleozoic sequence stratigraphy: views from the North American Craton. *Special Paper Geological Society of America*, 306, 87-115.
- [41] Aguilera-Franco, N. & Romano, U. H. (2004). Cenomanian - turonian facies succession in the guerrero-morelos basin, southern Mexico. *Sedimentary Geology*, 170, 135-162.
- [42] Dill RF, Shinn E. A, Jones, A. T., Kelly K & Steinen, R. P. (1986). Giant subtidal stromatolites forming in normal salinity water. *Nature*, 324, 55-58.
- [43] Scheibner, C. Speijer, R. P. & Marzouk, A. M. (2005). Turnover of larger foraminifera during the Paleocene-Eocene thermal maximum and paleoclimatic control on the evolution of platform ecosystems. *Geology*, 33(6), 493.
- [44] Hottinger, L. (1983). Processes determining the distribution of larger foraminifera in space and time. *Utrecht Micropaleont Bull*, 30, 239-253.
- [45] Amirshahkarami, M. Vaziri-Moghaddam, H. & Taheri, A. (2007a). Paleoenvironmental model and sequence stratigraphy of the Asmari Formation in southwest Iran. *Historical Biology: An International Journal of Paleobiology*, 19(2), 173-183.
- [46] Amirshahkarami, M. Vaziri-Moghaddam, H. & Taheri, A. (2007b). Sedimentary facies and sequence stratigraphy of the Asmari Formation at chaman-bolbol, Zagros basin, Iran. *Journal of Asian Earth Sciences*, 29, 947-959.
- [47] Zamagni, J. Mutti, M. & Košir, A. (2008). Evolution of shallow benthic communities during the late Paleocene-earliest Eocene transition in the northern Tethys (sw Slovenia). *Facies* 54(1), 25-43.
- [47] Scheibner, C. Rasser, M. W. Mutti, M. (2007). Facies changes across the Paleocene-Eocene boundary, the Campo section (Pyrenees, Spain) revisited. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 248, 145-168.
- [49] Rasser, M. Scheibner, C. & Mutti, M. (2005). A paleoenvironmental standard section for early ilderdian tropical carbonate factories (Corbieres, France; Pyrenees, Spain). *Facies*, 51(1), 218-232.
- [50] Wilson, J. L. (1975). *Carbonate facies in geological history*. Springer, 471p.
- [51] Flügel, E. (2004). *Carbonate Rocks, Analysis, Interpretation and Application*. Berlin, Springer-Verlag, 976.
- [52] Less, A. & Buller, A. T. (1972). Modern temperate - water and warm water shelf carbonate sediment contrasted. *Marine and Petroleum Geology*, 13, 67-73
- [53] Carannante, G. Esteban, M. Milliman, J. D. & Simone, L. (1988). Carbonate lithofacies as paleolatitude indicators: Problems and limitations. *Sedimentary Geology*, 60(1-4), 333-346.
- [54] Nelson, C. S. Keane, S. L. & Head, P. S. (1988). Non-tropical carbonate deposits of the modern New Zealand shelf. *Sediment Geol*, 160, 71-94.
- [55] James, N. P. (1997). The cool water carbonate depositional realm. In: James N. P. & Clarke, J. A. D. (eds) Cool-water carbonates. *SEPM. Spec Publ*, 56, 1-20.
- [56] Betzler, C. Brachert, T. & Nebelsick, J. H. (1997). The warm-temperate carbonate province: facies, zonations and delimitations. *Cour Forschinst Senckenberg*, 201, 83-100.
- [57] Lees, A. (1975). Possible influence of salinity and temperature on modern shelf carbonate sedimentation. *Mar Geol*, 19, 159-198.
- [58] Hallock, P. & Schlager, W. (1986). Nutrient excess and the demise of coral reefs and carbonate platforms. *Palaios*, 1(4), 389-398.
- [59] Alavi, M. (2007). Structures of the Zagros fold-thrust belt in Iran. *American Journal of Science*, 307(9), 1064-1095. doi:10.2475/09.2007.02.
- [60] Hallock, P. (2000). Symbiont-bearing foraminifera: harbingers of global change? *Micropaleontology* 46, supplement, 1, 95-104.
- [61] Hallock, P. Sheps, K. Chapronière, G. & Howell, M. (2006). Larger benthic foraminifers of the Marion plateau, northeastern Australia (odp leg 194): Comparison of faunas from bryozoan (sites 1193 and 1194) and red algal (sites 1196-1198) dominated carbonate platforms. Proceedings of the Ocean Drilling Program: *Scientific Results*, 194, 1-31.

بیو فاسیس و پالئواکولوژی سازند جهرم در منطقه لار، حوضه زاگرس (جنوب غربی ایران)

۱. نفریه^۱، ح. وزیری مقدم^۱، ع. طاهری^۲ و ع. غبیشاوی^۳

^۱گروه زمین شناسی، دانشگاه اصفهان، اصفهان، جمهوری اسلامی ایران
^۲گروه زمین شناسی، دانشگاه صنعتی شاهرود، شاهرود، جمهوری اسلامی ایران
^۳آداره زمین شناسی، شرکت ملی مناطق نفتخیز جنوب، اهواز، جمهوری اسلامی ایران

E-mails: el_2090na@yahoo.com, avaziri7304@gmail.com, ali.ataheri@gmail.com & aghabeishavi@yahoo.com

چکیده:

این پژوهش به مطالعه و بررسی رخساره های زیستی، محیط دیرینه و پالئواکولوژی سازند جهرم در برش های کوه گچ و کوه کورده در منطقه لار، جنوب غربی ایران (حوضه زاگرس) می پردازد. رسوبات سازند جهرم در برشهای مورد مطالعه در یک شلف باز و در دو محیط شلف داخلی و میانی نهشته شده اند که در شلف داخلی تنوع روزن داران کف زی با دیواره بدون منفذ و در شلف میانی روزن داران کف زی با دیواره منفذ دار می باشد. اجتماع دانه های کربناته از نوع فورامول و هتروزونن بوده و بازسازی عرض های جغرافیایی در منطقه مورد مطالعه بیانگر رسوب گذاری در شرایط های *SID* می باشد.