

Tanaffos (2002)1(1),46-49

©NRITLD, National Research Institute of Tuberculosis and Lung Disease, Iran

Comparison of Right Tei Index Between Normal Individuals and Patients with Pulmonary Disease

Behzadnia N, Sharif-Kashani B, Shokoofi-Moghiman Sh, Koroorian AR

Cardiology Unit ,NRITLD, Shaheed Beheshti University of Medical Sciences and Health Services, TEHRAN-IRAN

ABSTRACT

Background: Tei index is an echocardiographic indicator for studying the function of ventricles. In most studies, left Tei measurement was used to evaluate cardiac function in patients with heart failure, where its increase indicates left ventricular involvement. The following study was designed to compare right Tei index in healthy people with that of pulmonary patients in order to determine the effects of pulmonary diseases and their consequent cardiovascular complications.

Materials and Methods: The study population included 78 pulmonary patients and 37 normal individuals.

Doppler-echo imaging was performed by VingMed set, model CFM750 for all patients and the information was recorded in data sheets. Finally, results were analyzed by SPSS statistical software package, version 9.0.

Results: The mean (\pm SD) of age in the pulmonary patients and normal individuals was reported 49.28 ± 19.11 and 36.57 ± 15.46 respectively. Meanwhile, Tei index was revealed to be independent of age, sex, ventricular geometry, pulse, and blood pressure, thus no interference resulted from intervening factors. The mean (\pm SD) of right Tei index was measured 0.47 ± 0.25 and 0.38 ± 0.16 in the pulmonary patients and normal individuals respectively.

Conclusion: Regarding the difference found between Tei index of these two groups, it seems as if significant change in Tei index is strongly associated with pulmonary disease. Therefore, it is worth surveying, the correlation between Tei index and other cardiac indices in pulmonary disease. (Tanaffos 2002;1(1):46-49)

Keywords: Right Tei index, Normal individuals, Pulmonary disease

INTRODUCTION

The asymmetric feature of the right ventricle and its location has made common indices for left ventricle function studies such as ejection fraction, fractional shortening or Simpson volume change


assays to be ineffective for evaluating right ventricle function(1,2). Regarding independence of left ventricles Tei index from ventricular geometry and on the basis of reports considering effectiveness of Tei index in right ventricle pressure and volume overload(4,5,6), a study was designed to compare right Tei index measurement in two groups: the healthy individuals and the patients with pulmonary diseases. As it is shown in figure 1, to calculate the sum of IVC and IVR, the ejection time could be

Correspondence to: Sharif-Kashani B

Tel.-Fax.: +98-21-2803550

E-mail address: Babaksharifkashani@Yahoo.com

subtracted from the whole systole duration. Systole time may be measured from the interval between the two consequent flow of tricuspid valve and the ejection time is obtained from pulmonary valve flow time (3,8,9,11).



Schema of Doppler time intervals. The Tei index is the interval between cessation and onset of the mitral inflow. b = left ventricular ejection time. ICT = Isovolumetric Contraction Time, IRT = Isovolumetric Relaxation Time, ET = Ejection Time

MATERIALS AND METHODS

Between March 2000 and March 2001, a total of 115 patients were referred to the echocardiography clinic of Massih-Daneshvari Hospital. Echocardiographic right Tei index as well as other variables, including age, sex, ... were

studied. The study population included 78 as pulmonary patients and 37 as normal individuals. Pulmonary diseases consisted of COPD, tuberculosis, sarcoidosis, bronchiectasis, and malignancies. In the normal group, not even a trace of positive pathologic finding was elicited in history and physical examination. Results were recorded in special data sheets.

All patients underwent echocardiography in left lateral position by VingMed apparatus, CFM 750, using 3.25 MHz probe. 5-10 images were obtained from each patient. Measuring Tei index requires at least 5 images obtained from each position. Finally, our data were analyzed by SPSS statistical software package, version 9.0 and were reported using frequency, Cross Tab and Correlate (One Way ANOVA) options.

RESULTS

The results are summarized in table 1. The pulmonary patient population comprised of 48 women and 30 men with mean age (± SD) of 49.28 ± 19.11 years, whereas, the normal individuals were included 26 women and 11 men with mean age (± SD) of 36.57 ± 15.46 years.




Fig 1. Frequency of right Tei results in healthy and pulmonary groups

Table 1. Mean and Standard Deviation (S.D)of right Tei Index and age

Condition		Right-Tei	Age
<u>Pulmonary</u>	Mean	0.4704	49.28
	N	78	78
	S.D	0.2476	19.11
<u>Normal</u>	Mean	0.3822	36.57
	N	37	37
	S.D	0.1642	15.46
<u>Total</u>	Mean	0.4532	46.72
	N	134	134
	S.D	0.2234	19.14

DISCUSSION

Significant increase ($p < 0.05$) of right Tei index in these patients when comparing with normal individuals have implied that these changes should not be considered as a terminal phenomenon, indeed, these changes are developed gradually regardless of the underlying pulmonary disease. The mean (\pm SD) of right Tei index was measured 0.47 ± 0.25 and 0.38 ± 0.16 in these patients and normal individuals, respectively; however, this index was ranged 0.05-1.2 in patients with pulmonary disorders and 0.13-0.78 in normal individuals (Fig 1).

REFERENCES

- 1-Otto C. *Textbook of Clinical Echocardiography*. 2 nd ed. W.B Saunders; 2000
- 2- Feigenbaum H. *Echocardiography*. 5 th ed. Lea & Febiger; 1994.
- 3-Tei C. Doppler Echocardiography index for assessment of global Rt. Ventricular function. *J Am Soc Echo* 1996; 9: 838-47.
- 4- Tei C. Non invasive Doppler derived myocardial performance index correlation with simultaneous measurement of cardiac catheterization. *J Am Soc Echo* 1997,10:168-78.
- 5- Tsutsumi T. Serial evaluation for myocardial performance in fetuses and neonates using a new Doppler index. *Pediatr Int* 1999;41(6):722-7.
- 6-Bruch C. Tei index in patients with mild to moderate congestive heart failure. *Eur Heart J* 2000; 21(22):1888-95
- 7- Sutton M. The Tei index- a role in the diagnosis of heart failure. *Eur Heart J* 2000 ; 21(22): 1822-4
- 8- Kim WH. Estimation of left ventricular function in Right ventricular pressure and volume overload, Detection of early left ventricular dysfunction by tei index. *JPN Heart J* 1999; 40(2):145-54.
- 9-Genjyu Et. Assessment of global LV function in Normal children and in children with dilated cardiomyopathy. *J Am Soc Echo* 1999,12:1058-64.
- 10- Jacob E. Effect of preload alterations on a new Doppler Echocardiographic index of combined systolic and diastolic performance. *J Am Soc Echo* 1999;12: 1065-72.
- 11- Yeo TC. Value of a Doppler derived index combining systolic and diastolic time intervals in predicting outcome in primary pulmonary hypertension. *Am J Cardiol* 1998; 81:1157-61.

مقایسه شاخص Tei قلب راست بین بیماران ریوی و افراد سالم

دکترندا بهزاد نیا^۱، دکتر بابک شریف کاشانی^۱، دکتر شبنم شکوفی مقیمان^۲، دکتر علیرضا کروریان^۲
^۱استادیار بخش قلب، ^۲پزشک بخش قلب مرکز آموزشی، پژوهشی و درمانی سل و بیماری‌های ریوی، دانشگاه علوم پزشکی شهید بهشتی

سابقه و هدف: شاخص Tei، معیاری در اکوکاردیوگرافی است که برای ارزیابی عملکرد بطن‌ها بکار می‌رود. در بیشتر مطالعات شاخص Tei قلب چپ بیماران دچار نارسایی قلبی بررسی شده است. مطالعه حاضر برای بررسی شاخص Tei قلب راست در بیماران ریوی، انجام شده است.

مواد و روشها: مطالعه به صورت مورد - شاهدی بر روی ۷۸ بیمار ریوی (مورد) و ۳۷ فرد سالم (شاهد) انجام شد. از تمامی افراد مورد مطالعه، اکوکاردیوگرافی داپلر به عمل آمد.

یافته‌ها: میانگین سنی \pm SD دو گروه مورد و شاهد به ترتیب $49/28 \pm 19/11$ و $36/57 \pm 10/46$ سال بود. میانگین (\pm SD) شاخص Tei قلب راست در بیماران ریوی $0/47 \pm 0/25$ و در گروه شاهد $0/38 \pm 0/16$ بود.

نتیجه‌گیری و توصیه‌ها: شاخص Tei در بیماران ریوی بالاتر است و بررسی همبستگی بین شاخص Tei با سایر شاخص‌های عملکردی قلب در بیماران ریوی توصیه می‌شود.

واژگان کلیدی: شاخص Tei، قلب راست، بیماری ریوی