## Archive of SID



Material &

Method

Result





**Article Name** Pattern of Drug Resistance in Pulmonary TB Patients

AuthorsMajid Valioallahpour Amiri 1, Seyed Mehdi Mirsaeidi 1, Kiarash<br/>Mohajer 1, Seyed Davood Mansoori 1, Payam Tabarsi 1, Mohammad<br/>Reza Masjedi 2

Authors Specialization 1 Department of Infectious Disease, 2 Department of Pulmonary Medicine, NRITLD, Shaheed Beheshti University of Medical Sciences and Health Services, TEHRAN-IRAN

Tuberculosis (TB) is the most common cause of death due to infectious diseases globally after AIDS (1). Approximately, 2 billion people in the world are infected with mycobacterium tuberculosis, and 8.8 million new cases of tuberculosis occur annually, with over 50000 attributable deaths each week (2). The presence of multidrugresistant tuberculosis (MDR-TB) is a strong global threat to human health. Drug resistance is either primary (initial), in which TB bacilli are initially drug-resistant or secondary (acquired). The latter Introduction develops during treatment by patient?s inappropriate consumption of drug or inadequate and inappropriate treatment regimens. Tuberculous bacilli resistance is divided in to three categories including monoresistance, combined resistance, and multi-drug resistance (MDR). MDR is determined by resistance of bacilli at least to isoniazid and rifampicin. This study was performed for the evaluation of drug resistance of tuberculous bacilli in hospitalized patients with pulmonary TB in Massih Daneshvari Hospital.

> Between the year 2000 and 2002, we reviewed the files of hospitalized patients in TB ward of Massih-Daneshvari hospital and selected those who had been treated with suspicion of MDR TB. Among these, the files of patients whose antibiogram had confirmed drug resistance were selected. Demographic, paraclinical and radiologic data along with antibiogram and the kind of treatment given were obtained from records. Then data were analyzed by SPSS software.

The patients under study were 43 (27 men (63%) and 16 women (37%) with age range of 16-80 years). Among them, 25 cases (58%) were Afghan and 13(30%) were Iranian; the nationality of the remaining had not been recorded. Sputum smear was 1+, 2+, and 3+ in 14 (32.5%); 14 (32.5%); and 13 (30.1%) cases, respectively. Two patients (4.9%) had negative sputum smear but positive sputum culture. The sputum cultures of all patients were positive ranging from 1+ to 4+. Also, 14 (33%), 13 (30%) and 12 (28%) cases had 1+, 4+, and 3+ sputum cultures, respectively. Only 4(9%) patients had 2+ sputum culture. Around 95% of 43 cases had received anti-tuberculosis treatment for at least once. Meanwhile, the history of 1, 2, 3, and 4 times of treatment was obtained from 25 (58%), 12 (28%), 3(7%), and 1 cases respectively. Antibiogram was carried out for all patients in which bacilli were resistance to at least isoniazid and rifampicin (MDR-bacilli) in 38 (88%) cases. The status of TB bacilli resistance was as follows: there was resistance to isoniazid (INH), rifampicin (RIF), and streptomycin (STM) in 24 (56%) cases and the bacilli were only sensitive to ethambutol (EMB).

www.SID.ir

## Archive of SID

Discussion

In 4 (9%) and 2 (5%) cases, they were resistant to INH, RIF and INH as well as STM, respectively. Interestingly, TB bacilli were resistant at least to STM in 36 (85%) cases. Only 35 out of 43 cases had radiologic data (including chest X-ray and CT Scan), among whom 27 (63%) cases had cavitary lesions in their lungs, and the remaining 8 had other pulmonary lesions. Regarding treatment, 31 (72%) patients had received four-drug standard regimen plus ofloxacin and amikacin or only ofloxacin, so that the first drug regimen was used in 19 (44%) patients and the second in 12 (30%) patients. Other drug regimens were determined on the kind of antibiotic used including INH, pirazinamide (PZA), EMB, cycloserin, amikacin, ciprofloxacin, and ofloxacin.

In the United States 1997, about 8% of isolated Mycobacterium tuberculosis had primary resistant to INH, out of which 1.3% were resistant at least to INH and RIF (3). Estonia, Iran, China, India and some part of Russia were reported to be the most endemic regions of drug resistant and multidrug-resistant tuberculosis in the world. The rate of primary resistance in Japan was as high as the median of the world, but the rate of acquired (secondary) resistance was almost twice of the median (4). In the year 2000, the World Health Organization (WHO) registered 273000 cases of MDR-TB worldwide (5). Drug resistance has clinical and paraclinical determinations; MMM however, there is no single definition for it. In our center, a MDR bacilli should have the following three criteria: 1. Persistent positive .TANAFFOSJOURNAL.IR sputum smear in spite of ?Directory Observed Treatment Short Course? (DOTS). 2. Unchanged or progressive radiologic features. 3. Confirmed resistance at least to INH and RIF by paraclinical report. In our study, the number of cases having MDR- bacilli criteria was 38 (88%). Twenty-six (56%) patients were resistant to INH, RIF, STM, and EMB. This rate is more than that of other world medical centers which is probably due to referring patients to our center. In fact, those patients with clinical suspicion of MDR-tuberculosis had been referred to this hospital. The performed studies have shown that the majority of Mycobacterium tuberculosis strains resistant to INH and RIF are also resistant to other anti-TB drugs. Similarly in the present study, 38 cases were resistant at least to INH and RIF; 4 cases of them were resistant only to these two drugs, and the remaining had simultaneous resistance at least to one other drug. In a study by Liu et al. (6), single drug resistance to INH and STM was most common, respectively. The present study also showed that 41 (95%) and 36 (85%) cases had resistance at least to INH and STM, respectively. In a study conducted by Liu et al. (6) and other performed studies including in Ethiopia (7), single drug resistance to EMB has been very low. Similarly, minimum resistance to EMB was obtained in 26 patients (60%) in our study which is the least in comparison with other drugs. Regarding the high prevalence of resistance to STM, adding this drug in MDR-tuberculosis suspects is probably inappropriate and may induce more resistance to other concurrently prescribed drugs. Low prevalence of resistance to EMB causes its consumption as a low-cost drug in suspicion of MDR-TB suspects. In the present study, the majority of the study patients had TB with acquired drug resistance. Ninety-five percent of them had at least a single history of treatment, and most of them had responded to the standard drug regimen whom had once again in the form of MDR. As a conclusion, a single definition for MDR-TB is very necessary for control and appropriate treatment of this disease. One of the reasons for inadequate treatment is the delay in reporting drug resistance from the laboratory which is due to applying traditional methods for culture and antibiogram. Therefore, we recommend the application of

ation of www.SID.ir

## Archive of SID

5	new methods for culture and rapid testing of drug susceptibility such as BACTEC system and gene analysis (PCR) in order to achieve more rapid paraclinical diagnosis of MDR- bacilli resulting in more precise and faster treatment of the disease.
Refrences	<ol> <li>Dye C, Scheele S, Dolin P, Pathania V, Raviglione MC. Consensus statement. Global burden of tuberculosis: estimated incidence, prevalence, and mortality by country. WHO Global Surveillance and Monitoring Project. JAMA 1999; 18: 282 (7): 677-86.</li> <li>Kamholz SL. Drug resistance tuberculosis. J Assoc Acad Minor Phys 2002; 13(2): 53-6.</li> <li>Tuberculosis morbidity- United States, 1997. MMWR Morb Mortal Wkly Rep 1998; 47(13): 253-7.</li> <li>Suzuki K, Tomita M, Kinoshita Y, Yoshida A, Yoshida S, Tsuyuguchi K, et al. Present trends of drug-resistant tuberculosis and how to manage it by mycobacterial laboratories. Rinsho Byori 2002; 50(9): 847-52.</li> <li>Loddenkemper R, Brendel A, Sagebiel D, Forssbohm M. The state of tuberculosis in Germany in 2000. Pneumologie 2002; 56(9): 550-7. 6. Liu Y, Jiang G, Zhao L, Fu Y, Li Y, Bi Z, et al. Drug resistance of Mycobacterium tuberculosis in a nationwide epidemiological survey in China in the year of 2000. Zhonghua Je He He Hu Xi Za Zhi 2002; 25 (4): 224-7.</li> <li>Abate G. Drug resistant tuberculosis in Ethiopia: problem scenarios and recommendation. Ethiop Med J 2002; 40(1): 79-86.</li> </ol>

## Conclusion

Copyright 2003-2004 Tanaffos Jouranl, Email : info@tanaffosjournal.ir Designed by : Tooba Co.

www.SID.ir