

## Suicide Attempts as the Cause of Post Intubation Airway Stenosis (PIAS)

Azizollah Abbasi Dezfouli<sup>1</sup>, Shahrokh Adhami<sup>2</sup>, Roya Farzanegan<sup>1</sup>, Gholam Reza Karimi<sup>3</sup>, Mehrnaz Vali Zadeh<sup>1</sup>, Mohammad Behgam Shadmehr<sup>1</sup>, Mehرداد Arab<sup>1</sup>, Mojtaba Javaherzadeh<sup>1</sup>, Saviz Pejhan<sup>1</sup>, Abolghasem Daneshvar Kakhki<sup>1</sup>

<sup>1</sup>Department of Thoracic Surgery, NRITLD, Shaheed Beheshti University of Medical Sciences and Health Services, <sup>2</sup>Department of Thoracic surgery, Kurdistan University of Medical Sciences and Health Services, <sup>3</sup>Department of Psychiatry, Baqiyatallah Medical Sciences University, TEHRAN-IRAN.

### ABSTRACT

**Background:** The incidence of suicide attempt has been increasing in recent years. Presenting a group of patients who attempted suicide, underwent ventilatory support and developed postintubation airway stenosis (PIAS) may help us in prevention and better understanding of this complication.

**Materials and Methods:** Among patients who referred to our center for treatment of PIAS, those who had been intubated for suicide attempt were investigated in a prospective study. Information was entered in a questionnaire and regular follow ups were done in a 15-month period (April 2003 to July 2004).

**Results:** Among 100 patients with PIAS, 19 enrolled in this study including 10 females and 9 males (mean ( $\pm$ SD) age, 25.3 ( $\pm$ 9.96) yrs; ranging from 17 to 56 yrs). Type of disease and reasons of suicide were categorized by a psychologist as follows: Eleven patients with psychosocial stress along with an immature personality back-ground, 7 cases of psychological disorders and one with an unknown cause.

Direct causes of committing suicide included family problems in 10 cases, lovesick in 2, addiction in 3, depression in 6 and social problems in 2 cases (some patients mentioned two reasons and one refused to mention the reason).

Mean time of intubation was 14.78 days (3-30 days), and the mean length of stenosis was 35.12 mm (20-50 mm), 8 patients underwent tracheostomy. Three patients were treated with bronchoscopic dilation and 16 underwent laryngotracheal resection and reconstruction. There were 8 cases of recurrence after resection among which 4 were treated by second resection, 2 recovered by bronchoscopic dilation and 2 managed by stenting. This group of patients (study group) was compared with a similar group of patients in whom the causes of intubations were different (control group). Incidence of post-surgical recurrence ( $p=0.011$ ) and the length of stenosis ( $p=0.01$ ) were higher in the study group.

**Conclusion:** In our patients, social problems such as unemployment, illiteracy and singleness were the more frequent causes of suicide compared with psychological disorders. Patients who undergo mechanical ventilation due to suicide and develop PIAS could be treated by tracheal resection and reconstruction; although the incidence of post-surgical recurrence is higher in them compared with the other groups of patients with PIAS. (Tanaffos 2005; 4(15): 11-16)

**Key words:** Suicide, Post Intubation Airway Stenosis (PIAS), Suicide attempt

Correspondence to: Abbasi- Dezfouli A

Address: NRITLD, Shaheed Bahonar Ave, Darabad, TEHRAN

19569, P.O:19575/154, IRAN

Email: abbasidezfouli@nritld.ac.ir

## INTRODUCTION

Suicide has a history as old as mankind. Unfortunately, incidence of suicide attempts has been increasing in recent years (1, 2, 3, 4, 5, 6).

Based on World Health Organization (WHO) report, at least one million people in the world commit suicide annually (2). Suicide is the 8<sup>th</sup> cause of death in the United States of America, 2<sup>nd</sup> in 15-19 yrs old males residing in this country (7) and 10<sup>th</sup> in the world (1). Thus, suicide has been converted from an individual abnormality to an extensive important social obstacle. It is a social health problem which needs precise programming for its prevention (1, 8, 9).

Psychiatrists believe that suicide-related factors are different between men and women in different ages, various jobs, and different educational levels (10).

Furthermore, individuals commit suicide often as the result of psycho-somatic disorders. Among patients with psychological diseases who attempt suicide, some have depression and some are schizophrenic (which their hallucinations cause suicide attempt). Other psychological disorders are also responsible for suicide attempts to various degrees (11, 12). Some people attempt suicide because of their diseases (either psychotic or somatic); on the other hand, there are healthy ones with other types of problems. The latter group turned suicide into a big social problem (12).

Referring some patients with PIAS resulting from suicide attempts to our center, made some questions come to the physicians' minds as follows:

- 1- Is the possibility of PIAS higher in patients who attempt suicide?
- 2- Has airway care been insufficient in these patients while being under mechanical ventilation?

Presence of tracheal stenosis causes numerous difficulties for patients and their families and can also impose severe psycho-economical effects on family and society.

Since these patients formed a nearly harmonious group based on family and social aspects and related treatments, we studied them in a group as a descriptive study and compared them with a similar group of patients with PIAS to answer the

abovementioned questions.

## MATERIALS AND METHODS

This was a prospective study including individuals who had been referred to our center between Apr 2003 and Jul 2004 because of PIAS due to suicide attempt. Information was entered in a questionnaire and regular follow-up was performed. The questionnaire contained various questions including psychological status, cause of suicide attempt, drug history, family history, economical state of patients' parents, marital status, the type of treatment and its results. The patients' follow-ups were done actively via face to face interview and by phone. Details are summarized in table 1. The results were compared with those of a similar group of patients with PIAS which was not due to suicide attempt.

## RESULTS

The study patients were 10 females and 9 males with age range of 17-56 yrs; ( $25.3 \pm 9.96$  yrs). Only five of them were married (all of them were women). The educational level of these patients was as follows: one person was university student, 6 had high school diploma, and the remaining 12 had under diploma degree. Eight patients were unemployed. Based on psychologist's diagnosis, cause of suicide in 11 cases was psychosocial stress with an immature personality background, in 7 was psychological disorder and in one the cause was unknown.

Direct causes of suicide were family problems in 10 cases, lovesick in 2, addiction in 3, severe depression in 6, and social problems in 2 cases (in some patients there was more than one cause). None of them was the first offspring in their family and only five cases were under psychiatric treatment before attempting suicide.

Also, 5 patients had previous history of suicide attempt. The mean duration of intubation was 14.78 days (ranged 3-30 days). Eight patients had undergone tracheostomy procedure before being referred to us. Site of stenosis was the trachea in all patients and mean length of stenosis was 35.12 mm (20-59 mm).

Archive of SID

The procedures were included laryngotracheal resection and reconstruction (resection anastomosis), bronchoscopic stenosis dilation and polyflex stenting. Three patients were treated with bronchoscopic dilation out of which one had undergone resection anastomosis in another center. In 16 cases, resection anastomosis was done (one patient had been operated in another center and was referred due to recurrence of post- surgical stenosis); consequently, 6 of them were involved with post-surgical stenosis recurrence (totally 8 patients).

Among these 8 patients, 4 underwent resection anastomosis again and 2 underwent stenosis dilation by rigid bronchoscopy. As a result, 6 of them were recovered. Stent was placed for the remaining 2 other cases. Only one patient had recurrent laryngeal nerve paralysis as a permanent complication. The statistical significance of various factors in our patients is shown in table 2.

**Table 2.** Statistical significance of different factors related to patients with attempting suicide.

Factor		p-value
Age	Ranging 17-35 yrs (mean $\pm$ SD), 25.3 $\pm$ 9.96	
Sex	Female 52.6% Male 47.4%	p>0.05
Marital Status	Unmarried 73.7% Married 26.3%	P<0.05
Job	Unemployed 42.1%	
Educational level	Below diploma 63.1% Diploma and higher 36.9%	P>0.05
Suicide method	Pill 57.9% Poison 36.8% Opium 5.3%	P<0.05
Second offspring to next	second offspring to next 100% The first offspring 0	
Cause of suicide based on psychiatrist's diagnosis	Psychosocial stress 61.1% Psychiatric disorder 39.9%	P>0.05
Psychiatric disorder	Depression 71.4% Other disorders 28.6%	
History of suicide attempt	Negative 66.7% Positive 33.3%	

This group of patients (case group) was compared to similar group (control group) with PIAS which was not due to suicide attempt. Results are summarized in table 3. Of 55 patients in the control group, one had left vocal cord paralysis; one had right vocal cord paralysis, and one had fixed vocal cord. In addition, one patient died due to sudden cardiac arrest. In his history, he had heart failure and experienced several heart attacks before surgery.

**Table 3.** Comparison of patients group (who attempted suicide and involved with PIAS in Masih Daneshvari Hospital) with control group (who were intubated due to other causes and referred to this center simultaneously with the patients group).

	Control group	suicide group	p-value
Total	81	19	
gender			
Female	15	10	
Male	66	9	
Age range and mean (Yr)	17-66 24.42	17-56 25.32	
Subglottis stenosis	2	0	
Mean duration of intubation and range(day)	15.58 0-40	14.78 3-30	
Mean length of stenosis and range(mm)	28.28 10-60	32.72 20-50	P=0.011
Treatment			
resection-anastomosis	RA=55	RA=16	
bronchoscopy dilation	B-D=23	B-D=3	
Stent	S=2	S=0	
laser	L=1	L=0	
Post surgical recurrence in this center	5 55	6 16	P=0.011

## DISCUSSION

1. **Age:** mean ( $\pm$ SD) age of these patients was 25.3 ( $\pm$ 9.96) yrs. In fact, most of them were between 17 and 35 years old and only 3 patients were over 35 years. In a study conducted in Dubai, the age

range was 21-40 years (13). Another study showed that increasing the age results in increased suicide rate. Suicide rate seems to increase in men over 45 years old and women over 55 years of age (14). In contrast, we had only one female patient over 55 years old; additionally, suicide age range in men was under 37 years.

2. **Marital Status:** only five patients were married. The effect of marriage on decreasing the suicide rate in our patients was significant ( $p < 0.05$ ). Suicide incidence in unmarried or single persons is twice the married ones (14). Thus, it seems that marriage and giving birth significantly reduce suicide incidence (11, 14, 15). In a survey conducted in the Lorestan province (Iran), 40 suicides were reported by eating chalk between 1995 and 2000. Most patients were single (52.17% single women and 88.2% single men) (16).
3. **Sex:** most patients in our study were men. In two surveys conducted in Oulu state in North Finland (17) during 1998-2000, and Dubai (13) during 1992-2000, the majority of patients attempting suicide were male. Suicide rate was also high in men in Highland of Scotland (18). It was also true in the USA (14). In contrast, of 40 patients who were studied in the Lorestan province, 57.5% were women and 42.5% were men (16).
4. **Occupation/ Employment:** our study patients were consisted of 5 housewives, one medical student, one butcher, 3 students, one businessman and the others were unemployed. Unemployment rate did not show statistical significance as a suicidal cause. However, 42.1% of patients were unemployed. Numerous studies have shown employment as a protective factor against suicide (12, 14). In other words, there was a strong relationship between unemployment and suicide (11, 19, 20, 21, 22, 23, 24). In our study, most patients were unemployed. In a study performed on 40 patients in the province of Lorestan (16),

37.5% were students, 35% were housewives and 10% were unemployed. In the present study, these percentages were 15.8%, 26.3% and 42.1%, respectively.

5. **Suicide methods:** several methods have been reported for suicide, however, selecting these methods depends on cultural, geographical and socio-economical conditions. Among our study patients, pills, poison and opium had been used in 11, 7 and 1 cases, respectively. Drug consumption showed statistical significance ( $p < 0.05$ ). Some of these drugs were indral, phenobarbital and benzodiazepines. Hanging was the most common method of suicide in Dubai (13) and the USA (14). A survey performed in South-West of France (Aquitaine), showed hanging, drowning, and drug use as the most common methods of suicide (25). Hanging is one of the most common methods of suicide in the world (26). In a survey conducted in South Africa, hanging, use of firearms and self-poisoning consisted 57%, 30% and 13% of suicide attempts respectively (27). Suicide attempt by eating chalk is a strange method which has been reported only in Lorestan, Iran (16).
6. **Reason of suicide attempt:** one of the important risk factors in Denmark study (1981-1997) was psychological disorder or history of hospitalization due to that (21). Other studies have considered mood disorder as one of the risk factors for suicide attempt (11). Epidemiological studies in Lorestan province showed a higher frequency of psychological disorders in subjects in the age range of 26-40 years compared with those in the age range of 18-25 yrs. Some factors such as environmental stress, repeated child births, anxiety due to biologic alterations in women, occupational and economical problems, unemployment in men, divorce and high education cause high frequency of psychiatric disorders in these ages (28). However, in our

study only in 7 patients the suicide attempt was as the result of recognized psychiatric disorder and psychosocial tensions were the cause of suicide in the others. Psychological disorders in these 7 cases were mostly seen in the age group of 26-37 years compared with those less than 25 yrs; 57.14% were unemployed and 85.71% were male. It seems that attention must be paid to decrease these stresses in young adults. 39.9% of our study patients who went to coma due to suicide attempt had psychiatric disorder of which 71.4% had depression based on psychiatrist's diagnosis.

7. **Cause of stenosis:** the most important cause of PIAS is long duration of intubation and poor nursing care of airways during this period (29). This length of time in our patients was not different from other reports but we assume that the patient's care has not been appropriate. A hospitalized patient with suicide attempt may be ignored by his or her family and physician, or perhaps another cause might be responsible in this issue. One of the important differences between our study and other ones is using pills, especially diazepam and phenobarbital, for suicide by our patients. We suppose that maybe these drugs or those used to control symptoms like convulsion are responsible for stenosis. All study patients had used drugs or poisons for suicide attempts. Perhaps using these drugs or application of some methods for detoxification is responsible for causing scar and endotracheal fibrosis; however, evaluation of this hypothesis is difficult. Anti-convulsant drugs, particularly dilantin, were prescribed for our patients. It is believed that dilantin aggravates scar formation. This capability has been used for healing of bed sores (30-33). However, the effectiveness of these drugs is not known in our patients. Although the majority of study patients had been hospitalized in academic medical centers and emergency wards, this rate of PIAS seems unusual; however, precise statistical

judgement is not possible.

We assume that admission overload of these patients can decrease the caring level and impose an unbearable pressure on hospital personnel. It is suggested that those patients with acute respiratory failure due to drug poisoning should be transferred to centers with special facilities and capability of ventilation support one or two days after the admission.

## REFERENCES

1. EBR Desapriya and Iwas N. New trends in Suicide in Japan. *Inj Prev* 2003; 9: 284. 2003 BMJ Publishing group.
2. Polewka A, Kroch S, Chrostek Maj J. Suicidal behavior and suicide attempts in adolescents and young adults--epidemiology, risk factors, prevention and treatment. *Przegl Lek* 2004; 61 (4): 261- 4.
3. Jakuboniene D, Gurevicius R. Mortality from external causes: influence of age and gender. *Medicina (Kaunas)* 2003; 39 (8): 773- 87.
4. Jianlin J. Suicide rates and mental health services in modern China. *Crisis* 2000; 21 (3): 118- 21.
5. Neeleman J. [Suicidality]. *Ned Tijdschr Geneeskde* 2001; 145 (11): 515- 20.
6. Hafner H. The epidemiology of suicide and attempted suicide. *Psychiatr Neurol Med Psychol (Leipz)* 1989; 41 (8): 449- 75.
7. Vahhabzadeh J. Main discussions in psychology, Tehran, Scientific and cultural publications, 1993; 8-15.
8. De Leo D. Struggling against suicide: the need for an integrative approach. *Crisis* 2002; 23 (1): 23- 31.
9. Disley B, Coggan C. Youth suicide in New Zealand. *Crisis* 1996; 17 (3): 116- 22.
10. Kaplan and Sadok (2000) comprehensive text book of psychiatry, vol 3, suicide 92, Williams and wilkins. P: 2031- 2045.
11. Welch SS. A review of the literature on the epidemiology of parasuicide in the general population. *Psychiatr Serv* 2001; 52 (3): 368- 75.
12. PoorAfkari N. Technical psychology, clinical psychology, First Edition, Shahrab 1996; 1-20.

13. Koronfel AA. Suicide in Dubai, United Arab Emirates. *J Clin Forensic Med* 2002; 9 (1): 5- 11.
14. Suicide in synopsis of Psychiatry/ Behavioral sciences/ clinical. Psychiatry by KAPLAN and Sadock's Lippin Cott Williams and wilkins (2003)- 9<sup>th</sup> edition – (913-923).
15. Stack S. The effect of marital integration on African American suicide. *Suicide Life Threat Behav* 1996; 26 (4): 405- 14.
16. Evaluation the cases of chalk/plaster eating as a new and pdd method of suicide in Lorestan province, *Yafteh Journal*, 5<sup>th</sup> volume, winter 2003, n:19.28-32.
17. Timonen M, Viilo K, Hakko H, Sarkioja T, Meyer-Rochow VB, Vaisanen E, et al. Is seasonality of suicides stronger in victims with hospital-treated atopic disorders? *Psychiatry Res* 2004; 126 (2): 167- 75.
18. Stark C, Matthewson F, O'Neill N, Oates K, Hay A. Suicide in the Highlands of Scotland. *Health Bull (Edinb)* 2002; 60 (1): 27- 32.
19. Lewis G, Sloggett A. Suicide, deprivation, and unemployment: record linkage study. *BMJ* 1998; 317 (7168): 1283- 6.
20. Platt S. Unemployment and suicidal behaviour: a review of the literature. *Soc Sci Med* 1984; 19 (2): 93- 115.
21. Qin P, Agerbo E, Mortensen PB. Suicide risk in relation to socioeconomic, demographic, psychiatric, and familial factors: a national register-based study of all suicides in Denmark, 1981-1997. *Am J Psychiatry* 2003; 160 (4): 765- 72.
22. Voss M, Nylen L, Floderus B, Diderichsen F, Terry PD. Unemployment and early cause-specific mortality: a study based on the Swedish twin registry. *Am J Public Health* 2004; 94 (12): 2155- 61.
23. Gibb S, Beautrais A. Epidemiology of attempted suicide in Canterbury Province, New Zealand (1993-2002). *N Z Med J* 2004; 117 (1205): U 1141.
24. Fuller GN, Rea AJ, Payne JF, Lant AF. Parasuicide in central London 1984-1988. *J R Soc Med* 1989; 82 (11): 653- 6.
25. Maurice S, Pommereau X, Pueyo S, Toulouse C, Tilly B, Dabis F, et al. Epidemiological surveillance of suicides and attempted suicides in Aquitaine, south-west France, using an original computer network of sentinel general practitioners. *J Epidemiol Community Health* 1989; 43 (3): 290- 2.
26. Gunnell D, Bennewith O, Hawton K, Simkin S, Kapur N. The epidemiology and prevention of suicide by hanging: a systematic review. *Int J Epidemiol* 2005; 34 (2): 433- 42. Epub 2005 Jan 19.
27. Meel BL. Determinants of suicide in the Transkei sub-region of South Africa. *J Clin Forensic Med* 2003; 10 (2): 71- 6.
28. Mohammadi M, MalekAfzali H, Noorbala A, Seyedbagheri Yazdi A, Rahgozar M, Mesgarpour B, Safa M, Radfar R. Epidemiology of psychology disorders, in Lorestan province. *Yafteh Journal* 2001; 5<sup>th</sup> volume, N .19.
29. Grillo HC, Donahue DM, Mathisen DJ, Wain JC, Wright CD. Postintubation tracheal stenosis. Treatment and results. *J Thorac Cardiovasc Surg* 1995; 109 (3): 486- 92; discussion 492- 3.
30. Bhatia A, Prakash S. Topical phenytoin for wound healing. *Dermatol Online J* 2004; 10 (1): 5.
31. Pitiakoudis M, Giatromanolaki A, Iliopoulos I, Tsaroucha AK, Simopoulos C, Piperidou C. Phenytoin-induced lymphocytic chemotaxis, angiogenesis and accelerated healing of decubitus ulcer in a patient with stroke. *J Int Med Res* 2004; 32 (2): 201- 5.
32. Bhatia A, Nanda S, Gupta U, Gupta S, Reddy BS. Topical phenytoin suspension and normal saline in the treatment of leprosy trophic ulcers: a randomized, double-blind, comparative study. *J Dermatolog Treat* 2004; 15 (5): 321- 7.
33. Scheinfeld N. Phenytoin in cutaneous medicine: its uses, mechanisms and side effects. *Dermatol Online J* 2003; 9 (3): 6.