

*Tanaffos* (2007) 6(2), 93-95

©2007 NRITLD, National Research Institute of Tuberculosis and Lung Disease, Iran

### Diagnosis: *Pneumocystis Jirovecii* Pneumonia

Chest X-ray demonstrated bilateral fluffy alveolar densities in midzone of the lungs and parahilar peribronchial infiltrations were also seen (Figure 1).




Figure 1. Bilateral diffuse infiltration in 1<sup>st</sup> chest X-ray

High resolution computed tomography (HRCT) of the lung revealed bilateral nonhomogeneous ground glass opacities in mid and upper zones of the lungs. Mild bilateral peribronchial wall thickening was also seen (Figure 2). These findings were suggestive of *Pneumocystis Jirovecii* Pneumonia (1, 2).

Smear of bronchoalveolar lavage (BAL) for pneumocystis was suspicious. Transbronchial biopsy revealed mild distortion of alveolar architecture and prominent intra-alveolar deposition of foamy

eosinophilic exudates. Immuno-histochemistry staining with monoclonal antibody was strongly positive for *Pneumocystis* (Figure 3).




Figure 2. Extensive ground glass attenuation on lung HRCT




Figure 3. Immunostaining showing clusters of organisms in the alveoli

Acid fast staining of sputum and BAL was negative, but PCR for cytomegalovirus (CMV) was positive in BAL.

The serology for human immunodeficiency virus (HIV) including both enzyme-linked immunosorbent

assay (ELISA) and Western blot reported to be positive. Absolute CD4+ T cell count was 6/ $\mu$ l. The patient was interviewed in detail and acknowledged that he had intravenous drug abuse for a short period ten years ago with sharing syringes.

Pneumocystis pneumonia (PCP) is often the acquired immunodeficiency syndrome (AIDS)-defining illness in patients infected with HIV, occurring most frequently when the T-helper cell count (CD4+) is less than 200 cells per cubic millimeter.(1) Incidence of PCP has declined substantially with widespread use of prophylaxis and effective antiretroviral therapy (ART). The majority of cases occur among patients who are unaware of their HIV infection or not receiving ongoing HIV care or those with advanced immunosuppression(3).

Most people infected with HIV are not recognized in countries such as Iran and PCP may be the first presentation of their HIV infection.

Isolation of CMV from BAL specimens is relatively common, yet most of these patients have an alternative diagnosis (especially pneumocystis or bacterial pneumonia) and may improve without specific therapy directed at CMV. Nonetheless, in a patient with advanced HIV disease, interstitial infiltrates on chest X-ray, and no established alternative diagnosis, CMV may be the sole responsible pathogen. (4) This diagnosis is confirmed often when histopathology demonstrates intracellular inclusions typical of CMV (5), which was not seen in our patient.

The patient was treated with thrimethoprim-sulfamethoxazole (TMP/SMX) 15 mg/kg/day (TMP) and prednisolone 40 mg orally twice daily. After 2 days, his condition improved dramatically and the treatment was continued for 3 weeks with tapering of steroid.

Chest X-ray on the 6th day of treatment showed marked improvement (Figure 4).

Antiretroviral therapy (ART) with zidovudine, lamivudine and nevirapine and weekly azithromycin (1250 mg) for the very low CD4 count were initiated and the TMP/SMX was continued in a prophylactic dose.




Figure 4. Marked improvement in 6<sup>th</sup> day chest X-ray

In conclusion, we recommend considering HIV infection in differential diagnosis of all patients with history of opium addiction, even if the patient denies any intravenous abuse, particularly when they present with opportunistic infections such as Pneumocystis.

## REFERENCES

1. Thomas CF Jr, Limper AH. Pneumocystis pneumonia. *N Engl J Med* 2004; 350 (24): 2487- 98.
2. Rossi SE, Erasmus JJ, Volpacchio M, Franquet T, Castiglioni T, McAdams HP. "Crazy-paving" pattern at thin-section CT of the lungs: radiologic-pathologic overview. *Radiographics* 2003; 23 (6): 1509- 19.

3. Benson CA, Kaplan JE, Masur H, Pau A, Holmes KK; CDC; National Institutes of Health; Infectious Diseases Society of America. Treating opportunistic infections among HIV-infected adults and adolescents: recommendations from CDC, the National Institutes of Health, and the HIV Medicine Association/Infectious Diseases Society of America. *MMWR Recomm Rep* 2004; 53 (RR- 15): 1- 112. Erratum in: *MMWR Morb Mortal Wkly Rep* 2005; 54 (12): 311.
4. Walzer PD, Smulian AG. Pneumocystis Species. In: Mandell GL, Bennett JE, Dolin R. editors. Principles and practice of infectious diseases. 6<sup>th</sup> Ed, 2005, Churchill Livingstone, 1567-73.
5. Uberti-Foppa C, Lillo F, Terreni MR, Puglisi A, Guffanti M, Gianotti N, et al. Cytomegalovirus pneumonia in AIDS patients: value of cytomegalovirus culture from BAL fluid and correlation with lung disease. *Chest* 1998; 113 (4): 919-23.

Archive of SID