

## Evaluation of fruits cultivar and different harvest times of Damghan pistachio to early split complications and contamination to the *Aspergillus flavus* fungus

Amini F.<sup>1</sup>, Afshari H.<sup>2</sup>, Mohammadi Moghaddam M.<sup>3</sup>

<sup>1</sup>Graduate student, Damghan Branch, Islamic Azad University, Damghan, Iran

<sup>2</sup>Assistant Professor, Damghan Branch, Islamic Azad University, Damghan, Iran

<sup>3</sup>Researcher of Damghan Pistachio Research Station

**Abstract:** Study the effect of fruit cultivar (intact, early split, irregular crack) on (Akbari, Abbasali, Khanjari) cultivars in Damghan city to *Aspergillus Flavus fungus* and effect of harvest time in *early split pistachio* in 1389, sampling from orchards has been performed. Samples after grinding by dilute cultivating in a specific AFPA medium bottles in a completely randomized model were three times recultivated. Colonies of *A. flavus* after 3 to 7 days, were isolated, identified and counted. The results of Duncan grouping were analyzed by SAS software. Results of that study of contamination of samples to *A. Flavus* fungal Showed The contamination of the samples to the fungus *A. Flavus* differ from each other, there is less contamination in the Khanjari cultivar and more contamination in Akbari cultivar. With delay in the harvest time, percentage of intact fruits and without front skin cracking decrease. The lowest percentage of early split (0 first) was at first harvesting time (Shahrivar 10) in Akbari cultivar and the highest percentage of *early split* (Shahrivar 20) in the the second harvest time in the Akbari cultivar.

**Keywords:** *Aspergillus flavus*, cultivar, early split, harvesting times

### INTRODUCTION

Pistachio fruit is sensitive to contamination by Fungi such as *Aspergillus flavus* and *Aspergillus parasiticus* that produce *Aflatoxins*. Fruit skin cracking reduce the fruit quality that during harvest and after harvest will reduce fruit skin quality and affect its market friendliness [3].

Fruit cracking may occur when the fruit is still attached to the tree. Fruit cracking in stone fruit, seeded fruit, grapes and vegetables has been seen. skin cracking and surface cracks of fruit may be the result of a number of factors including water stress, insect, spraying.

Early split pistachios, are abnormal Pistachio nuts that both green skin and bony skin, simultaneously crack at the split place. In other words, cracks created on the Green skin, exactly coincide with the

split of bony skin and so the kernel of pistachio is visible. In this cultivar of cracks because the edible part of fruit (the kernel) is directly exposed to the attack of spores and *fungi*, is one of the most important factors in *Aflatoxin* production in orchards [1].

These Pistachios are abnormal because the. hard skin split occurs before maturity.(Prior to separation of the, soft skin from, bony skin), so in this case the soft skin is attached to the bony skin and does not have the necessary flexibility. By premature Splitting of hard skin the, soft skin also crack from the hard skin split spot. This condition is called early split [2].

When the harvest time for Pistachio nuts draws near, *Aflatoxin* generator *fungus* population density increase within orchards and it appears to be a

\*Corresponding author: Fatemeh Amini, Graduate student, Damghan Branch, Islamic Azad University, Damghan, Iran Email: aminiba.43@gmail.com

correspondence between the increase of the density of spores of this fungus and the growth of Pistachio nuts fruit [12].

Mojtahedi and colleagues in 1979 reported the Iran's Pistachio nuts orchard fruits contamination [6].

The *Aspergillus flavus* fungus group and *Aspergillus niger* population increases by approaching harvest time due to the increased abundance of cracked poured Pistachio nuts on the orchard floor and contamination increase by contact with the fungus mentioned above. We can also mention more time exposure to the spores in the atmosphere of Pistachio nuts orchards. Timely and Early harvesting reduce *Pistachio nuts* fruit contamination and also reduce contamination of new *Pistachio nuts* [8].

Fruit growth of Pistachio nuts in orchards is associated with soft skin and bony skin. When the fruit matures the soft skin separate from the Hard skin. And cracking of Hard skin is important for Fruits market friendliness [6].

The purpose of this study is assessing the effect of fruit cultivar (early split, irregular cracks and intact) and different times of harvesting phenomenon on early split of Pistachio nuts on (Akbari, Khanjari and Abbasali), cultivar.

## MATERIALS AND METHODS

**Sampling:** In order to study the extent of contamination of Pistachio nuts of Semnan province to *A.flavus* group of *fungus aflatoxin* different parts of Damghan city Pistachio nuts cultivation of fresh pistachios in 1389 were sampled.

From late Mordad, coincide with the product maturation, we select the orchards and collect samples of three varieties of Pistachio nuts (Akbari, Abbasaly Khanjari). of 1.5 kg from each cultivar, in

sampling with interval of 10 days during the three phases, at each phase the percentage of nut side pistachios (intact, early split and irregular cracks) are calculated and after separation of this Pistachio nuts from each other we separate the green skin and sun-dried them.

Since the distribution of toxic and contamination dispersion in different parts of the kernel of pistachio is different and estimating the extent of fungal *A.flavus* infection and *aflatoxin* requires achieving a quite uniform and homogeneous sample, pistachio samples are grinded. Grinded samples by dilute cultivating method cultured on surface environment of *Aspergillus flavus parasiticus* Agar (AFPA) medium. (Completely randomized design with three replications). Fungi of This group on the AFPA specific medium produce Conidiales of typical yellow to olive green.

From each cultivar the amount of 10 gr kernel of pistachio in diluted method was added to 90ml peptone water 1% and then diluted to  $10^{-1}$ , 0.01 ML of above sample were cultured on surface of AFPA medium petri dish. Petri dishes were kept at  $26^{\circ}\text{C}$ .

After 3 to 7 days of cultivation, colonies of *A. flavus* were counted and separated, the contamination in various samples were compared. Obtained Data using SAS software analyzed and compare with Duncan test was performed at 5% level.

## RESULTS AND DISCUSSION

Results of *A. flavus* colony counts is shown in Table 1. As can be seen in Table the results from these *A.flavus* colonies suggest that The fungal contamination of the samples of kernel of pistachio is different for various Pistachios. As such in some cases there was no fungal contamination of *A.flavus* and some samples showed high contamination of this fungus. (Table 1). Evaluating the result

(Duncan's multiple range tests) is indicated the difference between that average numbers of fungal colonies different samples of *Pistachio nuts* in 1% statistical level is of significance. (Table 2) The results of these investigations show the delay in the time of harvest reduces percentage of intact, and un-cracked front skin fruits .this difference between the times of first (Persian date Shahrivar 10), and

second (Persian date Shahrivar 20) and third (Persian date Shahrivar 30) is quite significant.

Effects of harvest time on the percentage of the early split product showed the lowest percentage of early split is zero in the first harvest (Persian date Shahrivar 10) in Akbari cultivar, and most early split of Pistachio nuts 7.5% in the second harvest (Persian date Shahrivar 20) in Akbari cultivar.

Table 1: Comparison of pistachio samples colony numbers of fungi *A.flavus* of Damghan

No.almond sample	The average number fungal colonies
1	0
2	$5.7 \times 10^2$
3	$2.1 \times 10^2$
4	0
5	$1.33 \times 10^2$
6	$6.3 \times 10^2$
7	0
8	0
9	$0.333 \times 10^2$
10	$.066 \times 10^2$
11	$1.6 \times 10^2$
12	$2.3 \times 10^2$
13	0
14	$1 \times 10^2$

Percentage difference of early split between the second and third times with first time means the early harvesting can be effective in reducing contamination of pistachio *A.flavus* fungi.

Assessment of different harvest time effects showed that the delay in harvesting increases the fungi *Aspergillus flavus* contamination of the product. (Figure 2)




Fig 1. Means of *A.flavus* fungi in different pistachio cultivars




Fig 2: Means of *A.flavus* fungi in harvesting time

The results showed that the delay in harvesting reduces the percentage of intact fruit (without cracking the green skin) Intact skin of fruits until harvest is important because as long as the front skin is intact it acts as a deep barrier against penetration of external agents such as insects, fungal spores and dust and also helps moisture loss of the fruit

In this study, the delay in harvest increase the fruit early split. Shatzky and colleagues (1997) reported that the early split is the main cause of fungal contamination and *aflatoxin* production.

The results of this study showed that the highest fungus *Aspergillus flavus* contamination occurs in the last harvest, and least contamination in the first harvest. In other words, harvest at the first weeks after physiological maturity of product, reduced

product contamination to fungus *Aspergillus flavus* also the delay in the harvest increases tarnished pistachio.

Evaluation the effects of different harvest times, showed that the contamination in the first harvest has a significant difference with the second and third times of harvest. The test results is consistent with the results of Taj Abadi Pur (1383) study about the effects of harvest time.

Results of research on Kerman Ohadi Pistachio shows that best time of harvest for highest weight dry product occurs in the first week of Mehr and maximum percentage of early split fruit occurs in the third week of Mehr. Most productive fruits occur at the fourth week of Mehr and least productive fruits occur at second week of Shahriwar [4].

#### REFERENCE

1. Taj Abadi Pur,A;1999. Early split complications and its percentage in various pistachios.

Pistachio Research Institute Publications No. 77/450, 15

2. Taj Pur Abadi,A; And M. Fooladi; 2005. Identification and determination of aflatoxin contaminated pistachio in processing stage. Pistachio Research Institute's final report. Rafsanjan 0.16
3. Syahkvhy,B;2010. The effects of concentration and time of spraying of gibberellic acid on the reduction of green pistachio shell cracking percentage, Journal of Horticultural Science
4. Crane ,J.C. 1978. Quality of *pistachio nuts* as affected by time of harvest. J.Amer.Soc.Hort.Sci.103 (3):332-333.
5. Doster;M.A.and T.J.Michailides.1995 the development of early split pistachnuts and the contamination by molds *Aflatoxine* and insect .first international symposium on pistachionuts.20-24 September 1994.Adana Turkey. *Acta Horticulturae* 419: 359-364.
6. Afshari.H; 2009. Differences in nutrient in early split and healthy pistachio's green skin and phenolic and Galic acids effect in green skin on the *aflatoxin* toxin, Journal of Horticultural Science.
7. Basirat,M; 2003. Preventing effect of Vapor Guard in *pistachio nuts* cracking . The work progress report from 1379 and 1380
- 8:M.Moradi,. 2003. Study of Accumulation of aflatoxin generator factors to determine the starting point of contamination in pistachio production process and its control. The final report of Pistachio Research Institute of Iran.
9. Moradi ,M; Taj Abadi Pur,A; and B. Panahi 2004. The early the date pistachio splitting relationship And changes in their physical properties and aflatoxin contamination in three Commercial varieties in Iran (Ohadi, Ahmdaqayy and Kaleghoochy). the final report of Pistachio Research Institute of Iran.
10. Moradi, M., and H. Masoomi . *Aspergillus fungus* in pistachio orchards in Kerman province. Selected Proceedings of the first academic - applied conference of Sirjan Pistachio. Page 68.. 1381. Fertilizers Role in plant and animal population density
11. Mohammadi Moghaddam,M; 2000, evaluation of sensitivity of pistachios to fungus *Aspergillus flavus*, master's thesis, Department of Plant Protection of Agriculture Faculty, Tarbiat Modarres University, page 140.
12. Rafizadeh, M;2000. Determination of the best harvest time of pistachio nuts to optimize product quality. final report of Pistachio Research Institute. 27 page.
- 13.Kader , A.A ;C.M.Heintz;J.I.Labavitch and H.I.Rae.1982.Studies related to the description and evaluation of *pistachio nut* quality. J.Amer.Soc.Hort.Sci.107:812-816.
- 14.Yazdanpanah ,H; *Mycotoxine* contamination of food and Feedstuffs in Iran.*journal Of pharamaceutical Kesearch*(2001)17:21-30
15. Z. PakKish, 2003. Effects of gibberellic acid and Benzyladenine spray on the reduction of *aflatoxin* and cracking of pistachio nut's green shell and fungus penetration into dried fruits. MS Thesis Department of Horticulture section, Agricultural faculty of Shiraz University.