

## The Effect of Salicylic Acid and Zinc on Inflorescence Bud Abscission, Leaf Chlorophyll Florescence and Characteristic of Pistachio Nut Cv. Owhadi

H.R. Karimi<sup>1\*</sup>, S. Sarmadi-Far<sup>1</sup>, B. Panahi<sup>1</sup>

<sup>1</sup>Department of Horticultural sciences, College of Agriculture, Vali-e-Asr University of Rafsanjan, Rafsanjan, Kerman, Iran

**Abstract:** In order to determine effect salicylic acid and Zn on characteristics of pistachio nut (*Pistacia vera* L. cv. Owhadi), the experiment was carried out as factorial in the framework of CRD design with three replications. The results showed that salicylic acid and Zn increased nut fresh weight and decreased blank nut percentage. The application of salicylic acid decreased splitting percentage and kernel dry weight. Simultaneous application of 250 mg L<sup>-1</sup> SA and 0.2% Zn (Zn<sub>2</sub>SO<sub>4</sub>) decreased inflorescent bud abscission at compared to control.

**Keywords:** Salicylic acid, *Pistacia vera*, Bud abscission, Chlorophyll florescence

### INTRODUCTION

Pistachio (*Pistacia vera* L.) a member of Anacardaceae family is an important and exportable nut crop of Iran cultivated since old times. At this time, Iran is the leading producer of this crop followed by USA, Turkey and Syria. The majority of the pistachio production areas is located in arid and semi-arid regions adjacent to desert regions. Low irrigation water quality, soil salinity, nutrient imbalance and heat and drought stress are the most limiting factors in these areas. Under environmental stress condition, reactive oxygen species (ROS) produced in plant. In order to avoid the harmful effects of ROS, plants evolve an effective scavenging system including of non-enzymatic antioxidants and enzymatic antioxidants [1].

Salicylic acid (SA) is considered as a hormone-like substance, which plays an important role in the regulation of plant growth and development, seed germination, fruit yield, rooting of cuttings and

resistance to abiotic stresses [2,3]. Ion uptake and transport [4] photosynthetic rate, stomata conductance and transpiration [5] could be affected by SA application. Exogenous SA could regulate the activities of antioxidant enzymes and increase plant tolerance to the abiotic stress [6]. Moreover, the effects of SA on reducing plant abiotic stress have reported by numerous reports [1, 7, 8, 9, 10, and 11].

Many reports were showed that Zn has an antioxidant role in plant under conditioning stress. Cakmak (2000) reported that zinc had critical roles in the defense system of cells against ROS, and thus represents an excellent protective agent against the oxidation of several vital cell components such as membrane lipids and proteins, chlorophyll, SH-containing enzymes and DNA [12]. Sharma and his colleagues (1995) also reported that Zn deficiency decrease CO<sub>2</sub> fixation, stomata conductance, chlorophyll level and photosynthetic efficiency in plants [13].

\*Corresponding author: H.R. Karimi, Department of Horticultural sciences, College of Agriculture, Vali-e-Asr University of Rafsanjan, Rafsanjan, Kerman, Iran E-mail: h\_karimi1019@yahoo.com

Zinc deficiency is now recognized as one of the most critical micronutrient deficiency in plants grown on calcareous, saline and sodic soils with high pH values. The foliar Zn application could alleviate this problem. Esmailizadeh and his colleagues (2009) and Brown and his colleagues (1993) reported that foliar zinc application improved fruit weight and splitting percentage in pistachio [14, 15].

There are limited data in the literature about SA effects on fruit trees. The objective of this study was to determine the effect of SA and zinc on quantity and quality characteristics of pistachio cv. Owhadi.

## MATERIALS AND METHODS

### *Experimental design*

This research was conducted in 2009 in 35- year-old commercial orchard of *Pistacia vera* L. cv. Owhadi grafted on 'Badami-e-zarand' rootstock.

The experiment was conducted as factorial in the framework of completely randomized design (CRD) with three replications, included one plant.

### *Treatments*

Treatments included distilled water as control ( $S_1Zn_1$ ), 0.1% Zn without SA ( $S_1Zn_2$ ), 0.2% Zn without SA ( $S_1Zn_3$ ), 250 mg L<sup>-1</sup> SA without Zn ( $S_2Zn_1$ ), 250 mg L<sup>-1</sup> SA plus 0.1% Zn ( $S_2Zn_2$ ), 250 mg L<sup>-1</sup> SA plus 0.2% Zn ( $S_2Zn_3$ ), 500 mg L<sup>-1</sup> SA without Zn ( $S_3Zn_1$ ), 500 mg L<sup>-1</sup> SA plus 0.1% Zn ( $S_3Zn_2$ ), 200 mg L<sup>-1</sup> SA plus 0.2% Zn ( $S_3Zn_3$ ). The trees sprayed with the solutions until dripping with an atomizer.

### *Nut characteristics*

Inflorescence bud retention was recorded during plant growth. At harvest time, the nut yield of each tree was recorded and 100 fruits from each tree were taken randomly and quantity and quality characteristics of fruit and nut (nut fresh weight, kernel dry weight, nut splitting percentage and

blank nut percentage and once) were measured. For kernel dry weight determination, plant materials were dried in an oven at 75 ° C for 48 hours. Splitting nuts were determined according to Zeng and Brown (1998) [16].

### *Chlorophyll fluorescence*

A chlorophyll fluorometer (Hansatech LTD Pocket PEA) was used to measure chlorophyll fluorescence of leaves of plants at 15 days after foliar application of the various treatments. For each plant measurements were taken at four locations on each leaf from recurrent season shoots on fully expanded leaves and then averaged.

### *Data analysis*

Data analysis was performed using MSTATC statistical software and the means were compared using Duncan's Multiple Range Test (DMRT,  $P=0.05$ ).

## RESULTS

### *Inflorescence bud retention*

Inflorescence bud retention was affected by salicylic acid and Zn so that the application of salicylic acid increased inflorescence bud retention up to 250 mg L<sup>-1</sup>. The application of Zn<sub>2</sub>SO<sub>4</sub> increased inflorescence bud retention so that the highest inflorescence bud retention was observed with 0.2% Zn<sub>2</sub>SO<sub>4</sub>. The interaction of salicylic acid and Zn<sub>2</sub>SO<sub>4</sub> indicated that the highest inflorescence bud retention was observed with S<sub>2</sub>Zn<sub>3</sub>, however there was no significant difference compared to control (Table 1).

### *Nut splitting percentage*

Nut splitting percentage was affected by salicylic acid and Zn, so that the application of 500 mg L<sup>-1</sup> salicylic acid decreased splitting percentage compared to control.

The application of Zn increased splitting

percentage and the highest splitting percentage was observed with 0.2% Zn<sub>2</sub>SO<sub>4</sub>. The results of interactive effects of salicylic acid and Zn showed that simultaneous application of

salicylic acid and Zn increased splitting percentage compared to control. The highest splitting percentage was observed with S<sub>2</sub>Zn<sub>3</sub> (Table 2).

Table 1. Interactive effects of salicylic acid and Zn<sub>2</sub>SO<sub>4</sub> on inflorescence bud retention and chlorophyll fluorescence of leaves.

Zn <sub>2</sub> SO <sub>4</sub>	Salicylic acid		
	S <sub>1</sub> (control)	S <sub>2</sub> (250 mgL <sup>-1</sup> )	S <sub>3</sub> (500 mgL <sup>-1</sup> )
100 nut fresh weight (g)			
Zn <sub>1</sub> (control)	141.03 ed	147.96 b	143.06 cd
Zn <sub>2</sub> (0.1 %)	144.53 c	144.50 c	148.26 b
Zn <sub>3</sub> (0.2 %)	159.23 a	157.43 a	139.73 e
100 kernel dry weight (g)			
Zn <sub>1</sub> (control)	52.89 e	55.22 c	53.05 ed
Zn <sub>2</sub> (0.1 %)	54.40 cd	53.34 ed	56.54 b
Zn <sub>3</sub> (0.2 %)	53.02 ed	62.63 a	61.40 a
Splitting (%)			
Zn <sub>1</sub> (control)	61.00 e	74.00 c	65.00 d
Zn <sub>2</sub> (0.1 %)	85.00 ab	62.66 e	66.00 d
Zn <sub>3</sub> (0.2 %)	84.00 b	86.00 a	65.00 d
Blank (%)			
Zn <sub>1</sub> (control)	2.00 a	1.00 c	1.33 b
Zn <sub>2</sub> (0.1 %)	1.33 b	1.00 c	1.00 c
Zn <sub>3</sub> (0.2 %)	1.00 c	1.00 c	1.00 c

Different letters in each column and row show significant differences at 5% of Probability (Duncan's test).

Table 2. Interactive effects of salicylic acid and Zn<sub>2</sub>SO<sub>4</sub> on nut fresh weight, kernel weight, splitting and blank.

Zn <sub>2</sub> SO <sub>4</sub>	Salicylic acid		
	S <sub>1</sub> (control)	S <sub>2</sub> (250 mgL <sup>-1</sup> )	S <sub>3</sub> (500 mgL <sup>-1</sup> )
Inflorescence bud retention (%)			
Zn <sub>1</sub> (control)	5.66 ab	6.33 ab	5.66 ab
Zn <sub>2</sub> (0.1 %)	6.33 ab	6.66 ab	4.66 b
Zn <sub>3</sub> (0.2 %)	7.66 ab	8.32 a	5.66 ab
F <sub>v</sub>			
Zn <sub>1</sub> (control)	43.56 cb	93.62 ab	93.44 c
Zn <sub>2</sub> (0.1 %)	93.63 ab	43.71 a	43.53 cb
Zn <sub>3</sub> (0.2 %)	60.52 cb	60.52 cb	77.63 ab
F <sub>m</sub>			
Zn <sub>1</sub> (control)	72.23 cb	78.43 ab	61.90 c
Zn <sub>2</sub> (0.1 %)	76.37 ab	87.37 a	66.70 cb
Zn <sub>3</sub> (0.2 %)	68.90 cb	69.77 cb	80.70 ab

Different letters in each column and row show significant differences at 5% of Probability (Duncan's test).

#### *Blank nut percentage*

The results showed that the application of salicylic acid and Zn decreased blank percentage of nut. The lowest blank percentage was obtained with 0.2% Zn<sub>2</sub>SO<sub>4</sub> and 250 mgL<sup>-1</sup> salicylic acid.

The interaction of salicylic acid and Zn identified that the combined application of salicylic acid and Zn decreased blank nut in comparison with control. The highest blank percentage was obtained with S<sub>1</sub>Zn<sub>1</sub> (control), and the lowest it was observed with other treatment except S<sub>2</sub>Zn<sub>1</sub> S<sub>1</sub>Zn<sub>3</sub> (Table 2).

#### *Nut fresh weight*

The mean comparisons indicated that the salicylic acid and Zn<sub>2</sub>SO<sub>4</sub> increased nut fresh weight so that the highest nut fresh weight was observed with 0.2 % Zn<sub>2</sub>SO<sub>4</sub> and 250 mgL<sup>-1</sup> salicylic acid, respectively. The nut fresh weight decreased with 500 mg L<sup>-1</sup> salicylic acid compared to control.

The interactive effect of salicylic acid and Zn<sub>2</sub>SO<sub>4</sub> showed that the highest yield was gained with S<sub>1</sub>Zn<sub>3</sub>, however no significant difference was found in comparison with S<sub>2</sub>Zn<sub>3</sub> (Table 2).

#### *Kernel dry weight*

Salicylic acid decreased kernel dry weight, and the lowest kernel dry weight was obtained with 500 mg L<sup>-1</sup> salicylic acid, however there was no significant different in comparison with 250 mg L<sup>-1</sup> Zn<sub>2</sub>SO<sub>4</sub> increased it and the highest kernel dry weight was obtained with 0.2% Zn<sub>2</sub>SO<sub>4</sub>. The interactive effect of salicylic acid and Zn<sub>2</sub>SO<sub>4</sub> showed that the highest yield was gained with S<sub>2</sub>Zn<sub>3</sub>, however no significant difference was found in comparison with S<sub>1</sub>Zn<sub>3</sub>(Table 2).

#### *Leaf chlorophyll florescence*

The results showed that F<sub>m</sub> and F<sub>v</sub> increased with 250 mg L<sup>-1</sup> salicylic acid and then decreased with 500 mg L<sup>-1</sup>. The Zn<sub>2</sub>SO<sub>4</sub> increased F<sub>v</sub> while had no effect on F<sub>m</sub>. The maximum F<sub>v</sub> was observed with 0.1% Zn<sub>2</sub>SO<sub>4</sub>. The interaction of salicylic acid and

Zn on Leaf chlorophyll florescence showed that the highest F<sub>m</sub> and F<sub>v</sub> were observed with S<sub>2</sub>Zn<sub>2</sub> (Table 1).

### **DISCUSSION**

Pistachio is one of the most important horticultural crops in Iran. The majority of the pistachio producing regions is located in arid and semi-arid area with stress conditions. Therefore, suitable treatments are necessary for increasing yield efficiency and quality of this important nut crops. Our findings showed that zinc increased splitting percentage, nut fresh weight and kernel dry weight. It may be associated with the role of zinc in assimilation of carbohydrates to growing fruit, as reported by Esmacilizadeh and his colleagues (2009) and Brown and his colleagues (1993) [14,15]. They showed that the application of Zn increased fruit weight and splitting percentage in pistachio. The present study indicated that salicylic acid decreased splitting percentage. The lignin compound hydrolyzes in suture layer during maturation phase and salicylic acid decreased splitting by inhibition of the hydrolysis of lignin compound by affecting polygalacturonase enzyme. Our results indicated that salicylic acid and Zn decreased blank nut. Carbohydrate pool is the main factor affecting blankness in pistachio. The high yield depletes the carbohydrates and on the other hand, Zn affects photosynthesis and assimilated carbohydrates to growing fruits. Furthermore, Zn and salicylic acid can alleviate environmental stress, which resulted to increasing photosynthetic efficiency and followed carbohydrate content [14]. The interaction of salicylic acid and Zn indicated that inflorescence bud abscission in pistachio decreased with S<sub>2</sub>Zn<sub>3</sub> and it may be related to alleviating environmental stresses and increasing carbohydrate content resulted to photosynthesis. The effects of SA and Zn on decreasing of abiotic stress and increasing photosynthesis on crop plants reported by numerous works [1, 7, 8, 9, 10, 11, 12,

13]. Roa and his colleagues (1997) reported that SA treatment increased Cu/Zn-SOD activity and also enhanced H<sub>2</sub>O<sub>2</sub> content, which was independent of changes in catalase activity [17]. In corn plants pretreatment with SA increased antioxidant enzymes activity as well as chilling tolerance [18]. Our findings showed that the higher concentration of salicylic acid (500 mg L<sup>-1</sup>) and Zinc (0.2%) combination (S<sub>3</sub>Zn<sub>3</sub>) was decreased splitting and blank nut and this may be related to the effect of salicylic acid on hormonal imbalance in plant.

In conclusion, according to this study, salicylic acid and Zn were effective on nut quality of pistachio and alleviate of abiotic stress.

#### REFERENCES

1. Eraslan, F., Inal, A., Gunes, A. and Alpaslan, M., 2007. Impact of exogenous salicylic acid on the growth, antioxidant activity and physiology of carrot plants subjected to combined salinity and boron toxicity. *Sci Hort.*, 113: 120-128.
2. sfendiyaroglu, M. and Ozeker, E., 2008. Rooting of *Olea europaea* 'Dormat' cuttings by IBA and salicylic acid treatments. *Pakistan J Bot.*, 40: 1135-1141.
3. Zhang, L., Gao, Y., Zhang, Y., Liu, J. and Yu, J., 2010. Change in bioactive compound and antioxidant activities in pomegranate leaves. *Sci Hort.*, 123: 543-546.
4. Harper, J.P. and Balke, N.E., 1981. Characterization of inhibition of K<sup>+</sup> absorption in oat roots by salicylic acid. *Plant Physiol.*, 68: 1349-1353.
5. Khan, W., Prithiviraj, B. and Smith, D.L., 2003. Photosynthesis responses of corn and soybean to foliar application of salicylates. *J Plant Physiol.*, 160: 485-492.
6. He, Y.L., Liu, Q., Chen, C. and Bian, A.H., 2002. Thermotolerance related to antioxidation induced by salicylic acid and heat acclimation in tall fescue seedlings. *J Plant Physiol Mol.*, 28: 89-95.
7. Amin, A.A., El-Sh, M., Rashad, H.M. and EL-Abagy, H., 2007. Physiological effect of indole-3- butric acid and salicylic acid on growth, yield and chemical constituents of onion plants. *J Appli Sci Res.*, 3: 1554-1563.
8. Abdel-Wahed, M.S.A., Amin, A.A. and El-Rashad. S.M., 2006. Physiological effect of some bioregulators on vegetative growth, yield and chemical constituents of yellow maize plants. *Word J Agri Sci.*, 20: 149-153.
9. Wajahatullan, K., Balakrishnan, P. and Donald, S., 2003. Photosynthetic responses of corn and soybean to foliar application of salicylic acid. *J Plant Physiol.*, 160: 485- 492.
10. Metwally, A., Finkemeier, I., George, M. and Dietz, K., 2003. Salicylic acid alleviates the cadmium toxicity in barley seedlings. *Plant Physiol.*, 132: 272- 281.
11. Tissa, S., Darren-Touchell, E.B. and Kingsley, D., 2000. Acetyl salicylic acid (Aspirin) and salicylic acid induce multiple stress tolerance in bean and tomato plants. *Plant Growth Regul.*, 30: 157-161.
12. Cakmak, I., 2000. Possible roles of zinc in protecting plant cells from damage by reactive oxygen species. *New Physiol.*, 146: 185-205.
13. Sharma, P.N., Kumar, P. and Tewari, P.K., (1995). Early signs of oxidative stress in wheat plants subjected to zinc deficiency. *J Plant Nut.*, 27: 451-461.
14. Esmaeilzadeh, M., Talaei, A., Lessani, H., Javanshah, A. and Hokmabadi, H., 2009. The investigation the effect of shoot girdling, fruit thinning and Zn-sulphate and sucrose spray on inflorescence bud retention and qualities characteristics of pistachio cv. Owhadi. Abstract book 5<sup>th</sup> International Symposium On Pistachio and Almond., 170 p.
15. Brown, P., Zhang, Q. and Beede, B., 1993. Effect of foliar fertilization on zinc nutritional

- status of pistachio trees. In: California Pistachio Industry. Annual Report.,73-78PP.
16. Zeng, D.Q. and Brown, P.H., 1998. Effect of potassium application on soil potassium availability, leaf potassium status, nut yield and quality in mature pistachio (*Pistacia vera* L.) tree. California Pistachio Industry, Annual Report., 90-60.
17. Rao, M.V., Paliyath, G., Ormrod, D.P., Mur, D.P. and Watkins, C.B., 1997. Influence of Salicylic Acid on H<sub>2</sub>O<sub>2</sub> Production, Oxidative Stress, and H<sub>2</sub>O<sub>2</sub>-Metabolizing Enzymes (Salicylic Acid-Mediated Oxidative Damage Requires H<sub>2</sub>O<sub>2</sub>). *Plant Physiol.*, 115:137-149.
18. Janda, T., Szalai, T., Tari, I. and Paldi, E., 1999. Hydroponic treatment with salicylic acid decreases the effects of chilling injury in maize (*Zea mays* L.) plants. *Planta.*, 208: 175-180.

Archive of SID