

Induction of Systemic Resistance by *Trichoderma harzianum* Isolates in Pistachio Plants Infected with *Verticillium dahliae*

Z. Fotoohiyan¹, S. Rezaee^{*1}, Gh.H. Shahidi Bonjar², A.H. Mohammadi³, M. Moradi³

¹Department of Plant pathology, Science and Research Branch, Islamic Azad University, Tehran, Iran

²Department of Plant Pathology, Shahid Bahonar University, Kerman, Iran

³Horticultural Science Research Institute, Pistachio Research Center, Agricultural Research, Education and Extension Organization (AREEO), Rafsanjan, Iran

Received: 2 June 2015

Accepted: 26 September 2015

Abstract

Twenty isolates of *Trichoderma harzianum* were isolated from the rhizosphere of healthy pistachio plants from different localities of Kerman Province, Iran. Five isolates with high antagonistic activity in *in vitro* assays against *Verticillium dahliae* (the causal agent of pistachio wilt), were investigated for their effect on the defense enzymes, peroxidase (PO), phenyl alanine-ammonia lyase (PAL) as well as the total phenol and protein contents in pistachio seedlings exposed to *V. dahliae* under greenhouse conditions for one month after inoculation. The results indicated that all of five isolates had the ability to induce defense enzymes in treated pistachio seedlings; the Tr8 isolate had the maximum PAL activity and a corresponding increase in the total phenol content. The maximum PO activity and increase in total of protein content were seen with the Tr5 and Tr19 isolates, respectively. The increase in the activity of these enzymes when pistachio seedlings treated with antagonist alone or in combination with pathogen was greater than for plants inoculated with pathogen alone. In addition, Tr8 induced a significantly higher level of resistance in pistachio seedlings; therefore it showed the highest inhibition about 45.4% of verticillium wilt disease. This study suggests that the increased induction of defense related enzymes results in increased total phenol and protein contents due to enhanced resistance to invasion of pistachio seedlings by verticillium wilt. Outcomes of the study will be useful in formulating *T.harzianum* isolates for control of verticillium wilt in pistachio plants.

Keywords: Induced resistance, Peroxidase, Phenyl alanine-ammonia lyase, *Trichoderma harzianum*, *Verticillium dahliae*.

Introduction

Verticillium dahliae, is one of the important soilborne plant pathogens. It causes vascular wilts in more than 300 plant species including pistachio (Agrios, 2005; Williamson *et al.*, 2007). In some countries, including Iran, verticillium wilt is a serious pistachio problem (Aminae and Ershad, 1999). Because of the lack of

specificity of the host and the extreme variability of *V. dahliae* pathogenicity, control of *V. dahliae* is difficult (Pegg, 2002). The use of chemical compounds, resistant rootstocks and soil disinfestation methods are particularly important elements in current management strategies. However, the effectiveness of these management prac

*Corresponding author: E-mail: srezaee@srbiau.ac.ir

tices is lowered because *Verticillium's* mode of protection in soil as microsclerotia and the occurrence of new physiological races and chemical control is expensive and may be subject to future governmental restrictions due to environmental and health concerns (Rowe and Powelson, 2002). Recently, there has been a worldwide tendency to use eco-friendly methods in plant protection management that complement current strategies (Hajieghrari *et al.*, 2008; Mbarga *et al.*, 2012). Hence the interest in applying biological controls, for example, by using beneficial microorganisms that occur naturally in the soil and are antagonists of the pathogen (Karkachi *et al.*, 2010, Abano and Sam-Amoah, 2012).

Trichoderma spp. are among the most important biocontrol agents used for management of different diseases (Papavizas, 1985; Harman, 2004). They are free living fungi that are common in soil and root ecosystems and currently are being successfully used and commercialized to combat a broad range of soil phytopathogenic fungi (Spiegel and Chet, 1998; Yedidia and Chet, 2001; Jabnoun-Khiareddine *et al.*, 2009; Kakvan *et al.*, 2013). *Trichoderma* spp. are well known to antagonise other fungi by a variety of active and passive mechanisms. One of the mechanisms of biocontrol is the defense response that occurs during early stages of root colonization by *Trichoderma* (Howell *et al.*, 2000; Howell, 2003). As a result of interaction of *Trichoderma* with the plant, various enzymes, avirulence-like gene products and low molecular weight compounds are released from fungal or plant cell walls by the activity of *Trichoderma* enzymes (Djonovic *et al.*, 2006; Woo *et al.*, 2006; Woo and Lorito, 2007). These compounds elicit a further reaction in the plant, by activating the mycoparasitic gene expression cascade thus enhancing the biocontrol ability of *Trichoderma* (Rasmussen, 1991; Ramanathan *et al.*, 2000; Mandal, 2010). In addition, the defense reaction is enhanced due to the accumulation of PR-proteins, phytoalexins, chalcone synthase, phenylalanine-ammonia lyase (PAL), peroxidase (PO), polyphenol oxidase (PPO), phenolics, lipoxygenase, superoxide

dismutase and β -1,3-glucanase in plants (Shivakumar *et al.*, 2002; Babitha *et al.*, 2004; Girish *et al.*, 2005). PAL, one of the most extensively studied enzymes in plants because it is the first enzyme in the phenyl propanoid pathway and, catalyses the conversion of L-phenylalanine to trans-cinnamic acid which in turn enters different biosynthetic pathways leading to lignin synthesis. Thus, changes in PAL activity are the key events in controlling the synthesis of phenyl propanoids and this defense mechanism is used for protection against pathogen invasion. Induction of PAL as a response to pathogen infection is well documented in various host pathogen interactions (Geetha *et al.*, 2005, Kavitha *et al.*, 2012). Also, peroxidase is a component of an early response in plants to pathogen infection and plays the most important role in cell wall lignifications, substrate oxidation, photosynthesis, respiration and growth regulation and the plants biochemical defense against pathogens (Srivastava, 1987; Bruce and West, 1989). The products of the enzyme in the presence of hydrogen donor and hydrogen peroxide have antimicrobial activity (VanLoon and Callow, 1983). PO is one of the key enzymes involved in phenyl propanoid pathway and it is associated with disease resistance in plants (Hammerschmidt *et al.*, 1982). Several studies reported that *Trichoderma* spp. could induce resistance in different plant species against a variety of fungal pathogens (De Meyer *et al.*, 1998; Han *et al.*, 2000; Yedidia *et al.*, 2003; Shores *et al.*, 2005; Moreno *et al.*, 2009). The induction of plant defense responses and an increased resistance to pathogens by *Trichoderma* spp., also was observed when pre-treated with biocontrol agents in the field (Yedidia *et al.*, 1999; Hanson and Howell, 2004; Shores *et al.*, 2010). Induced resistance may provide an alternative approach to plant protection especially for problems not satisfactorily controlled by various fungicides (Schoenbeck, 1996).

Recent reports suggest that *Trichoderma* isolates can stimulate production of biochemical compounds of a phenolic nature associated with host defense. However,

more knowledge about these biochemical responses is needed to improve efficient formulations and potential biocontrol agents with suitable antagonistic characteristics must be screened carefully for other traits relevant to their use in a given application. There is little information on the use of *T. harzianum* as biocontrol agent against pistachio wilt caused by *V. dahliae*. Therefore in the present study we have screened local isolates of *T. harzianum* isolated from rhizosphere soil samples of healthy pistachio plants in different locations of Kerman Province for their ability to induce protection against verticillium wilt in pistachio plants by production of biochemical compounds and their potential as biocontrol agents.

Materials and Methods

Isolation of microorganisms

During 2012 – 2013, *Verticillium dahliae* isolates were obtained on selective media (Christen, 1981) from pistachio shoots with wilt symptoms. In 2013 – 2014, *Trichoderma harzianum* isolates were obtained from the rhizosphere of plants in healthy pistachio orchards in different areas of Kerman Province, on DAVET selective medium (Davet, 1979) using the technique of Rifai (1969). After proper growth, isolates were purified and identified by standard keys according to their morphology and microscopic characteristics (Goud *et al.*, 2003; Rifai, 1969; Bissett, 1991; Samuels *et al.*, 2015). From 20 isolates of *T. harzianum*, five isolates that exhibited high antagonistic activity in *in vitro* assays against *Verticillium dahliae* (in dual culture tests (Morton and Strube, 1955) and production of volatile and non-volatile metabolites (Dennis and Webster, 1971)) were investigated for their ability to induce defense enzymes. The pathogenicity of *V. dahliae* isolates was tested by the root-dipping method on pistachio seedlings (Badami zarand cultivar) at the 3rd-4th true leaf stage (Singleton *et al.*, 1992). The collected isolates were preserved on potato dextrose agar (PDA) and stored at 4°C.

Greenhouse evaluations

Preparation of plants

Pistachio seeds (Badami zarand cultivar) were washed thoroughly with sterile distilled water and then the surface was sterilized by placing the seeds in 1% sodium hypochlorite for 1 minute, after which they were rinsed three times in sterilized distilled water and placed in sterilized perlite for germination. After 3-4 days and appearance of the plumule and radicle, three germinated seeds were transferred to pots and in a mixture of soil, sand, and perlite (1/1/1, v/v/v), that had been autoclaved at 121°C for 30 minutes two successive days, and were allowed to grow in a greenhouse at 25°C for two months (Mohammadi and Banihashemi, 2002).

Preparation of inoculum of pathogen

Microsclerotia of *V. dahliae* were produced on a liquid medium as described by Hall and Ly, 1972. The cultures were inoculated in darkness at 25°C, incubated for 3 weeks under continuous shaking at 120 rpm in sterile conditions and checked regularly for the formation of microsclerotia. The inocula were then separated from the media by vacuum filtration, rinsed with sterile distilled water, dried aseptically in the shade for 72 h, weighed and passed through a 200 mesh screen to obtain smaller sizes of microsclerotia, from which 0.5 g of prepared microsclerotia used to infect 1 kg of soil.

Preparation of inoculum of *Trichoderma*

Erlenmeyer flasks containing 100 g of wheat seed and 100 mL of sterilized water were autoclaved at 121°C for one hour on three successive days. After cooling, about 5-7 small plugs of seven day old culture of *T. harzianum* isolates were dropped into each Erlenmeyer under sterilized conditions. The flasks were kept at 27°C for 4 weeks. Colonized wheat grains were then transferred into paper pockets, and were dried and ground to a powder. Ten g of prepared powder was used to infect 1 kg of soil (Frommel *et al.*, 1991).

Greenhouse biocontrol tests

Five isolates viz. Tr8, Tr19, Tr4, Tr5 and Tr18, of *T. harzianum*, all having high antagonistic activities in *in vitro* assays against *V. dahliae*, were evaluated for biocontrol experiments in the greenhouse. The seedlings with 4-5 true leaves were inoculated with pathogen and antagonist in four treatments: 1) Control (neither pathogen nor *T. harzianum*); 2) *T. harzianum*; 3) Pathogen + *T. harzianum* and 4) Pathogen. In treatments containing the antagonist, soils were inoculated with *T. harzianum* isolates seven days before infection with microslerotia of *V. dahliae*. All pots were watered as needed. Pots were kept under greenhouse conditions at $25 \pm 2^\circ\text{C}$ for one month. Pot culture experiments were conducted in the greenhouse using a completely randomized design with 4 replicates. Wilt symptoms were recorded at 10 day intervals for one month after inoculation. Disease severity was evaluated from 0 – 5 using the following scale (Huang *et al.*, 2006) 0, Healthy plants, 1, <25% of the plants wilted with scarcely any browning of the crown; 2, 25% of plants wilted and showed slight browning; 3, 50% of the plants wilted and showed progressive browning; 4, $\geq 75\%$ of plants wilted and showed complete browning; 5, dead plants.

Trichoderma as inductor of plant defence responses

Preparation of sample

Freshly leaf samples were collected at 5, 10, 20 and 30 days after inoculation with pathogen and antagonist to assay the changes in activities of defense related enzymes viz., peroxidase, phenylalanine-ammonia lyase and total of phenol and protein contents. Samples (0.5 gram) were homogenized with 1 mL of 0.1 M sodium phosphate buffer (pH 7.0) at 4°C . The homogenate was centrifuged for 20 min at 10,000 rpm. The supernatant was used for estimating plant defense enzymes activity. Crude enzyme extract in 0.1 M sodium phosphate buffer (pH 7.0) was used for the estimation of peroxidase and phenylalanine-ammonia lyase activity. The enzyme extracts were stored in deep freezer (-70°C) and utilized for later biochemical analysis.

Estimation of peroxidase (PO)

Peroxidase activity was assayed by measuring the oxidation of guaiacol in the presence of hydrogen peroxide as described by Hammer-Schmidt *et al.* (1982). A 1.5-mL aliquot of 0.05 M pyrogallol and 0.1 mL of enzyme extract were added to a cuvette. To initiate the reaction 0.5 mL of 1% H_2O_2 was added. Crude enzyme preparations were diluted to give changes in absorbance at 470 nm with differences of 0.1 to 0.2 absorbance units/min. The change in absorbance was recorded at 470 nm at 30-sec intervals for three min from zero second of incubation at room temperature. The peroxidase enzyme activity was determined for all the treated as well as control plants. The specific activity of PO was expressed as Unit/mg protein.

Estimation of Phenylalanine Ammonia Lyase (PAL)

PAL activity was determined as the rate of conversion of L-phenylalanine to *trans*-cinnamic acid at 290 nm (Dickerson *et al.*, 1984). The reaction mixture contained 1 mL enzyme extract, 0.5 mL substrate (50 mM L-phenylalanine) and 0.4 mL 25 mM Tris-HCl buffer (pH 8.8). After incubation for 1 h at 30°C , the reaction was stopped by the addition of 0.5 mL of 2 N HCl and the absorbance was read at 290 nm against a blank consisting of the same volume of reaction mixture without L-phenylalanine. The specific activity of PAL was expressed as Unit/mg protein.

Estimation of the total phenols

One gram of plant sample was homogenized in 10 mL of methanol/water, 8/2 (v/v) and agitated for 15 min at 70°C (Zieslin and Ben-Zaken, 1993). One mL of the methanolic extract was added to 5 mL of distilled water and 250 mL of Folin-Ciocalteu reagent (1 N) and the solution was kept at 25°C for 3 min. The absorbance of the developed blue color was measured at 725 nm using a spectrophotometer. Catechol was used as the standard. The total phenol content was expressed in mg/g of fresh tissue.

Estimation of the total proteins

Total protein concentration in the filtrates was assayed by the Bradford method (1976) using Coomassie blue reagent (Coomassie Protein Assay Reagent, Piere) and bovine serum albumin (BSA) as the standard protein. The specific activity of the enzymes in the total filtrate was calculated using protein concentrations determined by this method.

Statistical analysis

Data were analysed on SAS system version 9.1 (SAS Institute Inc., 1996). Mean separation was tested using Duncan's multiple range test at $p=0.05$. The test for induction of defense related enzymes and the total phenolic and protein content of pistachio seedlings by *T. harzi*

anumisolates against *V. dahliae* was established under a factorial in completely randomized design with a control and four replications for each test pathogen.

Results

Isolation of microorganisms

One isolate of *Verticillium dahliae* with high pathogenicity was isolated and used for further biocontrol investigations. Twenty isolates of *T. harzianum* collected from pistachio orchards in different areas of Kerman Province (Fig. 1), were selected and designated as Tr1, Tr2, Tr3, ... Tr20. These 20 isolates showed the highest *in vitro* activity.



Fig. 1. Sites in Kerman Province where samples were collected and isolates of *Trichoderma harzianum* were obtained are shown as white diamonds.

Greenhouse evaluations

The results of the greenhouse experiments revealed that all five isolates of *T. harzianum* had the ability to reduce wilt disease in treated pistachio seedlings (Fig. 2). The maximum wilt disease reduction observed in pots treated with Tr8 (45.4%) and Tr5 (32.9%) isolates respectively. Statistical analysis of the greenhouse

experiments revealed that wilt disease reduction of the plants with the antagonist in combination with pathogen was enhanced in comparison with treatment in which the plants were inoculated with pathogen alone. The Tr8 isolate showed the highest degree of inhibition about 45.4% of verticillium wilt disease under greenhouse conditions after one month. (Fig. 3)




Fig.2. Effect of the treatments of *Trichoderma harzianum* isolates and *Verticillium dahliae* on wilt disease reduction in pistachio seedlings under greenhouse conditions one month after inoculation. V= *V. dahliae* and Tr= *T.harzianum*




Fig.3. Effect of the treatments of Tr8 isolate and *Verticillium dahliae* in pistachio seedlings under greenhouse conditions one month after inoculation. C= Control, V= *V. dahliae* and Tr= *T.harzianum*

Greenhouse assay for biological control

The results indicated that all five isolates had the ability to induce defensive enzymes in treated pistachio seedlings, and that Tr8 isolate had maximum PAL activity and led to the maximum increase in total phenol content. The maximum PO activity and increase in total protein content was seen with the Tr5 and Tr19 isolates, respectively. The increase in the activity of these enzymes was greater for pistachio seedlings treated with antagonist alone or in combination with pathogen than for plants inoculated with pathogen alone. In addition, Tr8 induced a significantly higher level of resistance in the pistachio seedlings, and therefore showed the highest

level of inhibition about 82.5% of *Verticillium* wilt disease.

Estimation of peroxidase (PO)

The specific activity of peroxidase was found to be increased in plants treated with all five *T. harzianum* isolates and *V. dahliae*. There was significant difference in the activity of peroxidase between *T. harzianum* isolates (Fig. 4). The maximum PO activity was detected in Tr5 (4.7 U/mg protein) whereas minimum activity was from Tr19 (3.7 U/mg protein). Also, the results indicated that pistachio seedlings treated with antagonist in combina

tion with pathogen showed a significant increase in PO activity and greater than plants inoculated with antagonist or pathogen alone. At 20th day after inoculation,

enhanced PO activity was observed in almost all treatments and then slowly decreased (Fig. 4).




Fig.4. Effect of *Trichoderma harzianum* isolates and *Verticillium dahliae* on the PO activity in pistachio seedlings under greenhouse conditions one month after inoculation. C= Control, V= *V. dahliae* and Tr= *T.harzianum*, DAI=Days after inoculation

Estimation of Phenylalanine Ammonia Lyase (PAL)

The results indicated that the specific activity of phenylalanine-ammonia lyase of the strains varied from 2.1 to 2.8 (U/mgprotein) (Fig. 5). The highest specific activity was recorded for Tr8 (2.8 U/mgprotein) whereas Tr19 produced the lowest specific activity of PAL (2.1 U/mgprotein). Also, the specific activity of PAL was found to be higher in plants pretreated with *T.*

harzianum and pathogen than in plants inoculated with antagonist or pathogen alone. The control seedlings without pathogen infection displayed the lowest PAL activity. The PAL specific activity increased significantly after the challenge inoculation and reached the highest level at the 10th day after inoculation after which it slowly decreased (Fig. 5).




Fig.5. Effect of *Trichoderma harzianum* isolates and *Verticillium dahliae* on the PAL activity in pistachio seedlings under greenhouse conditions one month after inoculation. C= Control, V= *V. dahliae* and Tr= *T.harzianum*, DAI=Days after inoculation

Estimation of the total phenols

From Fig. 6, it is clear that the total phenol content increased with treatment by the all five isolates of *T. harzianum* and *V. dahliae*. All plants pretreated with *T. harzianum* isolates and *V. dahliae* showed more accumulation of total phenol than the control and for plants

pretreated with *Trichoderma* alone. The maximum total phenol content (0.68 mg/g) shown for the Tr8 isolate and the minimum (0.39 Mg/g) for Tr18. The accumulation of phenol increased from the 5th day after challenge inoculation with *V. dahliae* and reached the maximum level on the 30th day (Fig. 6).




Fig.6. Effect of *Trichoderma harzianum* isolates and *Verticillium dahliae* on the total phenol content in pistachio seedlings under greenhouse conditions for one month after inoculation. C= Control, V= *V. dahliae* and Tr= *T. harzianum*, DAI=Days after inoculation

Estimation of the total proteins

The total protein content was significantly higher in all specimens treated with *Trichoderma* isolates as compared with the untreated control (Fig.7). The pistachio seedlings treated with *T. harzianum* isolates and challenged with the pathogen showed the maximum total protein content, which was higher than the corresponding pathogen challenged control. The total

protein content reached its maximum on the 30th day after challenge inoculation (Fig. 7). However, the pistachio seedlings treated by Tr19 isolate exhibited significantly higher total protein content (0.68mg/g) than other isolates. Also, pistachio seedlings treated by the Tr5 isolate led to decrease total protein content (0.35mg/g).




Fig.7. Effect of *Trichoderma harzianum* isolates and *Verticillium dahliae* on total protein content in pistachio seedlings under greenhouse conditions for one month after inoculation. C= Control, V= *V. dahliae* and Tr= *T. harzianum*, DAI=Days after inoculation

Discussion

Application of biological control agents assists in reducing use of chemical pesticides and controlling release of their residues into the environment (Baker and Paulitz, 1996). *Trichoderma* spp. are widespread in almost any soil and rhizosphere, and have been investigated as effective biocontrol agents because of their ability to reduce of diseases caused by number of pathogenic plant fungi particularly many common soil borne pathogens (Elad, 2000; Freeman *et al.*, 2004; Dubey *et al.*, 2007; Vinale *et al.*, 2008). The beneficial action of *Trichoderma* spp. is not limited to fighting pathogens; they have also been shown to be opportunistic plant symbionts, enhancing the systemic resistance of plants (Yedidia *et al.* 1999; Shores *et al.*, 2010). Some isolates are also known for their ability to induce systemic resistance by production of biochemical compounds of phenolic nature in plants that are active against different pathogens (Kavino *et al.*, 2008; Radjacommaro *et al.*, 2010). They can induce localized or systemic resistance to diseases and their causative pathogens through the release of metabolites (Wei *et al.*, 1996; Zhou and Paulitz, 1994; Liu *et al.*, 1995). Resistance results in an increase in the concentration of metabolites and enzymes related to defense mechanisms, such as the enzymes phenylalanine-ammonia lyase (PAL), peroxidase (PO), chalcone synthase (CHS) that are involved in the biosynthesis of phytoalexins, chitinases and glucanases and high levels of phenols. Induced PO and PAL activity in plants enhances the antimicrobial properties and plant phenolics and their oxidation products provide resistance to a wide range of pathogens (Vidhyasekaran, 1988). *Trichoderma* spp. have recently led to the proposal that besides their recognized antifungal properties, such organisms could also act as elicitors of plant defense reactions, thereby promoting the expression of plant defense related metabolites (Yedidia *et al.*, 2003, Segarra *et al.*, 2007).

In this study, we evaluated the ability of five strains of *T. harzianum* as biological agents, isolated from the rhizosphere soil of healthy pistachio plants from Iran, to induce systemic resistance to verticillium wilt by way of the defense enzymes, PO and PAL and total phenol and protein contents in pistachio seedlings grown under greenhouse conditions. PO is a useful marker of plant resistance to infection and stress (Welinder, 1992). POs are used primarily for the synthesis of secondary metabolites and are known to be induced by pathogen infection (Delannoy *et al.*, 2003; Sasaki *et al.*, 2005). The increased PO activity contributes to disease resistance in infected plants (Vidhyasekaran, 1997). Therefore, the increase of PO activity by *T. harzianum* isolates in the all pistachio seedlings infected with *V. dahliae*, even in the absence of *V. dahliae* infection, can be considered as a marker of disease resistance during fungal phytopathogenesis in plants possibly through its utilization in cell wall lignification. Our results corroborate studies of Bradley *et al.* (1992), who reported that increased PO activity has been correlated with resistance in many species of plants and that these enzymes are involved in the polymerization of proteins and lignin or suberin precursors into plant cell walls, thus constructing a physical barrier that can prevent pathogen penetration of cell walls or movement through vessels. Other studies, (Nawar and Kuti, 2003; Hassan *et al.*, 2007; Van Wees *et al.*, 2008) have delineated the induction of PO in plants infected by pathogens or insects, resulting in faster and stronger resistance to them. Also, *T. harzianum* isolates increased PAL enzyme activity in pistachio seedlings. PAL induced phenyl propanoid metabolism starts with the conversion of L-phenylalanine into *trans*-cinnamic acid thus supplying precursors for flavanoid pigments, lignin and phytoalexins (Massala *et al.*, 1980; Hahlbrock and Scheel, 1989). An increase in PAL activity subse

quently might have led to increased levels of the signaling molecule salicylic acid and the phenolic compounds in the host thereby contributing to disease resistance (Klessig and Malamy, 1994; Charitha Devi *et al.*, 2012). Induction of PAL by *Pseudomonads fluorescens* against *C. gloeosporioides* in mango and noni has been reported (Vivekananthan *et al.*, 2004; Manjunath, 2009). In our study, a significant increase in the level of PAL in the treated pistachio seedlings is in agreement with earlier reports (Yedidia *et al.* 2003; Verma *et al.* 2007).

This finding is well supported by the study of Shores *et al.*, (2005), who reported high induction of PAL in cucumber plants treated with *T. asperellum*. as well as that of Kavitha *et al.*, (2008), who reported the accumulation of PAL and PO mediated by the *T. asperellum* in tomato. Similarly, systemic resistance was enhanced due to a high accumulation of defense enzymes in response to *R. solanacearum* challenge in tomato (Vanitha *et al.*, 2009). Further, the interaction between tomato and *V. dahliae* elicited enhanced activities of PO and PAL, phenylpropanoid metabolism, and synthesis of lignins (Gayoso *et al.*, 2010). Also, it was observed that resistant plants contain more phenols and proteins. A multifold increase in phenol and protein content was observed in the all the pistachio plants treated with *T. harzianum* isolates alone or with the pathogen, compared with the control plants. Some phenolics may act as signal molecules or antioxidants and thus induce resistance (Malamy *et al.*, 1990). The accumulation of phenols may be due to excess production of H₂O₂ in infected plants through increased respiration (Farkas and Kiraly, 1962) or to the activation of hexose-monophosphate shunt pathway, acetate pathway and to the release of bound phenols (Goodman *et al.*, 1967; De Ascensao and dubery, 2000; Mandal and Mitra, 2007). These observations closely resemble findings of resistance in maize roots elicited with *T. harzianum* strain T-22 (Bigirimana *et al.*, 1997). Similarly, induction of synthesis of high amounts of phenols by *T. harzi-*

anum isolates, compared with controls, suggests their role in inducing resistance against wilt in pistachio plants. The level of defense-related enzymes determines the degree of host resistance. Increase in activity and accumulation of these enzymes also depends on the plant genotype, physiological conditions and the pathogen.

In this study, it has been concluded that the pistachio plants treated with native bioagents of *T. harzianum* followed by challenge inoculation of *V. dahliae* enhances induction of defense related enzymes and that these were very effective in the control of verticillium wilt of pistachio plants. In conclusion, plants treated with *Trichoderma* in the root zone can produce higher levels of PO, PAL, phenols, and pathogenesis related proteins. In this regard, other researchers have also shown in the greenhouse similar results; e.g. *Trichoderma* isolates have been shown to be successful in controlling soil borne diseases (Basin *et al.*, 1999). *Trichoderma* spp. decreased wilt incidence in chickpea plants (Dubey *et al.*, 2007). The selection of biocontrol agents and the understanding the mechanisms involved in the antagonistic effect of *Trichoderma* spp. on plant pathogens are important in designing effective and safe biocontrol strategies. The different isolates of *Trichoderma* have different combative abilities for pathogen; their indirect effects may also vary.

From the current study, it can be inferred that there is maximum disease reduction in the application of indigenous *Trichoderma* isolate Tr8 against pistachio wilt as compared to the other tested isolates. This may be to the case for a number of reasons, including pathogenicity of isolates, climatic adaptability, influence of the pathogen origin and even the influence of local pistachio cultivars used in this region (Harman, 2006; Sharon *et al.*, 2007). Therefore, the Tr8 isolate native to Kerman Province could be an excellent candidate for providing long term biocontrol agent against *V. dahliae* in pistachio plants with the aim of reducing the use of chemical pesticides in this region. The mechanisms involved in induced

plant resistance are still poorly understood and to exploit such applications of *Trichoderma*, more research is needed.

Conclusions

The present investigation was aimed at understanding the induced systemic resistance of pistachio plants treated with native, highly potent *T. harzianum* isolates possessing rhizosphere competence and antifungal properties against *V. dahliae* at different times after challenge inoculation. Pre and post inoculation studies demonstrated that the pistachio plants responded well to select *T. harzianum* isolates, which induced some systemic resistance to *V. dahliae* infection and caused significant changes in the host plant in terms of total protein, phenol content and levels of PO and PAL. Due to the increase in plant defense enzymatic activity, and the decrease in disease incidence, the *Trichoderma* inoculum can also be applied in the field in order to enhance the yield compared to control plants. Hence, this study shows clear evidence that the native isolates of *Trichoderma* inoculum can not only enrich soil fertility and crop yield but also induce disease resistance and help to increase production. Different enzymes, PAL and PO which are induced of all defense mechanisms. The mechanisms are similar for *Trichoderma* spp. against different pathogens (Bolton, 2009). Therefore, this beneficial impact of new strains of antagonist in experimental systems with different plants and pathogens aimed to determine the ability of *Trichoderma* strain to induce defense response and resistance in a variety of biotic and abiotic conditions.

Acknowledgements

The authors thank Mr. Aminae, Mr. Abosaidi and, Mr. Tahari and Ms. Lori for their generous technical assistance and Mr. Khazaei for his support and help.

References

- Abano EE, Sam-Amoah LK (2012) Application of antagonistic microorganisms for the control of postharvest decays in fruits and vegetables. *International Journal of Advances in Biology Research*. 2(1), 1-8.
- Agrios GN (2005) *Plant Pathology*. 5th Ed. San Diego, Elsevier Academic Press, 922 pp.
- Aminae MM, Ershad D (1999) Occurrence of Verticillium wilt on Pistachio trees in Kerman province (Iran). *Iranian Journal of Plant Pathology*. 35, 59.
- Asaki-Sekimoto Y, Taki N, Obayashi T, Aono M, Matsumoto F, Sakurai N, Suzuki H, Hirai MY, Noji M, Saito K (2005) Coordinated activation of metabolic pathways for antioxidants and defence compounds by jasmonates and their roles in stress tolerance in Arabidopsis. *Plant Journal*. 44, 653-668.
- Babitha MP, Prakash HS, Shetty HS (2004) Purification and properties of lipoxygenase induced in downy mildew resistant pearl millet seedlings due to infection with *Sclerospora graminicola*. *Plant Science*. 166, 31-39.
- Baker R, Paulitz TC (1996) Theoretical basis for microbial interactions leading to biological control of soilborne plant pathogens. In: *Principles and Practice of Managing Soilborne Plant Pathogens*. Hall, R. (ed.). The American Phytopathology Society. Pp.50-79.
- Basin H, Ozturk SB, Yegen O (1999) Efficacy of a biological fungicide (Planter Box *Trichoderma harzianum* Rifai T=22) against seedling root rot pathogens (*Rhizoctonia solani*, *Fusarium* sp.) of cotton, GAP-Environmental Symposium, Sanleurfa, Turkey. Pp. 137-144.
- Bigirimana J, De Meyer G, Poppe J, Elad Y, Hofte M (1997) Induction of systemic resistance on bean (*Phaseolus vulgaris*) by *Trichoderma*

- harzianum*. Med. Fac. Landbouww Univ Gent. 62, 1001–1007.
- Bissett J (1991) The revision of the genus *Trichoderma* III. Section *Pachybasidium*. Canadian Journal of Botany. 69, 2357-2372.
- Bolton MD (2009) Primary metabolism and plant defense fuel for the fire. Molecular Plant Microbe Interaction. 22, 487–497.
- Bradford MM (1976) A rapid and sensitive method for quantification of microgram quantities of protein utilizing the principle of protein-dyebinding. Analytical Biochemistry. 72, 248-254.
- Bradley DJ, Kjellborn P, Lamb C (1992) Elicitor and wound induced oxidative cross-linking of a plant cell wall proline-rich protein: A novel, rapid defense response. Cell. 70, 21-30.
- Bruce RJ, West CA (1989) Elicitation of lignin biosynthesis and isoperoxidase activity by pectic fragments in suspension cultures of cluster bean. Plant Physiology. 91, 889-897
- Charitha Devi M, Radha Y (2012) Induced biochemical changes in the CMV infected cucurbit plants. Annals of Biology Research. 3(2), 863-870.
- Christen AA (1981) A selective medium for isolating *Verticillium albo-atrum* from soil. Phytopathology 72, 47–49.
- Davet P (1979) Technique pour l'analyse des population et de *Trichoderma* et de *Gliocladium virens* dans le sol. Annual Review of Phytopathology. 11, 529-533.
- De Ascensao ARCF, Dubery IA (2000) Panama disease: Cell wall reinforcement in banana roots in response to elicitors from *Fusarium oxysporum* f.sp. cubense race four. Phytopathology. 90, 1173–1180.
- De Meyer G, Bigirimana J, Elad Y, Hofte M (1998) Induced systemic resistance in *Trichoderma harzianum* T39 biocontrol of *Botrytis cinerea*. European Journal of Plant Pathology. 104, 279–286.
- Delannoy E, Jalloul A, Assigbetsé K, Marmey P, Geiger JP, Lherminier JF, Daniel JF, Martinez C, Dickerson DP, Pascholati SF, Hagerman AE, Butler LG, Niholson RL (1984) Phenylalanine ammonia lyase and hydroxyl cinnamate: CoA ligase in maize mesocotyls inoculated with *Helminthosporium maydis* or *Helminthosporium carbonum*. Physiology and Plant Pathology. 25, 111-123.
- Djonovic S, Pozo MJ, Dangott LJ, Howell CR, Kenerley CM (2006) Sm1, a proteinaceous elicitor secreted by the biocontrol fungus *Trichoderma virens* induces plant defense responses and systemic resistance. Molecular Plant Microbe Interaction. 8, 838-53.
- Dubey SC, Suresh M, Singh B (2007) Evaluation of *Trichoderma* species against *Fusarium oxysporum* f.sp. ciceris for integrated management of chickpea wilt. Biology Control. 40, 118-127.
- Elad Y (2000) Biological control of foliar pathogens by means of *Trichoderma harzianum* and potential modes of action. Crop Protection. 19, 709-714.
- Freeman S, Minz D, Kolesnik I, Barbul O, Zreibil A, Maymon M, Nitzani Y, Kirshner B, Rav-David D, Bilu A, Dag A, Shafir S, Elad Y (2004) *Trichoderma* biocontrol of *Colletotrichum acutatum* and *Botrytis cinerea*, and survival in strawberry. European Journal of Plant Pathology. 110, 361-370.
- Frommel MI, Pazos GS, Nowak J (1991) Plant-growth stimulation and biocontrol of *Fusarium* wilt (*Fusarium oxysporum* f. sp. lycopersici) by co-inoculation of tomato seeds with *Serratia plymuthica* and *Pseudomonas* sp. Phytopathology. 26, 66-73.
- Gayoso C, Pomar F, Novo-Uzal E, Merino F, Martinez de Ilarduya O (2010) The Ve-mediated re-

- sistance response of the tomato to *Verticillium dahliae* involves H₂O₂, peroxidase and lignins and drives PAL gene expression. BMC Plant Biology. 10, 232.
- Geetha NP, Amruthesh KN, Sharathchandra RG, Shetty HS (2005) Resistance to downy mildew in pearl millet is associated with increased Phenylalanine ammonia lyase activity. Functional Plant Biology. 32, 267-275.
- Girish N, Umesha S (2005) Effect of plant growth promoting rhizobacteria on bacterial canker of tomato. Arch. Phytopathol. Plant Protection. 38, 235-243.
- Goodman RN, Kiraly E, Zaitlin M (1967) The biochemistry and physiology of infectious plant diseases. Princeton, NJ: D. VanNosti and Co. Farkas G.L. and Kiraly Z. 1962. Role of phenolic compounds in the physiology of plant disease resistance. Phytopathology. 44, 105-150.
- Goud JK., Termorshuizen AJ, Gams W (2003) Morphology of *Verticillium dahliae* and *V. tricorpus* on semi-selective media used for the detection of *V. dahliae* in soil. Mycology Reserch. 107, 822-830.
- Hahlbrock K, Scheel D (1989) Physiology and Molecular biology of phenylpropanoid metabolism. Annal of Review Plant Phsiology and Plant Molecular Biology. 40, 347-367.
- Hajiehgrari B, Torabi-Giglou M, Mohammadi MR, Davari MM (2008) Biological potential of some Iranian Trichoderma isolates in the control of soil borne plant pathogenic fungi. African Journal Biology Control. 7, 967-972.
- Hall R, Ly H (1972) Development and quantitative measurement of microsclerotia of *Verticillium dahliae*. Canadian Journal of Botany. 50, 2097-2102.
- Hammerschmidt R, Nuckles EM, Kuc J (1982) Association of enhanced peroxidase activity with induced systemic resistance of cucumber to *Colletotrichum lagenarium*. Physiology and Plant Pathology. 20, 73-82.
- Han DY, Coplin DL, Bauer WD, Hoitink HAJ (2000) A rapid bioassay for screening rhizosphere microorganisms for their ability to induce systemic resistance. Phytopathology. 90, 327-332.
- Hanson LE, Howell CR (2004) Elicitors of plant defense responses from biocontrol strains of *Trichoderma virens*. Phytopathology. 94, 171-176.
- Harman GE (2004) Trichoderma species opportunistic, avirulent plant symbionts. Nature Reviews Microbiology. 2, 43-56.
- Harman GE (2006) Overview of mechanisms and uses of *Trichoderma* sp. Phytopathology. 96, 190-194.
- Hassan E, Maggie M, Saieda S, Abd El-Rahman I, El-Abbasi H, Mikhail MS (2007) Changes in peroxidase activity due to resistance induced against faba bean chocolate spot disease. Egyptian Journal of Phytopathology. 35(1), 35-48.
- Howell CR (2003) Mechanisms employed by *Trichoderma* species in the biological control of plant diseases: the history and evolution of current concepts. Plant Disease. 87, 4-10.
- Howell CR, Hanson LE, Stipanovic RD, Puckhaber LS (2000) Induction of terpenoid synthesis in cotton roots and control of *Rhizoctonia solani* by seed treatments with *Trichoderma virens*. Phytopathology. 90(3), 248-252.
- Huang LC, Ryan AF, Cockayne DA, Housley GD (2006) Developmentally regulated expression of the P2X (3) receptor in the mouse cochlea. Histochem. Cell Biology. 125, 681-692.
- Jabnoun-Khiareddine H, Daami-Remadi M, Ayed F, El-Mahjoub M (2009) Biological control of tomato Verticillium wilt by using indigenous *Trichoderma* spp. African Journal of Plant Science and Biotechnology. 3(1), 26-36.
- Kakvan N, Heydari A, Zamanizadeh HR, Rezaee S, Naraghi L (2013) Development of new biofor-

- mulations using *Trichoderma* and *Talaromyces* fungal antagonists for biological control of sugar beet damping-off disease. *Crop Protection*. 53, 80-84.
- Karkachi NE, Gharbi S, Kihal M, Henni JE (2010) Biological control of *Fusarium oxysporum* f.sp. *lycopersici* isolated from Algerian tomato by *Pseudomonas fluorescens*, *Bacillus cereus*, *Serratia marcescens* and *Trichoderma harzianum*. *Research Journal of Agronomy*. 4, 31-34.
- Kavino M, Harish S, Kumar N, Saravanakumar D, Samiyappan R (2008) Induction of systemic resistance in banana (*Musa* spp.) against Banana bunchy top virus (BBTV) by combining chitin with root-colonizing *Pseudomonas fluorescens* strain CHAO. *European Journal of Plant Pathology*. 120, 353-362.
- Kavitha R, Umesh S (2008) Regulation of defense-related enzymes associated with bacterial spot resistance in tomato. *Phytoparasitica*. 36(2), 144-159.
- Klessig DF, Malamy J (1994) The salicylic acid signals in plants. *Plant Molecular Biology*. 26, 1439-1458.
- Liu L, Kloepper JW, Tuzun S (1995) Induction of systemic resistance in cucumber by plant growth-promoting rhizobacteria: duration of protection and effect of host resistance on protection and root colonization. *Phytopathology*. 85, 1064-1068.
- Malamy J, Carr JP, Klessig DF, Raskin I (1990) Salicylic acid: a likely endogenous signal in the resistance response of tobacco to viral infection. *Science*. 250, 1002-1004.
- Mandal S, Mitra A (2007) Reinforcement of cell wall in roots of *Lycopersicon esculentum* through induction of phenolic compounds and lignin by elicitors. *Physiological and Molecular Plant Pathology*. 71, 201-209.
- Mandal S (2010) Induction of phenolics, lignin and key defence enzymes in eggplant (*Solanum melongena* L.) roots in response to elicitors. *African Journal of Biotechnology*. 9(47), 8038-8047.
- Manjunath H (2009) Morphological and molecular characterization of *Alternaria alternata* and *Colletotrichum gloeosporioides* incitants of leaf blight and anthracnose diseases of noni and their management. M.Sc. (Agri.) Thesis, Tamil Nadu Agricultural University, Coimbatore.
- Massala R, Legrand M, Fritig B (1980) Effect of amino oxyacetate, a competitive inhibitor of phenylalanine ammonia lyase, on the hypersensitive resistance of tobacco to tobacco mosaic virus. *Physiology and Plant Pathology*. 16, 213-226.
- Mbarga JB, Ten Hoopen GM, Kuate J, Adiobo A, Ngonkeu MEL, Ambang Z, Akoa A, Tondje PR, Begoude BAD (2012) *Trichoderma asperellum*: A potential biocontrol agent for *Pythium myriotylum*, causal agent of cocoyam (*Xanthosomasagittifolium*) root rot disease in Cameroon. *Crop Protection*. 36, 18-22.
- Mohammadi AH, Banihashemi Z (2002) Effect of NaCl, Na²SO⁴ and their combination on different growth stages of *Verticillium dahliae*. *Proc. 15th Plant Protection. Cong., Iran, Kermanshah*.
- Moreno CA, Castillo F, González A, Bernal D, Jaimes Y, Chaparro M, González C, Rodríguez F, Restrepo S, Cotes AM (2009) Biological and molecular characterization of the response of tomato plants treated with *Trichoderma koningiopsis*. *Physiological and Molecular Plant Pathology*. 74, 111-120.
- Nawar HF, Kuti JD (2003) Weyerone acid phytoalexin synthesis and peroxidase activity as markers for resistance of broad bean to chocolate spot disease. *Phytopathology*. 151, 564-570.

- Nicole M (2003) Activity of class III peroxidases in the defense of cotton to bacterial blight. *Molecular Plant-Microbe Interaction*. 16, 1030-1038.
- Papavizas GC (1985) *Trichoderma* and *Gliocladium*: biology, ecology and potential for the biocontrol. *Annals of Review Phytopathology*. 23, 23-54.
- Pegg GF (ed.) (2002) *Verticillium Wilts*. CAB International, Wallingford, UK, 576 pp.
- Radjacommaro R, Venkatesan S, Samiyappan R (2010) Biological control of phytopathogenic fungi of vanilla through lytic action of *Trichoderma* species and *Pseudomonas fluorescens*. *Archives of Phytopathology and Plant Protection*. 43, 1-17.
- Ramanathan A, Samiyappan R, Vidhyasekaran P (2000) Induction of defence mechanisms in greengram leaves and suspension cultured cells by *Macrophomina phaseolina* and its elicitors. *Journal of Plant Disease Protection*. 107, 245-257.
- Rasmussen JB (1991) Systemic induction of salicylic acid accumulation in cucumber after inoculation with *Pseudomonas syringae* pv. *syringae*. *Plant Physiol*. 97, 1342-1347.
- Rifai MA (1969) A revision of the genus *Trichoderma*. *Mycological Papers*. 116, 1-56.
- Rowe RC, Powelson ML (2002) Potato early dying: Management challenges in a changing production environment. *Plant Disease*. 86, 1184-1193.
- Samuels GJ, Chaverri P, Farr DF, Mc Cray EB (2015) *Trichoderma* Online, Systematic Mycology and Microbiology Laboratory, ARS, USDA. <http://nt.arsgrin.gov/taxadescriptions/keys/TrichodermaIndex.cfm>.
- SAS Institute Inc (1996) SAS/STAT Software: Changes and Enhancements through Release 6.11. SAS 20 Institute Inc., Cary, NC.
- Schoenbeck F (1996) Induced resistance: mechanisms and evaluation. In: Lyr H., Russel P. and Sisler H.D. (Eds.). *modern fungicides and antifungal compounds*. Andover, UK: Intercept Ltd. Pp. 447-450.
- Segarra G, Casanova E, Bellido D, Odena MA, Oliveira E, Trillas I (2007) Proteome, salicylic acid and jasmonic acid changes in cucumber plants inoculated with *Trichoderma asperellum* strain T34. *Proteomics*. 7, 3943-3952.
- Sharon E, Chet A, Viterbo M, Bar-Eyal H, Nagan GJ, Samuels Y, Spiegel Y (2007) Parasitism of *Trichoderma* on *Meloidogyne javanica* and role of the gelatinous matrix. *European Journal of Plant Pathology*. 118, 247-258.
- Shivakumar PD, Geetha HM, Shetty HS (2002) Peroxidase activity and isozymes analysis of pearl millet seedlings and their implication in downy mildew disease resistance. *Plant Science*. 160, 1-9.
- Shoresh M, Harman GE, Mastouri F (2010) Induced systemic resistance and plant responses to fungal biocontrol agents. *Phytopathology*. 48, 21-43.
- Shoresh M, Yedidia I, Chet I (2005) Involvement of jasmonic acid/ethylene signaling pathway in the systemic resistance induced in cucumber by *Trichoderma asperellum* T203. *Phytopathology*. 95, 76-84.
- Singleton LL, Mihail JD, Rush CM (1992) *Methods for Research on Soilborne Phytopathogenic Fungi*. APS Press, USA, 265pp.
- Spiegel Y, Chet I (1998) Evaluation of *Trichoderma* spp., as biocontrol agent against soil borne fungi and plant parasitic nematodes in Israel. *Integrative Pest Management Review*. 3, 169-175.
- Srivastava SK (1987) Peroxidase and polyphenol oxidase in Brassica juncea plants infected with *Macrophomina phaseolina* (Tassi) Goid and their implications in disease resistance. *Phytopathology*. 77, 249-254.

- Van Loon LC, Callow JA (1983) Transcription and translation in the diseased plant In: Biochemical Plant Pathology, Callow JA (Eds.) John Wiley and Sons, Chichester, UK.
- Van Wees SCM, Van Der Ent S, Pieterse CMJ (2008) Plant immune responses triggered by beneficial microbes. *Current Plant Biology*. 11, 443–448.
- Vanitha SC, Niranjana SR, Mortensen CN, Umesh S (2009) Bacterial wilt of tomato in Karnataka and its management by *P. fluorescens*. *Biocontrol*. 54, 685-695.
- Verma M, Brar SK, Tyagi RD, Surampalli RY, Valéro JR (2007) Antagonistic fungi, *Trichoderma* spp.: panoply of biological control. *Biochemical Engineering*. 37, 1–20.
- Vidhyasekaran P (1988) Physiology of disease resistance in plants. Vol. 2. CRC Press, Boca Raton, Florida. 128 p.
- Vidhyasekaran P (1997) Fungal pathogenesis in plants and crops: In: Marcel Dekker, editors. *Molecular Biology and Host defense Mechanisms*. New York. Pp. 67-73.
- Vinale F, Sivasithamparan K, Ghisalberti EL, Marra R, Woo SL, Lorito M (2008) *Trichoderma*–plant–pathogen interactions. *Soil Biology and Biochemistry*. 40, 1-10.
- Vivekananthan R, Ravi M, Ramanathan A, Samiyappan R (2004) Lytic enzymes induced by *Pseudomonas fluorescens* and other biocontrol organisms mediate defense against the anthracnose pathogen in mango. *World Journal of Microbiology and Biotechnology*. 20, 235-244.
- Wei G, Kloepper JW, Tuzun S (1996) Induced systemic resistance to cucumber diseases and increased plant growth by plant growth promoting rhizobacteria under field conditions. *Phytopathology*. 86, 221-224.
- Welinder KG (1992) Superfamily of plant, fungal and bacterial peroxidases. *Current Opinion Structure Biology*. 2, 388–393.
- Williamson B, Tudzynski B, Tudzynski P, Van Kan JAL (2007) *Botrytis cinerea*: the cause of grey mould disease. *Molecular Plant Pathology*. 8, 561-580.
- Woo S, Scala F, Ruocco M, Lorito M (2006) The molecular biology of the interactions between *Trichoderma* spp., phytopathogenic fungi, and plants. *Phytopathology*. 96(2), 181-185.
- Woo SL, Lorito M (2007) Exploiting the interactions between fungal antagonists, pathogens and the plant for biocontrol. In: Vurro, M., Gressel, J. (Eds.), *Novel Biotechnologies for Biocontrol Agent Enhancement and Management*. IOS, Springer Press, Amsterdam, the Netherlands. Pp. 107–130.
- Yedidia I, Chet I (2001) Effect of *Trichoderma harzianum* on microelement concentrations and increased growth of cucumber plants. *Plant Soil*. 235, 235-242.
- Yedidia I, Benhamou N, Chet I (1999) Induction of defense response in cucumber plants (*Cucumis sativus* L.) by the biocontrol agent *Trichoderma harzianum*. *Applied and Environmental Microbiology*. 65, 1061–1070.
- Yedidia I, Shores M, Kerem Z, Benhamou N, Apulnik YK, Chet I (2003) Concomitant induction of systemic resistance to *Pseudomonas syringae* pv. lachrymans in cucumber by *Trichoderma asperellum* (T-203) and accumulation of phytoalexins. *Applied and Environmental Microbiology*. 69, 7343–7353.
- Yedidia I, Srivastava AK, Kapulink Y, Chet I (2001) Effect of *Trichoderma harzianum* on microelement concentrations and increased growth of cucumber plants. *Plant and Soil*. 235, 235–242.
- Zhou T, Paulitz TC (1994) Induced resistance in the biocontrol of *Pythium aphanidermatum* by

Pseudomonas spp. on cucumber. *Journal of Phytopathology*.142, 51–63.

Zieslin N, Ben-Zaken R (1993) Peroxidase activity and presence of phenolic substances in peduncles of rose flowers. *Plant Physiology and Biochemistry*. 31, 333-339.

Archive of SID

القای مقاومت سیستمیک توسط جدایه‌های *Trichoderma harzianum* در گیاهان پسته آلوده به *Verticillium dahliae*

ز. فتوحیان^۱، س. رضایی^{۱*}، غ.ح. شهیدی بنجار^۲، ا.ح. محمدی^۳، م. مرادی^۳

۱. گروه گیاهپزشکی، واحد علوم و تحقیقات، دانشگاه آزاد اسلامی، تهران، ایران

۲. گروه گیاهپزشکی، دانشگاه شهید باهنر، کرمان، ایران

۳. موسسه تحقیقات علوم باغبانی، پژوهشکده پسته، سازمان تحقیقات، آموزش و ترویج کشاورزی، رفسنجان، ایران

مسئول مکاتبات: Email: srezaee@srbiau.ac.ir

تاریخ پذیرش: ۴ مهر ۱۳۹۴

تاریخ دریافت: ۱۲ خرداد ۱۳۹۴

چکیده

۲۰ جدایه *Trichoderma harzianum* از ریزوسفر درختان سالم پسته در مناطق مختلف استان کرمان، ایران جدا شدند. پنج جدایه با فعالیت آنتاگونیستی بالا در شرایط آزمایشگاهی علیه قارچ *Verticillium dahliae* عامل پژمردگی پسته، تاثیرشان روی آنزیم‌های دفاعی، پراکسیداز (PO) و فنیل‌آلانین آمونیلایز (PAL)، همچنین محتوی فنل و پروتئین کل در نهال‌های پسته آلوده شده با *V. dahliae* در شرایط گلخانه به مدت یک ماه پس از تلقیح بررسی شد. نتایج نشان داد که تمامی پنج جدایه تریکودرما قادر به القای آنزیم‌های دفاعی در نهال‌های پسته تیمار شده بودند. در حالی که جدایه Tr8 دارای بیشترین فعالیت PAL و به طور متقابل افزایش محتوی فنل کل بود. بیشترین فعالیت PO و افزایش محتوی پروتئین به ترتیب در جدایه‌های Tr5 و Tr19 مشاهده شد. افزایش فعالیت این آنزیم‌ها در نهال‌های پسته تیمار شده با آنتاگونیست به تنهایی یا همراه با پاتوژن، بیشتر از نهال‌های تلقیح شده با پاتوژن به تنهایی بود. بعلاوه جدایه Tr8 به طور معنی‌داری سطح بالاتری از مقاومت را در نهال‌های پسته القاء کرد، بنابراین بالاترین میزان بازدارندگی حدود ۴۵/۴ درصد از بیماری پژمردگی ورتیسیلیومی را نشان داد. این مطالعه تایید کرد که افزایش القاء آنزیم‌های دفاعی و افزایش محتوی فنل و پروتئین کل منجر به بالا بردن مقاومت نهال‌های پسته در معرض پژمردگی ورتیسیلیومی می‌شود. نتایج این تحقیق در فرموله کردن جدایه‌های *T. harzianum* جهت کنترل پژمردگی ورتیسیلیومی در گیاهان پسته مفید خواهد بود.

واژه‌های کلیدی: *Trichoderma harzianum*، *Verticillium dahliae*، مقاومت القایی، پراکسیداز، فنیل‌آلانین آمونیلایز.