Journal of Economic Geology Vol. 10, No. 1 (2018) ISSN 2008-7306



زمین شناسی اقتصادی جلد ۱۰، شماره ۱ (سال ۱۳۹۷) صفحات ۳ ه ۴

# Petrology, Geochemistry and Tectonomagmatic Setting of Farmahin Volcanic Rocks (North of Arak)

Reza Zarei Sahamieh<sup>\*</sup>

Department of Geology, Faculty of Science, Lorestan University, Khorramabad, Iran

Submitted: Dec. 24, 2016 Accepted: June 12, 2017

Keywords: Calc-alkaline, Crustal contamination, Continental margin, Farmahin, Urumieh-Dokhtar Zone

### Introduction

The study area includes Alam Baghi, Vashaghan, Sar Band and Ghermez Cheshmeh and is located in the northeast of Farmahin and the southwest of Tafresh. Based on the structural subdivisions of Iran, the mentioned area is a part of Central Iran and the Urumieh-Dokhtar magmatic belt (Hajian, 1970).

The studied volcanic rocks consist of trachybasalt, trachyandesite, basaltic andesite, andesite, dacite, rhyodacite, rhyolite, ignimbrite, tuff and tuffit in composition and in terms of age they belong to the middle and upper Eocene. It seems that the volcanic activities are related to folding and faulting in the studied area. On the other hand, in addition to causing orogenic activity, at the middle and upper Eocene (Ghasemi and Talbot, 2006), locally extensional regime has played a main role in volcanic eruption. Similar to this scenario happened in other areas such as Taft and Khizrabad in Central Iran (Zarei Sahamieh et al., 2008). Porphyritic, microlite porphyritic and microlitic are the main textures in these rocks. they Mineralogically, contain plagioclase, clinopyroxene, amphibole, quartz and biotite as the main minerals and zircon, apatite, and opaque minerals as accessories.

## Materials and methods

The major and trace elements of mineral composition are determined by electron probe micro-analysis (EPMA) using a Cameca SX100 instrument in the Iran Mineral Processing Research Center (IMPRC). Moreover, the whole-rock major

and some trace elements analyses for a few samples were obtained by X-ray fluorescence (XRF), using an ARL Advant-XP automated X-ray spectrometer.

# Results

Based on EPMA analyses, plagioclase mineral in basaltic andesite and trachybasalt samples range from labradorite to bytownite in andesite and trachyandesite has oligoclase- andesine and in dacite, rhyodacite, rhyolite has an albite-oligoclase composition. In the Wo-En-Fs diagram, all clinopyroxenes show augitic and a lessor amount of clinoenstatite composition and in the O-J diagram located in the Mg-Fe-Ca (Quad) field and in the 2Ti+Cr+Al<sup>VI</sup> vs. Na+Al<sup>IV</sup> diagram (Morimoto et al., 1988) located above on the  $Fe^{3+}=0$  line that indicate high oxygen fugacity during crystallization. Microscopic study on these rocks such as oscillatory zoning, resorption rims in plagioclase and the presence of basic inclusions suggest the occurrence of magmatic contamination on the parent magma. The presence of oxidized amfibool rims (in hornbelende as oxy hornbelende) indicate the high temperature of the magma at the time of eruption.

According to the classification diagrams such as total alkaline vs.  $SiO_2$  (Irvine and Baragar, 1971) TAS (Le Bas et al., 1986) and tectonic discrimination diagrams (Pearce et al., 1984) samples are plotted in sub-alkaline, basaltic-andesite, andesite, dacite and rhyodacite, subduction and volcanic arc fields, respectively. The geochemical diagrams such as AFM are used

\*Corresponding authors Email: zareisah@yahoo.com; zarei.r@lu.ac.ir

for the identification of magma series and show that the studied rocks are calc-alkaline and A/NK *vs.* The A/CNK diagram shows the metaluminous to peraluminous nature. Incompatible and LIL elements such as Ba, K and Rb enrichment show that the contamination of magma with continental crust has occurred. The similarity between the REE patterns in all of the collected samples in Alam Baghi, Vashaghan, Sarban and Ghermez Cheshmeh areas suggest the same source for all of the volcanic rocks.

#### Discussion

The tectonic setting diagrams show that these rocks belong to the continental margin which has been involved in a subduction zone.

The position of the samples on the major elements vs. SiO<sub>2</sub> diagrams indicate that magma differentiation has occurred. Spider diagrams show a positive anomalous in Rb and a negative anomalous in Nb and Ti This phenomenon shows a contamination between the magma and the crustal rocks (Rollinson, 1993). Also, MORB-normalized incompatible element patterns of the Farmahin area show that the parent magma has been contaminated. It appears that assimilation and fractional crystallization (AFC) were the dominant processes in the genesis of the studied volcanic rocks.

As a conclusion and according to field evidence and geochemical characteristics presented in this article, the studied area is composed of lava flows and pyroclastic rocks such as andesite, dacite, rhyodacite, ignimbrite, tuff and tuffits that cross cut by younger dykes and belong to the middle to late Eocene age (middle to upper Lutetien). According to Sm/Yb vs. Sm diagram (Aldanmaz et al., 2000), all the studied samples in terms of composition are similar to enriched mantle-derived melts that are generated by varying degrees of partial melting (10% - 20%) from a spinel lherzolite to spinelgarnet lherzolite source.

Considering the evidences, all rocks in the studied area belong to the subduction zone and the parent magma originated from mantle and was contaminated with continental crust during eruption and rising.

# Acknowledgments

The authors wish to thank the Journal Manager and reviewers who critically reviewed the manuscript and made valuable suggestions for its improvement.

#### References

- Aldanmaz, E., Pearce, J.A., Thirlwall, M.F. and Mitchell, J.G., 2000. Petrogenetic evolution of late Cenozoic, post-collision volcanism in western Anatolia, Turkey. Journal of Volcanology and Geothermal Research, 102(1– 2): 67–95.
- Ghasemi, A. and Talbot, C.J., 2006. A new scenario for the Sanandaj-Sirjan zone (Iran). Journal of Asian Earth Sciences, 26 (6): 683–693.
- Hajian, J., 1970. Geological map of Farmahin, scale1:100000. Geological Survey of Iran.
- Irvine, T.N. and Baragar, W.R.A., 1971. A guide to the chemical classification of the common volcanic rocks. Canadian Journal of Earth Sciences, 8(5): 523–548.
- Le Bas, M.J., Le Maitre, R.W., Streckeisen, A. and Zanettin, B., 1986. A chemical classification of volcanic rocks based on the total alkali silica diagram. Journal of Petrology, 27 (3):745–750.
- Morimoto, N., Fabrise, J., Ferguson, A., Ginzburg, I.V., Ross, M., Seifert, F.A., Zussman, J., Akoi, K. and Gottardi G., 1988. Nomenclature of pyroxenes. American Mineralogist, 173(9– 10):1123–1133.
- Pearce, J.A., Nigel, B., Harris, N.B.W. and Tindle, A.G., 1984. Trace Element Discrimination Diagrams for the Tectonic Interpretation of Granitic Rocks. Journal of Petrology, 25(4): 956–983.
- Rollinson, H.R., 1993. Using Geochemical Data: Evaluation, Presentation and Interpretation. Longman scientific and technical, London, 352 pp.
- Zarei Sahamieh, R., Tabasi, H. and Jalali, M., 2008. Petrology and tectonomagmatic investigation of volcanic rocks of Ashtian. Journal of Science, Kharazmi University, 8(3):227–240. (in Persian with English abstract)