

## The Effects of cumulus cells on in vitro maturation of mouse germinal vesicle stage oocytes

Reza Mahmoudi,<sup>1</sup>Ph.D. student, Aligholi Subhani,<sup>1</sup>Ph.D., Parichehr Pasbakhsh,<sup>1</sup>Ph.D., Farid Abolhasani,<sup>1</sup>Ph.D., Iraj Amiri,<sup>2</sup>Ph.D., Mozdeh Salehnia,<sup>3</sup>Ph.D., Farideh Etesam<sup>1</sup>Ph.D.

1 Department of Anatomy, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

2 Department of Anatomy, Faculty of Medicine, Hamedan University of Medical Sciences, Hamedan, Iran.

3 Department of Anatomy, Medical School, Tarbiyat Modarres University. Tehran, Iran.

### Abstract

**Background:** In vitro maturation (IVM) of oocytes is a promising technique to reduce the costs and avert the side-effects of gonadotropin stimulation for in vitro fertilization (IVF). The pregnancy rates from oocytes matured in vitro are much lower than those of in vivo stimulation cycles, indicating that optimization of IVM remains a challenge.

**Objective:** In this study, we investigated the effect of cumulus cells on maturation and fertilization rate of immature oocytes (Germinal vesicle).

**Materials and Methods:** Germinal vesicle (GV) oocytes were recovered from 6-8 weeks old Balb C female mice 48hr after injection of 10 IU pregnant mare serum gonadotropin (PMSG). Collected oocytes were divided into two groups. Group A: GV oocytes without cumulus (denuded oocyte). Group B: GV oocytes with cumulus cells (cumulus-oocyte complex). The oocytes in both groups were cultured in TCM-199 medium in a humidified atmosphere of 5% CO<sub>2</sub> in air at 37°C. The maturation, fertilization and developmental rates were recorded after 24hr.

**Results:** Maturation, fertilization and developmental rates in denuded oocytes (DO) were 65.1%, 68.02%, 78.63% respectively, and in cumulus-oocyte complex (COC) were 78.20%, 85.57% and 85.05%, respectively. The maturation, fertilization and developmental rates of COC were significantly higher than those of DO (p<0.05).

**Conclusion:** The results show that cumulus cells have beneficial effects on maturation, fertilization and cleavage rates of mice oocytes.

**Key Words:** Immature oocyte, In vitro maturation, Mouse, Cumulus cells

### Introduction

In vitro maturation (IVM) of mammalian oocytes has become an efficient method to produce mature oocytes in order to use in assisted reproductive techniques such as In Vitro Fertilization (IVF), Intra cytoplasmic sperm injection (ICSI), and cloning. Induction of ovulation to obtain mature oocytes for IVF has become a routine procedure in many infertility clinics. Some women, however, rather fail to respond to the hormonal stimulation or are at risk of ovarian hyperstimulation (1). In vitro maturation of oocytes offers an alternative to obtain mature oocytes in these cases (2, 3). Immature human oocytes exhibit acceptable

meiotic competence to metaphase II (MII), but their subsequent developmental competence remains disappointingly low. Only 40–80% of fertilized IVM oocytes progress through early cleavage, and of those that do cleave and are transferred, 15% implant to form a viable fetus (3-6).

Oocyte maturation is often conceptually divided into nuclear and cytoplasmic processes. Nuclear maturation is a term that refers to the resumption of meiosis from the germinal vesicle (GV) stage and progression to MII. Cytoplasmic maturation is a more general term that refers to other maturational events (not directly related to meiotic progression) that prepares the oocyte for fertilization and preimplantation development (7, 8).

Within the follicle, the regulation of oocyte development and cellular function is dependent on gap junctional communication, paracrine communication and interaction with elements of extracellular matrix (9-11). Cumulus cells play an

### Corresponding Author:

Dr Aligholi Subhani, Department of Anatomy and Vali-e-Ashr Reproductive Health Research Center, Faculty of Medicine, Tehran University of Medical Sciences, Tehran, Iran.

E-mail: aligholi1177@yahoo.com

important role on oocyte maturation, since they provide and transfer several known and unknown factors that are essential for normal nuclear and cytoplasmic maturation of oocytes and subsequent embryonic development after fertilization (12-14). Moreover, cumulus cells also are known to play an important role in regulation of meiotic progression of oocytes (15). During the growth and accomplishment of meiotic competence of oocytes (before initiation of meiosis), cumulus cells are responsible for maintenance of nuclear arrest at GV stage via elevating intercellular cAMP level in the oocytes by transferring an inhibitor signal through gap junctions (16, 17). Initiation of meiosis is also related to cumulus-function, as there are evidences that cumulus cells secrete a meiosis-inducing factor (18).

According to the mentioned data, this study was proposed to investigate the effects of cumulus cells on in vitro maturation, fertilization and developmental rates of mouse GV stage oocytes.

## Materials and Methods


### Collection of GV oocytes

Oocytes were obtained from 6-8 weeks old Blab C female mice. The animal were kept under controlled condition (12hr light, 12hr dark) and fed with water and pellets ad libitum. The mice were stimulated by an i.p. injection of 10 IU pregnant mare serum gonadotropin (PMSG). The animals were killed 44hr later by cervical dislocation and the ovaries were removed into TCM-199 (Sigma) supplemented with 10% fetal bovine serum (FBS). The GV-stage oocytes of ovarian follicle were released by puncturing with a 28G sterile needle under a stereomicroscope. A total of 654 oocytes were obtained from 30 ovaries and they were used for in vitro maturation. The average number of collected oocytes was 21.8 per ovary.

### In Vitro Maturation (IVM)

The collected GV-stage oocytes were divided into two groups: Group I: denuded oocytes (DO) (Figure1A) and Group II cumulus oocyte complexes (COC) (Figure1B). Each group were placed in 25µl micro drops TCM-199 supplemented with 10% FBS, 0.23 mM sodium pyruvate, 50mg/l streptomycin, 60 mg/l penicillin and 1µg/l epidermal growth factor (EGF), overlaid with embryo-tested light mineral oil (Sigma) for 24hr in a humidified atmosphere of 5% CO<sub>2</sub> at 37°C. At various intervals from the onset of incubation, oocytes were observed by inverted microscopy and morphological changes in the

nucleus or the extrusion of first polar body (MII) were used as the criterion for nuclear maturation of GV-stage oocytes. After IVM, oocytes in group II were exposed to 3 mg/ml hyaluronidase in TCM199 and cumulus cell were removed by repeated pipetting and the matured oocytes were collected for in vitro fertilization.



**Figure 1.** In vitro maturation, fertilization, and cleavage of mouse germinal vesicle (GV) oocytes. (A) Cumulus oocyte complex. The attached cumulus-corona cells can be seen. (B). magnification:  $\times 320$ ; B $\times 200$ . Bar=32  $\mu$ m.

### In Vitro Fertilization (IVF)

Sperms were collected from epididymes of blab C male mice aged 12 weeks. The sperm suspension ( $1 \times 10^6$  motile spermatozoa/ml) was capacitated for 2hr in one ml T6 media supplemented with 16 mg/ml BSA. The in vitro matured MII stage oocytes from each treatment group were placed into 0.9 ml T6 and 0.1 ml capacitated spermatozoa was added to it. After 5hr incubation, the oocytes were washed through three droplets of T<sub>6</sub> medium. Then, the oocytes were

cultured in a droplet of T<sub>6</sub> (25μl) under mineral oil. They were assessed for cleavage to the 2-cells stage after 24hr.

**Statistical analysis**

Collected data were analyzed by chi-square test. The differences in the values of maturation, fertilization and developmental rates, were considered significant when p<0.05.

**Results**

**In vitro maturation of mouse oocytes**

Table I shows the number of oocytes arrived to MII stage (Figure 2C) after 24hr culture. The maturation rate of COC was significantly higher than that of DO (p<0.05).



**Figure 2C.** Denuded GV stage oocyte. magnification: ×320: B×200. Bar=32 μm.

**In vitro fertilization and development of mouse oocytes**

As shown in Table II, the number of fertilized oocytes (Figure 3D) were significantly higher in COC than those of DO (p<0.05). But the difference between the number of fertilized oocytes which

developed to 2-cells stage (Figure 3E) was not significant (p>0.05) in two groups.

**D**



**E**



**Figure 3D&E.** Completion of the second meiotic division (MII) after 24h culture. (D). A fertilized 2-pronuclear zygote (E). A cleaved embryo at 24h post fertilization. All pictures taken under Hoffman contrast optics. Original magnification: ×320: B×200. Bar=32 μm.

**Table I:** Maturation rate of mouse oocytes after 24hr culture. DO = Denuded oocyte, COC = Cumulus oocyte complex, GV= Germinal vesicle oocyte. GVBD= Germinal vesicle breakdown, MII=metaphase II. Significant differences between DO and COC are indicated by an asterisk (p<0.05).

| Group   | No. of GV stage oocytes | Final stage oocyte maturation |            |                  |
|---------|-------------------------|-------------------------------|------------|------------------|
|         |                         | GVBD (%)                      | MI I (%)   | Degeneration (%) |
| A (DO)  | 264                     | 25 (9.5)                      | 172 (65.1) | 67 (27.4)        |
| B (COC) | 390                     | 24 (6.1)                      | 305 (78.2) | 61 (15.7)        |

**Table II:** Fertilization and developmental rates of mouse oocytes. Significant differences between DO and COC are indicated by an asterisk (P<0.05).

| Group   | Matured (MII) oocytes | Fertilized Oocytes (%) | Degenerated Oocytes (%) | 2-cell stage Embryos (%) | Degenerated Embryos (%) |
|---------|-----------------------|------------------------|-------------------------|--------------------------|-------------------------|
| A (DO)  | 172                   | 117 (68.02)            | 55(31.97)               | 92 (78.63)               | 25(31.97)               |
| B (COC) | 305                   | 261(85.57)             | 44(14.42)               | 222(85.05)               | 39(14.95)               |

## Discussion

Cumulus cells have been considered to play an important role in oocyte maturation by keeping the oocyte under meiotic arrest, inducing meiotic resumption and by supporting cytoplasmic maturation. These functions have been attributed to their gap junctions and their specific metabolizing capabilities (12-15). Physical contact between oocyte and cumulus cells has been considered necessary for the transfer of nutrients and factors essential for oocyte development (16-19). In addition, dissociated cumulus cells have been reported to produce paracrine factors, which resume meiosis in denuded oocytes (20). The beneficial effects of cumulus cells on oocyte maturation and early development were reported in different species (12, 14, 19-23). Our study confirmed these mentioned results and demonstrated that maturation and fertilization rates were significant by higher in cumulus-oocyte complex compared with denuded oocytes ( $p < 0.05$ ). But there was no significant difference in cleavage rates. However, the obtained data in this study are different with some other published results. For example Nishi *et al* (2003) investigated on the mice GV oocytes and reported that maturation, fertilization and developmental rates of DO were 64.4, 51.7 and 43.2%, and cumulus-oocyte complex (COC) were 76.4, 70.5 and 50.5%, respectively. The maturation rate (MII) and fertilization rate (2PN) of COC was significantly higher than that of DO ( $p < 0.05$ ) (25). On the other hand, Aono *et al* (2003) investigated on murine GV oocyte and reported that the maturation (MII) and cleavage rate (2-cell) of COC were 90.2%, 94.1% respectively (25). In our study, these amounts are higher than Nishi *et al* investigation in both COC and DO, and lower than that of Aono *et al*. This difference maybe is due to the animal, oocyte collection method and medium that used.

## Conclusion

In conclusion, our study indicates that in vitro maturation and fertilization and cleavage rates of mouse cumulus-oocyte complex (COC) are more successful than denuded oocytes. Further studies should explore the factors in cumulus cells that are responsible for these results. We think by exploring and extracting these factors and using them in culture media, oocytes maturation and embryonic preimplantation development in vitro can be improved.

## References

- Child TJ, Abdul-Jalil AK, Gulekli B, Tan SL. In vitro maturation and fertilization of oocytes from unstimulated normal ovaries, polycystic ovaries, and women with polycystic ovary syndrome. *Fertile Sterile* 2001; 76: 936-942.
- Cha KY, Chian RC. Maturation in vitro of immature human oocytes for clinical use. *Hum Reprod Update* 1998; 4: 103-120.
- Hardy K, Wright CS, Franks S, Winston RM. In vitro maturation of oocytes. *Br Med Bull* 2000; 56: 588-602.
- Chian RC, Lim JH, Tan SL. State of the ART in in vitro oocyte maturation. *Curr Opin Obstet Gynecol* 2004; 16: 211-219.
- Moor RM, Dai Y, Lee C, Fulka J. Oocyte maturation and embryonic failure. *Hum Reprod Update* 1998; 4: 223-236.
- Trounson A, Anderiesz C, Jones G. Maturation of human oocytes in vitro and their developmental competence. *Reproduction* 2001; 121: 51-75.
- Motlik J. Cytoplasmic aspects of oocyte growth and maturation in mammals. *J Reprod Fertile* 1989; 38(Suppl.): 17-25.
- Combelles CM, Cekleniak NA, Racowsky C, Albertini DF. Assessment of nuclear and cytoplasmic maturation in in vitro matured human oocytes. *Hum Reprod* 2002; 17: 1006-1016.
- Sutton ML, Gilchrist B, Thompson G. Effects of in-vivo and in vitro environments on the metabolism of the cumulus-oocyte complex and its influence on oocyte developmental capacity. *Hum Reprod Update* 2003; 9: 35-48.
- Roberts R, Franks S, Hardy K. Culture environment modulates maturation and metabolism of human oocytes. *Hum Reprod* 2002; 17: 2950-2956.
- Albertini F, Combelles M, Benecchi E, Carabatsos MJ. Cellular basis for paracrine regulation of ovarian follicle development. *Reproduction* 2001; 121: 647-653.
- Warriach M, Chohan KR. Thickness of cumulus cell layer is a significant factor in meiotic competence of buffalo oocytes. *J Vet Sci* 2004; 5 (3): 247-51.
- Tanghe S, Soom AV, Nauwynck H, Coryn M, Dekruif A. Minireview: function of the cumulus ophorous during oocyte maturation, ovulation and fertilization. *Mol. Reprod Dev* 2002; 61: 414-424.
- Janssenswillen C, Nagy ZP, Van Steirteghem A. Maturation of human cumulus-free germinal vesicle-stage oocytes to metaphase II by Coculture with monolayer Vero cells. *Hum Reprod* 1995; 10: 375-376.
- Downs SM. A gap-junction-mediated signal, rather than an external paracrine factor, predominates during meiotic induction in isolated mouse oocytes. *Zygote* 2001; 9: 71-82.
- Eppig JJ, Schultz RM, O'Brien M, Chesnel F. Relationship between the developmental programs controlling nuclear and cytoplasmic maturation of mouse oocytes. *Dev Biol* 1994; 164: 1-9.
- Shimada M, Terada T. FSH and LH induce progesterone production and progesterone receptor synthesis in cumulus cells, a requirement for meiotic resumption in porcine oocytes. *Mol Hum Reprod* 2002; 8: 612-618.
- Xia GL, Kikuchi K, Noguchi J, Izaike Y. Short time priming of pig cumulus-oocyte complexes with FSH and forskolin in the presence of hypoxanthine stimulates cumulus cells to secrete a meiosis-activating substance. *Theriogenology* 2000; 53: 1807-1815.
- Motlik J, Nagai T, Kikuchi K. Resumption of meiosis in pig oocytes cultured with cumulus and parietal granulosa cells: The effect of protein synthesis inhibition. *J Exp Zool* 1991; 259: 386-391.

20. Vanderhyden BC, Armstrong, DT. Role of cumulus cells and serum on the in vitro maturation, fertilization and subsequent development of rat oocytes. *Biol Reprod* 1989; 40: 720–728.
21. Isobe N, Terada T. Effect of the factor inhibiting germinal vesicle breakdown on the disruption of gap junctions and cumulus expansion of pig cumulus-oocyte complexes cultured in vitro. *Reproduction* 2001; 121: 249–257.
22. De Loos, Zeinstra E, Bevers MM. Follicular wall maintains meiotic arrest in bovine oocytes cultured in vitro. *Mol Reprod Dev* 1994; 39: 162–165.
23. Motta PM, Makabe S, Naguro T, Correr S. Oocyte follicular cells association during development of human ovarian follicle. A study by high resolution scanning and transmission electron microscopy. *Arch Histol Cytol* 1994; 57: 369–394.
24. Nishi Y, Takeshita T, Sato K, Araki T. Change of the mitochondrial distribution in mouse ooplasm during in vitro maturation. *J Nippon Med Sch* 2003; 70(5): 408-15.
25. Aono N, Naganuma T, Abe Y, Hara K, Sasada H, Sato E, Yoshida H. Successful production of blastocysts following ultrarapid vitrification with step-wise equilibration of germinal vesicle-stage mouse oocytes. *J Reprod Dev* 2003; 49 (6): 501-506.

Archive of SID