

Ultrastructural and morphometrical study of preimplantation endometrium in superovulated mice treated with progesterone or Sildenafil

Leila Roshangar^{1,2} Ph.D., Jafar Soleimani-Rad² Ph.D., Bahman Rashidi³ Ph.D., Hossein Mazochian⁴ M.D. Student, Behzad Nikzad⁵ M.D. Sara Soleimani Rad⁶ M.D.

1. Neuroscience Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

2. Drug Applied Research Center, Faculty of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran.

3. Department of Anatomical Sciences, Faculty of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran.

4. Medical Education Research Center, Tabriz University of Medical Sciences, Tabriz, Iran.

5. Department of Neuroscience, Tabriz University, Tabriz, Iran.

6. Faculty of medicine, Tabriz University of Medical Sciences, Tabriz, Iran.

Corresponding Author:

Jafar Soleimani Rad, Department of Anatomical Sciences, Faculty of Medicine, Tabriz University of Medical Sciences, Valiasr, Tavaneer Ave. No 7, Tabriz, Iran.
Email: SoleimaniRJ@yahoo.com
Tel: (+98) 9143154736

Received: 19 May 2012
Revised: 13 January 2013
Accepted: 8 June 2013

Abstract

Background: Endometrial development has an important role in blastocyst adhesion and implantation. During IVF cycles, endometrial development is enhanced by progesterone.

Objective: The aim of this study was to compare ultrastructural and morphometrical characteristics of mice uterine endometrium in natural cycle with those in superovulated cycles received progesterone or Sildenafil.

Materials and Methods: In This study, 60 female bulb/c mice were divided into 4 groups: a control and 3 experimental; gonadotropin, gonadotropin+ Sildenafil and gonadotropin+ progesterone. In experimental groups the mice superovulated mated. In the gonadotropin+ progesterone and gonadotropin+ Viagra groups, the mice respectively received 1mg progesterone and 3 mg Sildenafil citrate. Their uterine specimens were prepared for morphometrical and ultrastructural study. Height of the epithelial cells was measured, using motic software. Statistical analysis was performed using ANOVA.

Results: Microscopy revealed that in control group the cells had numerous apical microvilli and the height of the cells was $20.52 \pm 2.43 \mu\text{m}$. In gonadotropin+ progesterone group, the granules were found in basal and apical portions and cellular height were $17.91 \pm 2.78 \mu\text{m}$ which were significantly shorter than in the control and gonadotropin groups ($p < 0.001$). In this group, the apical membrane also contained pinopodes. In gonadotropin +Sildenafil group, the granules were found in both apical and basal portions and the height of the cells were $17.60 \pm 2.49 \mu\text{m}$ which were significantly shorter than in the control and gonadotropin groups ($p < 0.001$). In this group, pinopodes appeared slightly extensive than the other groups.

Conclusion: It is concluded that superovulatory drugs in mice stimulate endometrial maturation but injection of Sildenafil is nearly more positive.

Key words: Implantation, Progesterone, Sildenafil, Mice, Uterine.

This article extracted from Ph.D. thesis. (Bahman Rashidi)

Introduction

The low pregnancy rate in assisted reproductive technology (ART), is one of challenging issues which extensively is under investigation. Since the beginning of use of in vitro fertilization for treatment of infertility, the superovulatory methods and drugs and embryo culture media has progressed tremendously. However, the implantation rate in ART is still low (1, 2). It appears that low implantation rate is partly due to the interference of super ovulatory drugs with endometrial maturation and also lack of

synchronicity between endometrial and blastocysts development (3-6).

In ART protocols, in addition to the use of superovulatory drugs, for acceleration of endometrial maturation, after oocyte collection, progesterone is also used. The effect of this hormone on endometrial receptivity is established and reported that progesterone could extends endometrial implantation period possibly by modulation of uterine cell proliferation and expression of genes involved in implantation (7-16). Sildenafil citrate (Viagra) is a drug that regarding to its balancing effect between contraction and relaxation, in smooth muscle,

is primarily used for erectile dysfunction (17-25).

There are evidence that Sildenafil citrate is a vasodilator which act by increasing intracellular nitric oxide (NO) and it also enhances vasodilatory effect of NO by inhibiting the degradation of cGMP (26-28). Similar actions are proposed for Viagra in the myometrium and it is shown that vaginal use of Sildenafil improves uterine artery blood flow (29-31). Therefore, it appears that Viagra by increasing endometrial circulations could facilitate endometrial development and receptivity (32). Morphological characteristics of endometrium, especially endometrial luminal epithelium at preimplantation period, are correlated well with its receptivity at implantation period (6).

Consequently, it could be used as valuable criteria for evaluating endometrial development. The aim of the present study was to compare morphological characteristics of endometrium such as extension of pinopods, height of endometrial epithelial cells and location of secretory granules as an induction of endometrial development in superovulated mice after receiving progesterone or Viagra and immediately before implantation.

Materials and methods

Animals and treatment

In this experimental study, 60 adult female bulb/c mice with average weight of 25-30 gr and 30 adult male mice were used. The mice housed at room temperature under a standard condition, light cycle (12-h light/dark) and free access to food and water. The ethical committee of Tabriz University of Medical Sciences (TUMS) approved all aspects of this study following laboratory practice guidelines. The female mice were divided into 4 groups of 15 as: control, gonadotropin, gonadotropin+ progesterone and gonadotropin+ Sildenafil.

Gonadotropin was purchased from IBSA institute biochimique SA, Switzerland and Sildenafil was purchased from Rooz Daroo Co. Tehran, IRAN. Except in control group, in other experimental groups the mice received

7.5 IU hCG as intraperitoneal injection and 48 hours later 7.5 IU hMG. Then in all groups, two female mice at their oestrous cycle were put with one male mouse in a cage for mating. The presence of vaginal plug in the next morning was designated as 0 day post coitum (d.p.c).

In the gonadotropin+ progesterone and gonadotropin+ Sildenafil groups, the mice respectively received 1mg progesterone and 3mg Sildenafil citrate at 24, 48 and 72 hours after HMG injection. 96 hours after HMG injection using 23 gauge needles, the mice in experimental groups together with control mice were sacrificed and uterine specimens were obtained and processed for microscopic evaluation.

Light microscopy

For having uterine specimens exactly at preimplantation stage, the specimens only from those that their uterine contained blastocyst were fixed in 10% formalin, embedded in paraffin and 5 µm thick sections were stained with H & E and PAS and studied with light microscope. For measurement of height of epithelial cells, the images were transferred to monitor, using Motic image analyzer system plus 2, and the measurement was carried out at higher magnification as which was shown in previous study (33).

Transmission electron microscopy

Uterus from both control and experimental group fixed in 2.5% glutaraldehyde (Proscitech, Thuringowa, Australia) in phosphate buffer and processed for transmission electron microscopy at Tabriz University of medical sciences, drug applied and research center, histology department EM facility.

Samples were postfixed in 1% OsO₄ (TAAB, Berkshire- UK) dehydrated through an ethanol series, equilibrated in propylene oxide, and embedded in Araldite (Proscitech, Thuringowa, Australia). Thin sections were stained with uranyl acetate and lead citrate. The specimens were studied with LEO 906 transmission electron microscope for pinopodes morphology of nuclei and

localization of granules. The images recorded with AMT advantage plus CCD camera.

Statistical analysis

The values in every time point from control and treated groups were analyzed with ANOVA using SPSS 13 and the level of $p < 0.05$ was considered significant.

Results

Light microscopic study revealed that the cells of endometrial luminal epithelium were tall columnar in both control and those that only received HMG and HCG. But the cells were low columnar in those groups that received progesterone or Viagra in addition to superovulatory drugs. On the other hand, in the first two groups, PAS positive granules were restricted to sub nuclear and supranuclear regions, while in the experimental groups which were received progesterone or Viagra, they were dispersed in the cytoplasm. Electron microscopy confirmed the light microscopic findings and showed that in control group the endometrial luminal cells contained euchromatic nuclei,

several basal granules, numerous mitochondria and a well-developed rough endoplasmic reticulum (Figure 1, 2).

The cells had also several apical microvilli (Figure 2). The HMG-HCG group had euchromatic nuclei and granules were more localized at apical region and pinopodes were frequent at luminal surface (Figure 3). In the group that received progesterone, in addition to HMG+HCG, granular localization was similar to HMG-HCG group but apical microvilli were as in control group (Figure 4). In the group that received Viagra, instead of progesterone, the apical granules and pinopodes were much similar to those in superovulated group (Figure 5) and several phagolysosomes were present in the stromal cells (Figure 6). The results of morphometric study are summarized in table I.

As it is shown in the table the height of the endometrial epithelial cells in the control group was $20.52 \pm 2.43 \mu\text{m}$ and in gonadotropin group was 22.64 ± 1.64 . In gonadotropin+progesterone group, the heights of the cells were $17.91 \pm 2.78 \mu\text{m}$ and in gonadotropin+sildenafil was 18.23 ± 0.71 . The differences were statistically significant ($p < 0.001$).

Table I. The height of the endometrial epithelial cells (μm) in control and experimental groups

Groups	Control	Gonadotropin	Gonadotropin + progesterone	Gonadotropin+ Sildenafil
height of the endometrial cells	20.52 ± 2.43	$22.64 \pm 1.64^*$	$17.91 \pm 2.78^*$	$18.23 \pm 0.71^*$

*: The values indicated with mean \pm SD were significantly ($p < 0.001$) different from control group.




Figure 1. Electron micrograph from epithelial cell of endometrium in control mouse. Note basal lamina (BL), Nucleus (N), mitochondrion (M), granules (G), smooth endoplasmic reticulum (SER). 13700 X.




Figure 2. Electron micrograph from endometrial epithelial cells from control mouse. Note, Junctional complex (JC), mitochondria (M), microvilli (MV). 13500X.




Figure 3. Electron micrograph from endometrial epithelial cells from the group received gonadotropin. The micrograph shows Pinopods (P), mitochondria (M), and numerous apical granules (G). 6646X.




Figure 4. Electron micrograph from endometrial epithelial cells from experimental group received gonadotropin+progesterone. Note, Junctional complex (JC), several mitochondria (M), Nucleus (N) and numerous supranuclear granules (G). 13500X.




Figure 5. Electron micrograph from endometrial epithelial cells from experimental group received gonadotropin+Viagra. Note, the presence of extensive pinopods (P) and euchromatic Nuclei (N). 9500X.




Figure 6. Electron micrograph from endometrial epithelial and stromal cells from experimental group received gonadotropin+progesterone. Note, the presence of intracellular phagolysosomes. (G) 13500X.

Discussion

The results of the present study showed that the height of the endometrial luminal epithelium in control and HMG-HCG groups were higher than the height of cells in gonadotropin+ progesterone or Viagra groups. This is probably due to the stimulatory effect of gonadotropin on folliculogenesis which could result in higher estrogen level. The hypothesis of progesterone injection in IVF cycles is that, endometrial epithelial maturation is facilitated by progesterone

secretion during secretory phase. However, it appears that early progesterone injection interfere with the effect of estrogen and affect the height of the cells.

In the support of our finding it is shown that estrogen results in hyperplasia, hypertrophy and increasing of cell height (34). On the other hands high level of progesterone is accompanied by decreased cellular height in endometrial luminal epithelium (35-40). Ultrastructural studies showed that apical granules and pinopodes on luminal surface of endometrial epithelium were more frequent in

gonadotropin and gonadotropin+ Viagra groups in comparison to control and progesterone groups. Taking the above morphological features as indications of endometrial maturation, it is indicated that gonadotropin accelerates the endometrial maturation and receptivity.

While progesterone slightly suppresses the stimulatory effect of gonadotropins (41). Viagra has almost not any negative effect on gonadotropin-induced changes in endometrial epithelium. Similarly it is shown that superovulation protocol accelerates development of implantation window (42). The results of the present study are unique in that, the morphological changes of the endometrium are studied immediately, before implantation.

A condition that its study in human is impossible (43-45). To make sure that all the samples are from the same stage of development. The uterine was flushed and specimens was obtained only from those that blastocyst could be collected.

Conclusion

It is concluded that administration of Sildenafil rather than progesterone in superovulation protocol may have positive effect on implantation. However further studies and detection of molecules, involved in endometrial receptivity and implantation, would be needed to come to a definite conclusion.

Acknowledgments

The authors are grateful to research department of Tabriz University of Medical Sciences for their financial support.

Conflict of interest

The authors declare that there is no conflict of interest.

References

1. Keye W Chang R. Infertility evaluation and treatment. W.B. Saunders company, Philadelphia; 1995: 115-126.
2. Landgren B, Johannisson E. A new method to study the process of implantation of a human blastocyst is in vitro. *Fertil Steril* 1996; 65: 1967-1970.
3. Can A, Tekebiogbu M, Biberoglu K. Structure of premenstrual endometrium in HMG/HCG induced an ovulatory women. *Eur J Obstet Gynecol Reprod Biol* 1991; 42: 119-124.
4. Csemiczky G, Wramsly H, Johannisson E. Importance of endometrial quality in women with tubal infertility during a natural menstrual cycle for the outcome of IVF treatment. *J Assist Reprod Genet* 1998; 15: 55-61.
5. Bourgain C, Smitz J, Camus M. Human endometrial maturation is markedly improved after luteal supplementation of GnRH/ HMG stimulated cycles. *Hum Reprod* 1994; 9: 32-40.
6. Narkar M, Kholkute S, Chitlange S, Nandedkar T. Expression of hormone receptors, proliferation and apoptotic markers in primate endometrium. *Mol Cell Endocrinol* 2006; 246: 107-113.
7. Hewitt SC, Korach KS. Progesterone action and responses in the ERKO mouse. *Steroid* 2000; 65: 551-557.
8. Sengupta J, Ghosh D. Role of peri-implantation stage endometrium-embryo interaction in the primate. *Steroid* 2000; 45: 753-762.
9. Bourgain C, Devroey P. The endometrium in stimulated cycles for IVF. *Hum Reprod Update* 2003; 9: 515-522.
10. Fossum GT, Davidson A, Paulson RJ. Ovarian hyperstimulation inhibits embryo implantation in the mouse. *In vitro Fert Embryo Transfer* 1989; 6: 7-10.
11. Basir GH, Wai-sum O, Hung yu, Ng E, Chung Ho P. Morphometric analysis of pre-implantation endometrium in patients having excessively high oestradiol concentration after ovarian stimulation. *Hum Reprod* 2002; 16: 435-440.
12. Ertzeid GStoreng R. The impact of ovarian stimulation on implantation and fetal development in mice. *Hum Reprod* 2001; 16: 221-225.
13. Haengseok S, Kyuyong H, Hyunjung L. Progesterone supplementation extends uterine receptivity for blastocyst implantation in mice. *Reproduction* 2007; 133: 487-493.
14. Lindhard A, Bentin-Ley U, Ravn VIslin H. Biochemical evaluation of endometrial function at the time implantation. *Fertil Steril* 2002; 78: 221-233.
15. Jelink J, Ylikorkala O, Jarvinen PA, Alapiessa U. Effect of endogenous and exogenous progesterone on human endometrial enzymes. *Int J fertil* 1978; 23: 23-37.

16. Bussi M, Murphy CR. Hormonal control of enzyme activity during the plasma membrane transformation of uterin epithelial cell. *Cell Biol Int* 2001; 25: 859-871.
17. Harrold LR, Gurwitz JH, Field TS. The diffusion of a novel therapy into clinical practice: The case of Sildenafil. *Arch Intern Med* 2000; 160: 340-3405.
18. Bivalacqua J, Champion HC, Hellstrom WJ, Kadowitz PJ. Pharmacotherapy for erectile disfunction. *Trends Pharmacol Sci* 2000; 21: 484-489.
19. Wallis RM, Corbin JD, Francis SH, Ellis P. Tissue distribution of phosphodiesterase families and the effects of Sildenafil on tissue cyclic nucleotides, platelet function, and the contractile responses of trabeculae carneae and aortic rings in vitro. *Am J Cardiol* 1999; 83: 3-12.
20. Andersson KE, Wanger G. Physiology of erection. *Physiol Rev* 1995; 75: 191-236.
21. Burnett A. Nitric oxide in the penis: physiology and pathology. *J Urol* 1997; 157: 320-324.
22. Christ GJ, Richards S, Winkler A. Integrative erectile biology: the role of signal transduction and cell-to-cell communication in coordinating corporal smooth muscle tone and penile erection. *Int J Impot Res* 1997; 9: 69-84.
23. Giuliano FA, Rampin O, Benoit G, Jardin A. Neural control of penile erection. *Urol Clin North Am* 1995; 22: 747-766.
24. Porst H. The rational for prostaglandin E1 in erectile failure: a survey of worldwide experience. *J Urol* 1996; 155: 802-815.
25. Saenz de Tejada I. In the physiology of erection, signposts to impotence. *Contemp Urol* 1992; 7: 52-68.
26. Ballard SA, Gingell CJ, Tang K. Effects of Sildenafil on the relaxation of human corpus cavernosum tissue in vitro and on the activities of cyclic nucleotide phosphodiesterase isohyets. *J Urol* 1998; 159: 2164-2171.
27. Chuang AT, Strauss JD, Murphy RA, steers WD. Sildenafil, a type Cgmp Phospdiesterase inhibitor, specifically amplifies endogenous Cgmp-dependent relaxation in vitro. *J Urol* 1998; 160: 257-61.
28. Watanabe N, Kabasawa Y. 4-Benzylamino 1-chloro 6-substituted phthalazines: Synthesis and inhibitory activity toward phosphodiesterase-5. *J Med Chem* 1998; 41: 3367-3372.
29. March D. Viagra (Sildenafil) effectively treats enlarged hearts, mouse study show. Johns Hopkins Medical Institutions. Medical Video News 2005. Available at: http://www.hopkinsmedicine.org/Press_releases/2005/01_23_05.htm.
30. Wareing M, Myers JE, Ohara M, Baker PN. Sildenafil citrate (Viagra) enhances vasodilatation in fetal growth restriction. *J Clin Endocrinol Metab* 2007; 90: 2550-2555.
31. Khan RN, Hamoud H, Warren A, Wong LF, Alurkumalan S. Relaxant action of Sildenafil citrate (Viagra) on human myometrium of pregnancy. *Am J Obstet Gynecol* 2004; 191: 315-321.
32. Paulus WE, Strehler E, Zhang M, Jelinkova L, Danasouri E, Isterzik K. Benefits of vaginal Sildenafil citrate in assisted reproduction therapy. *Fertil Steril* 2002; 77: 856-857.
33. Roshangar L, Soleimani Rad J, Nikpoo P, Sayyah Melli M. Effect of oxytocin injection on folliculogenesis, ovulation and endometrial growth in mice. *Iran J Reprod Med* 2009; 7: 91-95.
34. Hosi J, Murphy CR. A scanning and light microscope study comparing the effect of clomiphene citrate. Estradiol 17beta and progesterone on the structure of uterine luminal epithelial cell. *Eur J Morphol* 1995; 33: 39-50.
35. Li T, Rogers A, Dockery P. A new method of histologic dating of human endometrium in the luteal phase. *Fertil Steril* 1988; 50: 52-60.
36. Sarani S, Ghaffari-Novin M, Warren M. Morphological evidence for the "implantation window" in human laminal endometrium. *Hum Reprod* 1999; 14: 3101-3106.
37. Dockery P, Rogers A. The effect of steroids on the fine structure of the endometrium. *Bailliere's Clin Obstet Gynecol* 1989; 3: 227-247.
38. Gunian A. Realization of estradiol effects in the uterus of ovariectomized rats under acute stress. *Eur J Obstet Gynecol Reprod Biol* 1995; 60: 69-74.
39. Karmer B, Wet GD. Exogenous gonadotropin administration affects the glycocalyx of rat endometrial epithelium cell during the period of implantation. *J Assist Reprod Genet* 1994; 11: 405-509.
40. Risek B, Klier FG, Phillips A, Hahn D, W Gilula NB. Gap junction regulation in the uterus and ovaries of immature rats by estrogen and progesterone. *J Cell Sci* 1995; 108: 1017-1032.
41. Tavaniotou A, Smits J, Bougain C, Devroey P. Ovulation induction disrupts luteal phase function. *Ann N Y Acad Sci* 2001; 943: 55-63.
42. Stein B, Kramer B. The effect of exogenous gonadotropic hormones on the endometrium of the rat. *H Ant* 1989; 164: 123-140.
43. Kramer B, Magan A, DeWet G. Hyperstimulation affect vascular permeability at implantation site in the rat endometrium. *J Assist Reprod Genet* 1993; 10: 163-168.
44. Dursun A, Sendag F, Terek MC. Morphometric changes in the endometrium and serum leptin levels during the implantation period of the embryo in the rat in response to exogenous ovarian stimulation. *Fertil Steril* 2004; 82: 1121-1126.

45. Salehnia M, Arianmanesh M, Beigi M. The impact of ovarian stimulation on mouse endometrium: a morphometrical study. *Iran J Reprod Med* 2006; 4: 7-11.

Archive of SID