### Alternation of apoptotic and implanting genes expression of mouse embryos after re-vitrification

Nasrin Majidi Gharenaz M.Sc., Mansoureh Movahedin Ph.D., Zohreh Mazaheri Ph.D., Shahram Pour beiranvand M.Sc.

Department of Anatomical	Abstract
Sciences, Faculty of Medical Sciences Tarbiat Modares	Background: Nowadays, oocytes and embryos vitrification has become a routine
University, Tehran, Iran.	technique. Based on clinical judgment, re-vitrification maybe required. But little is
	known about re-vitrification impact on genes expression.
	<b>Objective:</b> The impact of re-vitrification on apoptotic and implanting genes, Bax,
	<i>Bcl-2</i> and <i>ErbB4</i> , at compaction stage embryos were evaluated in this study.
	Materials and Methods: In this experimental study, 8 cell embryos (n=240) were
	collected from female mature mice, 60-62 hr post HCG injection. The embryos were
	divided randomly to 3 groups included: fresh (n=80), vitrified at 8 cell stage (n=80),
	vitrified at 8 cell stage thawed and re-vitrified at compaction stage (n=80). Embryos
	were vitrified by using cryolock, (open system) described by Kuwayama. Q-PCR
	was used to examine the expression of Bax, Bcl2 ErbB4 genes in derived
	blastocysts.
	Results: Our result showed that expanded blastocyst rate was similar between
	vitrified and re-vitrified groups, while re-vitrified embryos showed significant
	decrease in expanded blastocyst rate comparing with fresh embryos (p=0.03). In
~	addition, significant difference was observed on apoptotic gene expression when
Corresponding Author: Mansourah Movahadin Jalala	comparing re-vitrified and fresh embryos (p=0.004), however expression of Bax and
Ale-Ahmad Highway, Tarbiat	Bcl-2 (apoptotic) genes didn't demonstrate a significant difference between re-
Modares University, Faculty of	vitrified and vitrified groups. The expression rate of ErbB4, an implantation gene
Medical Sciences, Department of	was decreased in re-vitrified embryos comparing with fresh embryos (p=0.003), but
Anatomical Sciences, Tehran, Iran. P O Box: 14115-175	it was similar between re-vitrified and vitrified embryos.
Email: movahed.m@modares.ac.ir	Conclusion: Re-vitrification can alter the expression of Bax, Bcl-2 and ErbB4 genes
Tel: (+98) 21 82884502	and developmental rate of mouse embryos in compaction stage.

Received: 26 November 2015 Revised: 24 May 2016 Accepted: 1 June 2016 Key words: Vitrification, Embryo, Apoptosis, Gene This article extracted from M.Sc. thesis. (Nasrin Majidi Gharenaz)

### Introduction

ocyte and embryo cryopreservation can contribute to a reduction of ovarian hyper stimulation syndrome improvement (OHSS), of cumulative pregnancy rates and reduction of multiple gestations during assisted reproductive (1-4). techniques Furthermore, several findings also confirm a better obstetrical outcome of children being born after a frozen embryo transfer (FET) (5-7). Additionally, this method is useful to fertility preservation of patients who are candidate for chemotherapy or radiotherapy (8).

Occasionally, unexpected extra embryos are available after thawing the embryos and

frozen-thawed re-cryopreservation after embryos transfer is inevitable (9). These embryos could be re-vitrified for a future transfer. It has been revealed that previously frozen embryos during in vitro fertilization (IVF) could be thawed for genetic evaluation and re-vitrified for transfer at a later date (10). Providing the option to re-cryopreserve a patient's previously stored embryos, may encourage more patients to take a cautious approach with regard to number of embryos transferred. et al reported first Farhat successful pregnancy following clinical transfer of re-vitrified embryos (11).

Ito *et al* evaluated impact of re-vitrification embryos (2 cell stage, 4 cell stage, and morula, blastocyst stage) on the same stage on the embryos developmental competence. They reported re-vitrification with cryotop did not affect the developmental ability of mouse embryos (12). Shehan also reported similar results. But they did not evaluate the revitrification effect on gene expression. Severe injuries may occur at all phases of these procedures (13). Hence, cryopreservation may cause fragmentation in frozen thawed embryos (14-16). Moreover, profile of apoptotic and implanting genes can be affected by vitrification procedure (17, 18).

publications postulated Several that vitrification could induce apoptosis in vitrified embryos and alter the expression of apoptosis related genes (17-20). Members of Bcl-2 gene family are known to have an important role in regulating apoptosis. Bcl-2 is known for its anti-apoptotic activity and responsibility in cell survival, whereas Bax is pro-apoptotic and responsible for the induction of cell death (20). ErbB4 is a gene involved in implantation which interacts with heparin binding-epithelial arowth factor (HB-EGF) to mediate attachment of the blastocyst to uterine luminal epithelium (21, 22). There is insufficient data on the effect of repeated use of vitrification and slow freezing on genes expression; there have been several studies on the efficiencies of re-vitrification and double slow freezing on embryo viability (11, 2325).

Therefore re-vitrification effects on developmental rate and *Bax, Bcl-2* and *ErbB4* genes expression pattern were investigated.

### Materials and methods

This experimental study was performed on mouse embryos. Female NMRI mice (6-8 wks) were purchased from the Pasteur Institute (Tehran, Iran). The mice were housed in a room under standard laboratory conditions (12 hr light/dark at 22°C) with free access to water and standard food. The current study was conducted under the protocol approved by the animal experimentation committee of Medical Sciences Faculty in Tarbiat Modares University.

### Embryo collection and culture

Mice were super ovulated by an intraperitoneal (IP) injection of 7.5 IU pregnant mare's serum gonadotropin (PMSG; Intervet Inc, Netherlands) followed 48 hr later by 7.5 IU human chorionic gonadotropin (HCG; Pregnil; Organon, Oss, Netherlands). The mice were mated with a single male of the same age (6-8 wks old). Presence of vaginal plugs was a confirmation of a pregnancy. The pregnant female mice sacrificed by cervical dislocation 60-62 hr post HCG injection to collect 8-cell embryos (26).

240 embryos were divided to three groups. Group 1; 8-cell embryos were cultured in HTF (Geneocellideal, Iran) medium supplemented with 10% human serum albumin (HSA, Biotest, Germany) until 72 hr. Group 2; 8-cell embryos were vitrified- warmed at 8-cell stage, then were cultured until 72 hr. Group 3; initially 8-cell embryos were vitrified- warmed. After 6-8 hr, alive embryos re-vitrified at compaction stages. Then, compacted embryos were cultured until 64 hr.

### Vitrification procedure

Mouse embryos were vitrified by a two-step procedure with vitrification kit (Geneocellideal, Iran) using the cryolock (Biotech, USA) as a carrier, as described by Kuwayama (27). Embryos were initially equilibrated in equilibration solution (ES) at room temperature for 5-15 min. ES included 7.5% (v/v) ethylene glycol (EG, Sigma Germany) and 7.5% (v/v) dimethylsulfoxide (DMSO, Sigma Germany) dissolved in HTF medium supplemented with 10% HSA.

After initial shrinkage, the embryos regained their original volume and were transferred to vitrification solution (VS) for 1 min. Vitrification solution consisting of 15% (v/v) EG, 15% (v/v) DMSO and 0.5 mol/L sucrose (Sigma Germany) dissolved in HTF medium supplemented with 10% HAS. Within less than 60 sec, 2-3 embryos in minimal VS (<1.0  $\mu$ I) were placed on to the inner surface of the cryolock carrier. The cryolock was plunged vertically into liquid nitrogen.

### Warming technique

After cryo-storage, the embryos were warmed using a three steps dilution procedure with vitrification kit (Geneocellideal, Iran). Briefly, the cryolock containing the embryos were removed from liquid nitrogen and dipped solution (Geneocellideal, into T1 Iran) containing 1.0 mol/L sucrose at 37°C. After 1 min equilibration in T1 solution, the embryos were moved into T2 solution containing 0.5 M sucrose for 3 min. Then the embryos were moved into T3 solution containing 0.25 M sucrose for 3 min. Finally the embryos were washed by HTF medium. Re-vitrification procedure was done by a two-step procedure as described by Kuwayama.

### Assessment of embryo survival

Viability of warmed embryos was determined based on visual examination of the integrity of embryo membrane, zona pellucida and the normality of the cytoplasm two hours after warming. During in vitro culture, embryo development was evaluated every day.

# Molecular assessment by quantitative polymerase chain reaction (qPCR)

Total RNA was isolated from the embryos in each aroup usina QIAzol (Qiagen, Germany) according to the manufacturer's recommendations. To eliminate any genomic DNA contamination, RNA samples were treated with DNase using a kit (EN0521; Fermentas). The RNA concentration was determined by spectrophotometery, and the RNA samples were stored at -80°C until use. The cDNA was synthesized in a total volume of 20 µl containing 5 µg total RNA either with reverse transcriptase (+RT cDNA) or without the enzyme (-RTcontrol) using the cDNA kit (Fermentas, EU) and stored at -20°C until use (26).

All experiments were carried out in triplicate. For PCR reactions, primers were designed by the NCBI website and gene runner software and synthesized by Cinnagen (Iran) (Table I). PCRs were performed using Master Mix and SYBR Green in an Applied Biosystems, StepOne™ thermal cycler (Applied PCR Biosystems, USA). The program started with an initial melting cycle for 5 minutes at 95°C to activate the polymerase, followed by 40 cycles of melting (15 sec at 95°C), annealing (30 sec at 58°C) and extension (15 sec at 72°C). The quality of the PCR reactions was confirmed by melting curve analyses.

Efficiency was determined for each gene using a standard curve (logarithmic dilution series). For each sample, Glyceraldehyde 3phosphate dehydrogenase (GAPDH) was used as the reference gene and target gene were amplified in the same run. Reference gene was approximately equal. The target genes were normalized to a reference gene and expressed relative to a calibrator (Group 1 as control group).

### **Statistical analysis**

Developmental rates of vitrified-warmed embryos were analysed with the Chi square test. qPCR data were presented as mean±SD and were analysed using One-way analysis of variance (ANOVA) followed by Tukey's post hoc test. p<0.05 were considered statistically significant.

#### Results

# Survival rate finding in experimental groups

There was significant difference between re-vitrified embryos (87.5%) and fresh embryos (100%) regarding survival rate (p=0.03), but survival rates of vitrified and revitrified embryos didn't show any significant difference (Table II).

### Development to expanded blastocyst

Based on the current findings, expanded blastocyst rate didn't show any significant difference between vitrified and re-vitrified groups. However, there was significant difference between re-vitrified embryos (82.9%) and fresh embryos (92.5%) (p=0.03) (Table II).

### Degeneration rates investigation

The results of degeneration rate showed no significant differences between vitrified and revitrified groups. While, there was significant difference between re-vitrified and fresh embryos (p=0.03) (Table II).

### Profile of genes expression

Expression of *Bax* (as pro apoptotic gene) was significantly higher in re-vitrified embryos compare to fresh embryos (p=0.004).

Moreover, *Bcl-2* (as anti-apoptotic gene) expression rate, was lower in re-vitrified embryos compare to fresh embryos (p=0.004). Meanwhile, expression rate of *Bax* and *Bcl-2* was similar between re-vitrified and vitrified groups. The expression rate of *ErbB4*, an implantation gene, was similar between re-vitrified and vitrified groups. Although it was significantly lower (p=0.003) in re-vitrified embryos compared to fresh embryos (Figure 1).

**Table I.** Genes, primers, and sizes of amplification products (bp) for quantification of gene expression by real-time quantitative polymerase chain reaction

Gene	Primer sequence	Length (bp)	Code number	Tm (°C)
GAPDH	F:5´-GACTTCAACAGCAACTCCCAC-3´ R:5´- TCCACCACCCTGTTGCTGTA-3´	125	NM_001289726.1	80
Bax	F:5´- CGGCGAATTGGAGATGAACTG-3´ R:5´-GCAAAGTAGAAGAGGGCAA-3´	161	XM_006540584.1	83.5
Bcl2	F:5´- ACCGTCGTGACTTCGCAGAG -3´ R:5´- GGTGTGCAGATGCCGGTTCA -3´	239	NM_009741.1	84
ErbB4	F:5'-TACGAGCCTGCCCAAGTTC- 3' R:5'- GTGCCGATTCCATCACATCCT- 3'	103	XM_006536907.1	76.5

Table II. Embryonic developmental rate in 3 groups after 72 hr culture

Groups	Survival rate (%)	Blastocyst formation (%)	<b>Degeneration</b> (%)
Control	80 (100)	74 (92.5)	6 (67.5)
Experiment 1	71 (88.8 <sup>a</sup> )	71 (88.8)	9 (11.2)
Experiment 2	70 (87.5 <sup>a</sup> )	70 (82.9 <sup>a</sup> )	10 (17.1 <sup>a</sup> )

1; fresh embryos, 2; vitrified 8 cell embryos, 3; vitrified 8 cell and re-vitrified at compact

 $\alpha$  on error bar indicated valued differs with group 1 (p<0.05). Developmental rates of embryos were analysed with the Chi square.



**Figure 1.** Profile of apoptotic and implanting genes expression. 1; fresh embryos, 2; vitrified 8 cell embryos , 3; vitrified 8 cell embryos and re-vitrified at compact stage.  $\alpha$  indicated valued differs with control (p<0.05).



161 239 125

Figure 2. Expression of Bax, Bcl-2, ErbB4, and GAPDH genes

### Discussion

This research evaluated the impact of revitrification on developmental rate and profile of apoptotic and implanting genes expression in mouse embryos. Result of the current study demonstrated clearly that in vitro development in re-vitrified embryos reduced comparing to fresh embryos. However some investigator reported that re-vitrification did not negatively affect the developmental rate of embryos (12, 25, 28).

The reasons for this difference might be vitrification device. concentration of cryoprotectant, stage of embryos, species, and culturing time between vitrification and revitrification. Severed osmotic stress during vitrification and warming may alter integrity of embryo morphology and cellular actin filament organization (19). The factors which have been known to affect the cryopreservation of embryos and subsequently their developmental rate are: types of cryoprotectants, cooling and warming rates, dilution methods, embryo stage and quality of embryos (29).

Some investigators demonstrated that supplementing the cryopreservation media with an antioxidant reduces oxidative stress and increases the survival rates of embryos (30). Shehan et al added ascorbate to cryopreservation solution as an antioxidant, then re-vitrified mouse embryos at pronuclear stage, 2-cell stage, 8-cell stage and blastocyst stage and reported that re-vitrification of embryos didn't affected the survival rates and degeneration rates of embryos (25). An antioxidant treatment can be useful for development of embryos and cells differentiation (30). It is hypothesized that antioxidant can scavenge free radicals, which could damage DNA and the phospholipid membrane of the embryo (31).

Presence of an antioxidant in cryopreservation solution maybe explanation for the disparity between our results and Shehan's data. Gayar et al reported that in vitro and in vivo developmental rate of mouse blastocysts vitrified once or twice, did not show any significant differences with fresh embryos. Additionally, the viability of the embryos after single or repeated vitrification, was virtually identical. A third time vitrification, however, was not tolerated by the majority of embryos (32). This may be indicative of embryos experiencing some stress during successive vitrification procedures. It was reported that repeated freezing mouse embryos without a culture time intervention between each cycle after thawing contained lower mean cell numbers at hatching compared to unfrozen embryos (27).

It seems that repeated exposure of embryos to high cryoprotectant concentration could have a toxic effect on development of embryos. High concentration of cryoprotectants could alter enzymatic activity and plasma membrane integrity of embryos

(33-35). Additionally, disturbance in sodiumpotassium pump during blastocyst formation cause of cellularity may be а and developmental rates reduction after vitrification and re-vitrification (36). Also, the exposure of blastocysts to cryoprotectant decreases the survival rates and increases the DNA-fragmented nuclei in embryos (37). mechanisms Fundamental of oocvtes degeneration and embryo fragmentation is apoptosis (20).

Apoptosis is a regulated program that initiate cell death. Memebers of *Bcl-2* genes family play an important role in apoptosis regulation. Hence, we evaluated the possible impact of re-vitrification on expression of *Bax* and *Bcl-2*. The current findings demonstrate clearly that re vitrification down regulated *Bax* and up regulated *Bcl-2*. *Bcl-2* is anti-apoptotic and is responsible for cell survival, whereas *Bax* is pro-apoptotic and induced cell death (20).

Therefore, the decreased developmental capability of re-vitrified embryos, has a relation to regulation of apoptosis pathways. Similar to the current study, Dhali el al reported that lower survival and development of vitrified zygotes and 2-cell embryos may be attributed to the occurrence and regulation of embryonic apoptosis. The results of Dalhi's studied, clearly indicated a strong relationship between the compromised developmental transcriptional competence and altered activities of Bax, Bcl-2, and p53 genes in the treated embryos (17).

We also evaluated the possible impact of re-vitrification on implanting related gene *ErbB4*. It was demonstrated that expression of *ErbB4* remained similar in re-vitrified embryos compared to vitrified embryos, however, revitrification down regulated *ErbB4* gene expression. Interaction between ErbB4 and HB-EGF mediated the initial attachment of blastocyst to uterine luminal epithelium (22, 38). Signalling by HB-EGF back to the embryo, in turn, activates the program of trophoblast differentiation required for adhesive functions during subsequent attachment and invasion (22).

Successful implantation is an absolute necessity for the reproduction of species. The process by which a foreign blastocyst is accepted by the maternal endometrium is complex and requires interplay of many systems. Many genes, of embryonic and endometrial origin are involved in this process (38-40). Therefore, changes in the gene expression being involved in this process should be further evaluated. Successful live re-vitrified birth from pronucleus and blastocyst have been reported after re-slow freezing embryos; therefore, progress has already been made in this aspect of clinical treatment (11, 23).

Consequently, the re-freezing of blastocysts has resulted in a successful birth (23). In conclusion the current findings clearly demonstrate that re-vitrified and vitrified embryos are similar in developmental rate and expression of Bax, Bcl2 and ErbB4 genes. However, blastocyst rate and the expression Bcl2 and ErbB4 genes differ of Bax. significantly between re-vitrified and fresh embryos. The impacts of the current findings should be further evaluated on obstetrical outcome and long term follow up of the children being born after re-vitrification.

### Conclusion

Re-vitrification can alter the expression of *Bax, Bcl-2* and *ErbB4* genes and developmental rate of mouse embryos in compaction stage.

### Acknowledgements

We would like to thank the office of applied research of Tarbiat Modares University for its support of the project.

### **Conflicts of interest**

The authors declared no conflict of interest.

#### References

- 1. Liebermann J, Dietl J, Vanderzwalmen P, Tucker MJ. Recent developments in human oocyte, embryo and blastocyst vitrification: where are we now? Reprod Biomed Online 2003; 7: 623-633.
- 2. Vajta G, Nagy ZP. Are programmable freezers still needed in the embryo laboratory? Review on vitrification. *Reprod Biomed Online* 2006; 12: 779-796.
- Wennerholm U-B, Söderström-Anttila V, Bergh C, Aittomäki K, Hazekamp J, Nygren K-G, et al. Children born after cryopreservation of embryos or oocytes: a systematic review of outcome data. *Hum Reprod* 2009; 24: 2158-2172.
- Bhattacharya S, Kamath MS. Reducing multiple births in assisted reproduction technology. *Best Pract Res Clin Obstet Gynaecol* 2014; 28: 191-199.
- Belva F ,Henriet S, Van den Abbeel E, Camus M, Devroey P, Van der Elst J, et al. Neonatal outcome of 937 children born after transfer of cryopreserved embryos obtained by ICSI and IVF and comparison with outcome data of fresh ICSI and IVF cycles. *Hum Reprod* 2008; 23: 2227-2238.
- Pelkonen S, Koivunen R, Gissler M, Nuojua-Huttunen S, Suikkari A-M, Hyden-Granskog C, et al. Perinatal outcome of children born after frozen and fresh embryo transfer: the Finnish cohort study 1995-2006. *Hum Reprod* 2010; 25: 914-923.
- Roque M, Lattes K, Serra S, Solà I, Geber S, Carreras R, et al. Fresh embryo transfer versus frozen embryo transfer in in vitro fertilization cycles: a systematic review and meta-analysis. *Fertil Steril* 2013; 99: 156-162.
- Bedoschi G, Oktay K. Current approach to fertility preservation by embryo cryopreservation. *Fertil Steril* 2013; 99: 1496-1502.
- Elnahas A, Alcolak E, Marar EA, Elnahas T, Elnahas K, Palapelas V, et al. Vitrification of human oocytes and different development stages of embryos: An overview. *Middle East Fertil Soc J* 2010; 15: 2-9.
- 10. Matsumoto H, Jiang J, Tanaka T, Sasada H, Sato E. Vitrification of large quantities of immature bovine oocytes using nylon mesh. *Cryobiology* 2001; 42: 139-144.
- Farhat M, Zentner B-S, Lossos F, Bdolah Y, Holtzer H, Hurwitz A. Successful pregnancy following replacement of embryos previously refrozen at blastocyst stage: case report. *Hum Reprod* 2001; 16: 337-339.
- 12. Ito J, Kuramochi M, Inoue A, Yabe K, Fujiwara K, Sonoki S, et al. Cryotop facilitates high developmental ability of re-vitrified mouse embryos. J Reprod Engineer 2010; 13: 21-26.
- Koutlaki N, Schoepper B, Maroulis G, Diedrich K, Al-Hasani S. Human oocyte cryopreservation: past, present and future. *Reprod Biomed Online* 2006; 13: 427-436.
- 14. El-Toukhy T, Khalaf Y, Al-Darazi K, Andritsos V, Taylor A, Braude P. Effect of blastomere loss on the outcome of frozen embryo replacement cycles. *Fertil Steril* 2003; 79: 1106-1111.
- 15. Inaba Y, Miyashita S, Somfai T, Geshi M, Matoba S, Dochi O, et al. Cryopreservation method affects DNA fragmentation in trophectoderm and the speed of re-

expansion in bovine blastocysts. *Cryobiology* 2016; 72: 86-92.

- 16. Park SY, Kim EY, Cui XS, Tae JC, Lee WD, Kim NH, et al. Increase in DNA fragmentation and apoptosisrelated gene expression in frozen-thawed bovine blastocysts. *Zygote* 2006; 14: 125-131.
- Dhali A, Anchamparuthy V, Butler S, Pearson R, Mullarky I, Gwazdauskas F. Gene expression and development of mouse zygotes following droplet vitrification. *Theriogenology* 2007; 68: 1292-1298.
- Anchamparuthy V, Dhali A, Butler S, Pearson R, Jiang H, Gwazdauskas F. Effect of Vitrification of Zygotes in Nylon Mesh on the Differential Expression of Apoptotic Genes in Mouse Embryos. *World J Dairy Food Sci* 2007; 2: 54-61.
- Dhali A, Anchamparuthy V, Butler S, Pearson R, Mullarky I, Gwazdauskas F. Effect of droplet vitrification on development competence, actin cytoskeletal integrity and gene expression in in vitro cultured mouse embryos. *Theriogenology* 2009; 71: 1408-1416.
- 20. Yang MY, Rajamahendran R. Expression of Bcl-2 and Bax proteins in relation to quality of bovine oocytes and embryos produced in vitro. *Anim Reprod Sci* 2002; 70: 159-169.
- 21. Cha J, Sun X, Dey SK. Mechanisms of implantation: strategies for successful pregnancy. *Nature Med* 2012; 18: 1754-1767.
- 22. Davidson LM, Coward K. Molecular mechanisms of membrane interaction at implantation. *Birth Defects Res Part C Embryo Today* 2016; 108: 19-32.
- 23. Estes SJ, Laky DC, Hoover LM, Smith SE, Schinfeld JS, Somkuti SG. Successful pregnancy resulting from cryopreserved pronuclear and cleaved embryos thawed and cultured to blastocysts, refrozen and transferred. A case report. *J Reprod Med* 2003; 48: 46-48.
- 24. Smith LK, Roots EH, Dorsett MJO. Live birth of a normal healthy baby after a frozen embryo transfer with blastocysts that were frozen and thawed twice. *Fertil Steril* 2005; 83: 198-200.
- 25. Sheehan CB, Lane M, Gardner DK. The CryoLoop facilitates re-vitrification of embryos at four successive stages of development without impairing embryo growth. *Hum Reprod* 2006; 21: 2978-2984.
- 26. Shin MR, Choi HW, Kim MK, Lee SH, Lee HS, Lim CK. In vitro development and gene expression of frozen-thawed 8-cell stage mouse embryos following slow freezing or vitrification. *Clin Exp Reprod Med* 2011; 38: 203-209.
- 27. Kuwayama M, Vajta G, Ieda S, Kato O. Comparison of open and closed methods for vitrification of human embryos and the elimination of potential contamination. *Reprod Biomed Online* 2005; 11: 608-614.
- Fathi R, Valojerdi MR, Yazdi PE, Ebrahimi B, Alipour H, Hassani F. Development of 4-cell mouse embryos after re-vitrification. *Cryobiology* 2012; 64: 23-26.
- 29. Rall W. Factors affecting the survival of mouse embryos cryopreserved by vitrification. *Cryobiology* 1987; 24: 387-402.
- 30. Lane M, Maybach JM, Gardner DK. Addition of ascorbate during cryopreservation stimulates subsequent embryo development. *Hum Reprod* 2002; 17: 2686-2693.
- 31. Halliwell B, Gutteridge JM. Free radicals in biology and medicine. Oxford University Press, USA; 2015.

- 32. El-Gayar M, Gauly M, Holtz W. In vitro and in vivo survival of mouse blastocysts after repeated vitrification with the open pulled straw (OPS) method. *Cryoletters* 2010; 31: 454-459.
- 33. Toner M, Cravalho E, Stachecki J, Fitzgerald T, Tompkins R, Yarmush M, et al. Nonequilibrium freezing of one-cell mouse embryos. Membrane integrity and developmental potential. *Biophys J* 1993; 64: 1908-1021.
- 34. Ahn HJ, Sohn IP, Kwon HC, Jo DH, Park YD, Min CK. Characteristics of the cell membrane fluidity, actin fibers, and mitochondrial dysfunctions of frozen-thawed two-cell mouse embryos. *Mol Reprod Dev* 2002; 61: 466-476.
- 35. Lane M, Lyons EA, Bavister BD. Cryopreservation reduces the ability of hamster 2-cell embryos to regulate intracellular pH. *Hum Reprod* 2000; 15: 389-394.
- 36. Kasai M , Ito K, Edashige K. Morphological appearance of the cryopreserved mouse blastocyst

as a tool to identify the type of cryoinjury. *Hum Reprod* 2002; 17: 1863-1874.

- 37. Rajaei F, Karja N, Agung B, Wongsrikeao P, Taniguchi M, Murakami M, et al. Analysis of DNA fragmentation of porcine embryos exposed to cryoprotectants. *Reprod Domest Anim* 2005; 40: 429-432.
- Paria B, Das S, Andrews G, Dey S. Expression of the epidermal growth factor receptor gene is regulated in mouse blastocysts during delayed implantation. *Proc Nat Acad Sci* 1993; 90: 55-59.
- 39. Valles CS, Domínguez F. Embryo-endometrial interaction. *Chang Gung Med J* 2006; 29: 9-14.
- 40. Kaneko Y, Lecce L, Day ML, Murphy CR. β1 and β3 integrins disassemble from basal focal adhesions and β3 integrin is later localised to the apical plasma membrane of rat uterine luminal epithelial cells at the time of implantation. *Reprod Fertil Dev* 2011; 23: 481-495.