# Acute and chronic effects of gold nanoparticles on sperm parameters and chromatin structure in Mice

Mahsa Nazari<sup>1</sup> M.Sc., Ali Reza Talebi<sup>2</sup> Ph.D., Mohammad Hosseini Sharifabad<sup>3</sup> Ph.D., Abolghasem Abbasi<sup>3</sup> M.Sc., Arezoo Khoradmehr<sup>2</sup> M.Sc., Amir Hossein Danafar<sup>2</sup> M.Sc.

- 1. International Campus, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- 2. Research and Clinical Center for Infertility, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.
- 3. Department of Biology and Anatomical Sciences, Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

#### **Corresponding Author:**

*Ali Reza Talebi*, Research and Clinical Center for Infertility, Bouali Ave., Safaeyeh, Yazd, Iran. **Tel:** 09131532174 **Email:** prof\_talebi@ssu.ac.ir

Received: 27 December 2015 Revised: 8 June 2016 Accepted: 9 July 2016

#### Abstract

**Background:** The particles in the range of 1-100 nm are called nanoparticles. Gold nanoparticle is one of the most important metal nanoparticles with wide usage.

**Objective:** This study investigated the effects of gold nanoparticles on sperm parameters and chromatin structure in mice.

**Materials and Methods:** In this experimental study, 72 male bulb-c mice were divided into 9 groups including: 4 Sham groups (Sc 1-4), 4 experimental groups (Au 1-4), and 1 control group (C). Experimental groups received 40 and 200  $\mu$ g/kg/day soluble gold (Au) nano-particles for 7 and 35 days, by intra peritoneal injection, respectively. Sham groups were treated with 1.2 mM sodium citrate solution with 40 and 200  $\mu$ g/kg/day doses for same days and control group did not receive any materials. Motility and Morphology of spermatozoa were analyzed. Chromatin quality was also evaluated using AB (Aniline blue), TB (Toluidine blue) and CMA3 (Chromomycin A3) staining methods.

**Results:** The sperm analysis results showed that motility and morphology of sperm in experimental groups (especially in groups that have been treated for 35 days with nano-particles) had significant decrease in comparison with control group. TB, AB and CMA3 results showed a significant increase in abnormal spermatozoa from all Au-treated groups.

**Conclusion:** Gold nano-particles firstly can reduce the sperm parameters such as motility and normal morphology and secondly affect sperm chromatin remodeling and cause the increase instability of chromatin and also increase the rate of sperm DNA damage. These deleterious effects were more obvious in maximum dose and chronic phase.

Key words: Nanoparticles, Sperm, Chromatin, Mouse. This article extracted from M.Sc. thesis. (Mahsa Nazari)

#### Introduction

ano is a Greek prefix meaning 10<sup>-9</sup> and nowadays this word is used in Nanotechnology (1). Nanotechnology as an emerging technology has been expected to influence the areas of Information technology, medical treatment and environment protection (2). Unique characteristic of nanoparticles in comparison with their raw material is the fact that their diameters are less than 100 nm and cause high ratio of surface to volume (3). Gold is an expensive metal which its major application is the other on jewelry. On hand, the researchers have found the medical applications of gold salts like in dentistry and treatment of rheumatoid arthritis (4-5).

The nanoparticles are capable of penetrating inside the cells and combining with DNA molecules due to their extremely small sizes and unique physical and chemical properties which are different from those of larger materials made up of the same components. Furthermore, the germ cells may interfere with fertilization process and formation of fetus (6). In some experimental studies, the ability of gold nanoparticles with different sizes in penetration into the DNA molecule and positioning in DNA large groove has been shown (7).

It is shown that these nanoparticles have spermatotoxic effects. They suppress the movement of sperm cells and prevent the decondensation of sperm chromatin (8). It has been also shown that low levels of chromosome mutations induced by gold

nanoparticles in initial spermatocytes cells do not result from direct impact of these nanoparticles on genetic structure of masculine gems cells, but rather from inducing their impacts by creating disorders in function of the enzymes which are in charge of repairing spontaneous chromosome (9). damages During transition of spermatozoa, they undergo vast changes which prepare them for fertilization (10). However, sperm chromatin and DNA are very sensitive to exogenous and endogenous stresses during the above mentioned journey.

Unlike somatic cell chromatin that contains histones, the spermatozoa have an specific protein called protamine. During spermatogenesis, about 85% of histones in human and 95% in mouse are replaced by protamines and the reason of remaining of the rest histones is probably because of the role of this genes in fertilization, early embryo development and expression of a number of genes, including influencing genes (11).

Studies have indicated that in the cases of fertilization of ovum by a DNA-damaged sperm. the rates of implantation and pregnancy are significantly reduced (12-14). Although, damaged paternal DNA can be repaired during growth of embryo, but in cases of large amounts of sperm DNA damages, the repair is not enough and the process of reproduction will be affected and finally these damages may cause after birth defects (15). Causes of sperm DNA damage such as causes of male infertility are complex and related to intrinsic and extrinsic factors (12, 16-18). Zakhidov reported that gold nanoparticles cause abnormal chromatin condensation in mouse sperm (6). According to another study selenium nanoparticles can impact the sperm DNA integrity (19).

In spite of widespread application of nanoparticles. there insufficient are information on their impact on human health and environment. Hence, evaluating the probable adverse side effects of gold nanoparticles and their impacts on various sperm parameters as well as chromatin and DNA integrity in men who are exposed to gold nanoparticles are very important. So, the purpose of present study was to investigate the acute and chronic effects of gold nanoparticles with different doses on sperm parameters and chromatin structure of mouse as an experimental model.

#### Materials and methods

#### Nanoparticles

Spherical shape gold nano-particles with a diameter of 10-30 nm were purchased from Tehran University in a form of prepared Colloidal solution with black color and concentration of 1000 ppm per 100 ml for injection to the mice.

#### Animals and experimental design

In this experimental study which was done in research and clinical center for infertility, Yazd, Iran during 2015-2016, totally 72 male bulb-c mice with age of 5 wks and mean weight 22±4 gr were randomly divided into 9 groups (n=8 in each group) including 4 sham, 4 experimental and 1 control groups (C). The sham animals treated with sodium citrate solution of 1.2 mM in doses of 40 and 200 µg/kg/day for 7 and 35 days (Sc 1-4 (20). The treated groups respectively) received gold nanoparticles solution with the same doses and same times (Au 1-4 respectively) by i.p injection and the control group did not receive any material during experiments (20). The mice were kept for at least 2 wks before experiments at standard optical (12 hr light/dark) and appropriate thermal condition (22-25°C) in the clean cages and feed by special food and access to water ad libitum.

#### Epididymal sperm sampling

One day after the last injection in each group, the mice were sacrificed and the cauda epididymis of each animal was removed and placed in a petri dish containing 1000 µl of Ham's F10 medium. The dishes were incubated for 30 min at 37°C and 5% CO<sup>2</sup> (21). Sperm motility and morphology were analyzed and chromatin quality was assessed using aniline blue (AB), toluidine blue (TB) and chromomycin A3 (CMA3) staining methods.

#### Sperm analysis

For sperm motility we used the standard method of Makler chamber (Sefi Medical Co., Haifa) and evaluated by optical microscope (Olympus, Tokyo, Japan) at 20x magnification. The spermatozoa were divided into immotile (grade d), progressive motile (grades a+b) and non-progressive sperm (grade c). To investigate sperm morphology,

we used Papanicolaou staining. Briefly, the sperm smears were fixed in ethanol-ether solution (1:1) for 4 min and then the slides were stained according to the WHO guideline (22).

# Assessment of chromatin quality Aniline blue staining

Histone protein has many lysine residues which react with acidic stains like AB and become blue in color. Spermatozoa with residual histones are considered as AB<sup>+</sup> or immature cells. To do this test, after air-drying of smears, the fixation was done by 3% of glutaraldehyde in phosphate buffer for 30 min. In the next step, samples were stained by 5% solution of AB (Merck, Germany) in acetic acid 4% with the pH of 3.5 for 5 min. After washing by distilled water and mounting by DPX (Merck, Germany), the slides were examined by light microscope (Olympus, Tokyo, Japan) at 100x magnification. The percentage of colorless sperm cells (mature) and blue sperm cells (immature) were obtained (23).

#### Toluidine blue staining

TB is a metachromatic dye binds to the exposed phosphate groups of DNA and shows both chromatin condensation and DNA integrity (21). To do this test, the smears were air-dried and then fixed by ethanol-acetone (1:1) at 4°C for 30 min. For each sample, acidic hydrolysis was done by HCI solution (0.1 N) at 4°C for 5 min, and then washed by distilled water 3 times for 2 min. The staining was done by 0.05% TB in 50% Mcilvian buffer for 10 min and then evaluated by light microscopy at 100x magnification. In this staining we have a range of colors: light blue (sperm with normal chromatins), dark blue (sperm with slightly abnormal chromatin) and (sperm with sever chromatin purple abnormality) (23).

#### Chromomycin A3 staining

CMA3 is used for the evaluation of degree of protamination in spermatozoa (24). After drying of smears, we fixed slides by Carnoy's solution (methanol and Glacial acetic acid, 1: 3) for 10 min at 4°C and then we stained them with (0.25 mg/ ml) in McIlvain buffer (Sigma, USA) for 20 min. After washing and mounting of slides, spermatozoa were counted under florescent microscopy (BX51, Olympus, Tokyo, Japan) with a 475-nm filter and X100 eyepiece magnification and the results were expressed as percentage of CMA3+ spermatozoa (24).

#### Statistical analysis

After collecting data, we used the Spss software version 18. Differences between variables with normal distribution were analyzed by ANOVA test and between each 2 groups were determined by post hoc tests. The term 'statistically significant' was used for P-value  $\leq 0.05$ .

#### Results

#### Sperm parameters

The progressive motility and nonprogressive motility showed significant differences between all groups (p=0.000 and p=0.002 respectively). The treatment of mice with gold nanoparticles causes a significant increase in abnormal sperm morphology (p=0.000). In microscopic assessments, the different forms of sperm morphological defects such as deformed heads, bending and fracture of the necks and coiled tails especially in groups treated with 40 and 200 nanoparticles µg/kg/day of gold were observed. It should be noted that the duration and dose of injection have effect on sperm parameters. In other words these effects were more obvious in maximum dose (200 µg/kg/day) and chronic phase (35 days). The results of sperm parameters analysis of 9 groups are listed in table I.

#### Chromatin quality

As shown in table II, the rates of immature spermatozoa with abnormal chromatin (AB<sup>+</sup>) in 200 Au/7D and 200 Au/35D groups were significantly higher than sham and control groups with p<0.05 (Figure 1). The results of TB staining also showed a significant increase in abnormal spermatozoa in Au-treated animals (p=0.003). There were significant differences between 40 Au/7D group and control/Sc, 40, 35 (91.38±6.88, 98.00±1.41, 98.25±2.76, respectively; p<0.05) (Figure 2). In CMA3 test, the nuclei of spermatozoa with no brightness (CMA3) and spermatozoa with bright nuclei (CMA3<sup>+</sup>) were counted by fluorescent microscope. The rates of CMA3<sup>-</sup> sperm cells were difference between all experimental and sham/control groups. In this test the difference was significant between Au, 200, 35 with control, sham Sc, 40, 35 and sham Sc, 200, 35 (97.13±1.45, 99.75±0.46,  $100.00 \pm 0.00$ , 99.63±0.51, respectively, p<0.05) (Table II) (Figure 3).

Table I. Mean and standard deviation of sperm motility and morphology in groups

	Control	40Sc/7D	200Sc/7D (Sa 2)	40Sc/35D	200Sc/35D	40Au/7D	200Au/7D	40Au/35D	200Au/35D	p-value
Total motility	30.81±5.49	38.63±5.83	(SC 2) 21.49±6.70	27.62±7.41	23.75±7.87	(Au I) 22.12±6.65	(Au 2) 26.13±10.75	(Au 3) 18.37±6.75	(Au 4) 15.94±6.39	0.000
Normal morphology	65.63±5.63	64.88±7.68	69.00±4.34	51.25±9.42	57.63±8.36	45.50±9.59	50.88±8.54	25.63±13.00	29.63±6.63	0.000

P<0.05 represents a significant difference between groups. All data are present as mean ±SD. D: treatment days Sc: sham group Au: experimental group

Table II. Characteristics of sperm nuclear integrity in groups

		1	<u> </u>							
	Control	40Sc/7D	200Sc/7D	40Sc/35D	200Sc/35D	40Au/7D	200Au/7D	40Au/35	200Au/35	n-vəlue
	(C)	(sc 1)	(Sc 2)	(Sc 3)	(Sc 4)	(Au 1)	(Au 2)	(Au 3)	(Au 4)	p-value
AB	98.13±1.45	94.75±3.32	96.25±2.55	97.50±1.41	96.75±1.58	94.63±2.20	98.13±1.64	95.00±2.72	93.00±2.13	0.000
TB	98.25±2.76	92.38±3.85	94.25±2.31	98.00±1.41	96.88±3.22	91.38±6.88	96.00±5.01	96.50±2.00	95.00±2.00	0.003
CMA3 <sup>-</sup>	99.75±0.46	$97.50 \pm 1.51$	$97.88 \pm 2.90$	$100.00 \pm 0.00$	99.63±0.51	$97.38 \pm 2.20$	99.75±0.46	$98.50 \pm 1.06$	97.13±1.45	0.000
		41.00				6 m				

P<0.05 represents a significant difference between groups. All data are present as mean  $\pm$ SD.

D: treatment days Sc: sham group Au: experimental group



Figure 1. Aniline blue staining in 200Au/35 group. Normal spermatozoa are seen in light blue and abnormal spermatozoa are seen in dark blue  $\times 100$ .



**Figure 2.** Toluidine blue staining in 200Au/35 group.  $TB^+$  spermatozoa are dark blue and  $TB^-$  spermatozoa are seen in light blue ×100.



Figure 3. CMA3 staining in 200Au/35 group. Abnormal sperm CMA3<sup>+</sup> is seen in bright green and normal sperm CMA3<sup>-</sup> is seen in non-bright green  $\times 100$ .

#### Discussion

According to our results, the gold nanoparticles firstly can reduce the sperm parameters such as motility and normal morphology and secondly impair sperm chromatin remodeling leading to increase instability of chromatin. It also increased the rate of sperm DNA damage. Regard to the sperm parameters, we showed that the percentage of sperm with progressive motility decreased especially at the dose of 200 µg/kg/day for 35 days.

Because the duration of mouse spermatogenesis is about 31 days and to ensure that the nanoparticles effect on all of the cells in the process of spermatogenesis, administrated them for 5 we weeks. Furthermore, the impact of these nanoparticles on previously produced epididymal sperm cells was evaluated due to formation of disulfide bonds between adjacent protamines in chromatin structure of the sperm which takes place after one week in epididymis. It was observed that, epididymal spermatozoa are also influenced by gold nanoparticles.

The first part of our results showed that gold nanoparticles can affect different sperm morphological and motility parameters. In fact, treated groups showed more percentage of spermatozoa with abnormal morphology and low motility in comparison with sham and control groups. Taking together, the highest reduction in sperm motility and normal morphology was seen at dose of 200 g/kg/dayµ and duration of 35 days. It should be noted that the most abnormalities in sperm morphology were observed in tail (the twisted tail) and then in sperm head. Zakhidov et al demonstrated that the gold nano-particle shows spermatotoxic effects and may impact mouse spermatogenesis (6).

In accordance to our results, Wiwanitkit *et al* in a study on donor sperm stated that the gold nano-particles show dose-dependent effects on sperm motility (8). Also, like our results, Moretti *et al* showed that both silver and gold nanoparticles affect the sperm motility in dose-dependent manner (28) .Of

course, it should be noted that the studies indicating the effects of gold nano-particles on male reproductive system are very limited and the majority of researches were done on other metal nano-particles like silver, zinc and selenium. Grodzka-Ostrowska *et al* also showed that although silver nano-particles can reduce the count of spermatozoa, but these particles do not decrease the sperm normal morphology (22).

However, we saw different forms of sperm abnormalities such as head and tail gold-treated malformations in animals. Another study conducted by Rezvanfar et al showed that nano-selenium changes sperm parameters, including morphology, motility, viability and count (19). Also, in another study it was shown that zinc nanoparticles may affect mouse spermatogenesis and cause a significant decrease in sperm motility and increase in abnormal sperm morphology (1).

In recent years, gold nanoparticles have attracted particular attention and have found widespread use in scientific research, industry, medical, and especially to recognize and destroy cancer cells (25). Spermatogenesis is a complex process and is very sensitive that different factors can affect its quality (26). Nano-particles have negative effects on many organs, such as testes (24). Experimental in vivo studies have shown that nano-particles may easily transfer through the blood-brain and blood-testis barriers (28).

Although, about gold nanoparticle, the exact mechanism of toxicity is not clear yet, but, it is demonstrated that this particle can incorporate to the major groove of DNA and cause different abnormalities in cells (29). In etiological terms, it should be mentioned that the gold nanoparticles can bind to DNA molecules and impact on chromosome repairing enzymes. In addition, the deleterious effects of other nanoparticles, such as increasing ROS production, production of mutagenic molecules may also come true about gold nanoparticles (30).

Consequently, more investigations are demanded in this area. In present study we used the cytochemical staining like AB, TB, and CMA3 to evaluate the quality of sperm chromatin. The results of AB, TB, and CMA3 tests showed a significant difference between treated and controls. Since AB indicates the presence of excessive histones, TB indicates the amount of chromatin condensation and CMA3 test indicates the protamine deficiency, it can be concluded that the process of spermiogenesis is affected by dold nanoparticles. In this part, the greatest impact was also observed in a dose of 200 g/kg/day and duration of 35 days. So, regard to the effects of nanoparticle on sperm chromatin, we can say that gold in the form of nanoparticle may cause increase residual histones, DNA fragmentation and protamine deficiency in mouse spermatozoa as an experimental model.

In accordance to our results, Zakhidov et al investigated the effects of gold nanoparticles on mouse sperm chromatin using TB and showed that this metal nano-particle impedes the density of chromatin and cause abnormal chromatin condensation in mouse (6) .In addition to gold nanoparticles, Rezvanfar et al considered the effect of selenium the quality nanoparticles on of sperm chromatin using AB and AO staining methods and showed that these nanoparticles affect the sperm DNA integrity (19). As it was mentioned before, although, the mechanisms of biological effects of gold nanoparticles need to be investigated in future, but the role of free radicals and production of mutagenic following molecules nanoparticles intake should be considered.

#### Conclusion

In conclusion, our results showed that gold nanoparticles can change the mouse sperm parameters and chromatin structure and these deleterious effects are more obvious in maximum dose and chronic phase. Finally, due to the action of gold on sperm fertility potential, the use of golden jewelries and other forms of gold is not recommended especially in young men that want to have a baby.

#### Acknowledgements

This thesis was supported by a grant from the International Campus of Shahid Sadoughi University of Medical Sciences, Yazd, Iran.

#### **Conflict of interest**

The authors have no conflicts of interest.

#### References

- 1. Thakkar KN, Mhatre SS, Parikh RY. Biological synthesis of metallic nan- oparticles. *Nanomedicine* 2010; 6: 257-262.
- Komatsu T, Tabata M, Kubo-Irie M, Shimizu T, Suzuki K, Nihei Y, et al. The effects of nanoparticles on mouse testis Leydig cells in vitro. *Toxicol In Vitro* 2008; 22: 1825-1831.
- Talebi AR, Khorsandi L, Moridian M. The effect of zinc oxide nanoparticles on mouse spermatogenesis. *J Assist Reprod Genet* 2013; 30: 1203-1209.
- 4. Taylor A. herapeotic uses of trace elements. *Clin Endocrinol Metab* 1985; 14: 703-724.
- Sigler JW, Gilbert BB, Howard D, John TS, Dwight C, Mccrum WR. Gold Salts in the Treatment of Rheumatoid Arthritis: A Double-Blind Study. *Ann Int Med* 1974; 80: 21-26.
- Zakhidov ST, Marshak TL, Malolina EA, Yu A, Kulibin IA, Zelenina S M, et al. Gold Nanoparticles Disturb Nuclear Chromatin Decondensation in Mouse Sperm in vitro. *Biochem* 2010; 4: 293-296.
- Appasamy M, Muttukrishna S, Pizzey AR, Ozturk O, Groome NP, Serhal P, Jauniaux E, et al. Relationship between male reproductive hormones, sperm DNA damage and markers of oxidative stress in infertility. *Reprod Biomed Online* 2007; 14: 159-165.
- 8. Wiwanitkit V, Sereemaspun A, Rojanathanes R. Effect of Gold Nanoparticles on Spermatozoa. The First World Report. *Fertil Steril* 2009; 91: 7-8.
- Zakhidova ST, Pavlyuchenkovaa SM, Marshak TL, Rudoy VM, Dement'eva OV, Zelenina IA, et al. [Effect of Gold Nanoparticles on Mouse Spermatogenesis]. *Izv Akad Nauk Ser Biol* 2012; 39: 229-236. (In Russian)
- Greco E, Iacobelli M, Rienzi L, Ubaldi F, Ferrero S, Tesarik J. Reduction of the incidence of sperm DNA fragmentation by oral antioxidant treatment. *J Androl* 2005; 26: 349-353.
- Tarozzi N, Bizzaro D, Flamigni C, Borini A. Clinical relevance of sperm DNA damage in assisted reproduction. *Reprod Biomed Online* 2007; 14: 746-757.
- 12. Morris ID, Ilott S, Dixon L, Brison DR. The spectrum of DNA damage in human sperm assessed by single cell gel electrophoresis (Comet assay) and its relationship to fertilization and embryo development. *Hum Reprod* 2002; 17: 990-998.
- Tavalaee M, Razavi S, Nasr-Esfahani MH. [Effects of spermiogenesis defecys on Fertilization and pregnancy rate in IVF patients]. Yakhteh 2007; 9: 103-101. (In Persian).
- 14. Razavi S, Nasr-Esfahani MH, Tavalaee M, Ameri A. [Assessing the sperm DNA damage on ART outcome]. *J Arak Univ of Med Sci* 2008; 11: 21-33. (In Persian)
- 15. Shayesteh M, Nasr-Esfahani MH, Razavi S, Nazem HA, Deemeh MR, Tavalaee M. The efficiency of Zeta method in separation of sperm with normal morphology and chromatin structure. *Shahrekord Univ Med Sci J* 2008; 10: 20-27.

- 16. Nasr-Esfahani MH, Tavalaee M, Deemeh MR. Origins and Evaluation of DNA Damage in Infertile Individual. *J Iran Anat Sci* 2008; 24: 489-500.
- Tavalaee M, Razavi R, Nasr-Esfahani MH. Influence of sperm chromatin anomalies on assisted reproductive technology outcome. *Fertil Steril* 2009; 91: 1119-1126.
- Mardani M, Nasr-Esfahani MH, Razavi S, Shirazi R, Tavalaee M. [Differentiation between the effect of protamine deficiency and failed oocyte activation on fertilization post ICSI]. *J Iran Anat Sci* 2006; 4: 95-103. (In Persian)
- 19. Rezvanfar M A, Shahverdi AR, Ahmadi A, Baeeri M, Mohammadirad A, Abdollahi M, et al. Protection of cisplatin-induced spermatotoxicity, DNA damage and chromatin abnormality by selenium nano-particles. *Toxicol Appl Pharmacol* 2013; 266: 356-365.
- 20. Lasagna-Reeves C, Gonzalez-Romero D, Barria MA, Olmedo I, Clos A, Sadagopa Ramanujam VM, et al. Bioaccumulation and toxicity of gold nanoparticles after repeated administration in mice. *Biochem Biophys Res Commun* 2010; 393: 649-455.
- Talebi AR, Moein MR, Tabibnejad N, Ghasemzadeh J. Effect of varicocele on chromatin condensation and DNA integrity of ejaculated spermatozoa using cytochemical tests. *Andrologia* 2008; 40: 245-251.
- 22. Gromadzka-Ostrowska J, Dziendzikowska K, Lankoff A, Dobrzyn ska M, Instanes C, Brunborg G. Silver nanoparticles effects on epididymal sperm in rats. *Toxicol Lett* 2012; 214: 251-258.
- 23. Talebi AR, Khalili MA, Hossaini A. Assessment of nuclear DNA integrity of epididymal spermatozoa following experimental chronic spinal cord injury in the rat. *Int J Androl* 2007; 30: 163-169.
- 24. Iranpour FG, Nasr-Esfahani MH, Valojerdi MR, Al-Taraihi TM. Chromomycin A3 staining as a useful tool for evaluation of male fertility. *J Assist Reprod Genet* 2000; 17: 60-66.
- 25. Reed JA, Cook A, Halaas DJ, Parazzoli P, Robinson A, Matula TJ, Grieser F. The Effects of Microgravity on Nanoparticle Size Distributions Generated by the Ultrasonic Reduction of an Aqueous Gold-Chloride Solution. *Ultrasonics Sonochem* 2003; 10: 285-289.
- 26. Pryor JL, Hughes C, Foster W, Hales BF, Robaire B. Critical windows of exposure for children's health: The reproductive system in animals and humans. *Environ Health Perspect* 2000; 108: 491-503.
- 27. Li C, Taneda S, Taya K, Watanabe G, Li X, Fujitani Y, Nakajima T, Suzuki AK. Effect of in utero exposure to nanoparticle-rich diesel exhaust on testicular function in immature male rats. *Toxicol Lett* 2009; 185: 1-8.
- Moretti E, TerzuoliG, Renieri T, Iacoponi F, Castellini C, Giordano C, CollodeG.In vitro effect of gold and silver nanoparticles on human spermatozoa. *Andrologia* 2013; 45: 392-396.
- 29. Skuridin SG, Dubinskaya VA, Rudoi VM. Effect of Gold Nanoparticles on DNA Package in Model Systems. *Dokl. Biochem Biophys* 2010; 432: 141-143.
- Pawar K, Kaul G. Toxicity of titanium oxide nanoparticles causes functionality and DNA damage in buffalo (Bubalusbubalis) sperm in vitro. *Toxicol Ind Health* 2012; 30: 520-533.