



Research Article

Curcumin improves growth factors expression of bovine cumulus-oocyte complexes cultured in peritoneal fluid of women with endometriosis

Hendy Hendarto¹ Ph.D., Muhammad Yohanes Ardianta Widyanugraha¹ M.D.,
Widjiati Widjiati² Ph.D.

¹Department of Obstetrics and Gynecology, Faculty of Medicine, Airlangga University/Dr Soetomo Hospital Surabaya, Indonesia.

²Laboratory of Embriology, Faculty of Veterinary Medicine, Airlangga University, Surabaya, Indonesia.

Abstract

Background: Peritoneal fluid (PF) from infertile women with endometriosis contains inflammatory mediators that may interfere with folliculogenesis.

Objective: The aim was to evaluate the effects of curcumin on growth factors expression by evaluating Growth Differentiation Factor-9 (GDF-9), Kit Ligand (KitL), and Tumor Necrosis Factor α (TNF α) expressions in bovine cumulus-oocyte complexes (COC)s cultured with PF from infertile women with endometriosis.

Materials and Methods: In this experimental study, 21 infertile women (aged between 20 and 40 years) who referred to Dr Sutomo Hospital from January to July 2015 were enrolled. COCs were aspirated from antral follicles of bovine ovaries. PF was collected from infertile women with endometriosis undergoing laparoscopy for infertility evaluation. Curcumin, a strong anti-inflammatory turmeric, was added in Tissue Culture Medium 199 (TCM199) and PF for culture medium. Bovine COCs were cultured into three groups of the medium: 1. TCM199, 2. TCM199 + PF, and 3. TCM199 + PF + curcumin. GDF-9, KitL, and TNF α expressions were examined using immunohistochemistry technique.

Results: GDF-9 expression of bovine COCs cultured in PF with curcumin addition (2.67 ± 0.98) was found to increase compared to those cultured without curcumin (0.50 ± 0.67) ($p \leq 0.001$). It was similar to KitL expression of bovine COCs cultured with curcumin (2.67 ± 1.23), which increased compared to those without curcumin (0.33 ± 0.49) ($p \leq 0.001$). A significant difference in TNF α expression was noted between groups with or without curcumin ($p \leq 0.001$).

Conclusion: In the culture of PF from infertile women with endometriosis, curcumin addition improves the growth factors expression of bovine COCs. The increase of GDF-9 and KitL expressions will improve folliculogenesis.

Key words: Curcumin, Bovine, GDF9, Kit Ligand, TNF α , Endometriosis.

Corresponding Author:

Hendy Hendarto;

email:

hndhendy@yahoo.com

hendy.hendarto@fk.unair

.co.id

Tel: (+62) 8123007080

Received 8 March 2018

Revised 11 August 2018

Accepted 13 August 2018

Production and Hosting by
Knowledge E

© Hendy Hendarto et al. This article is distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use and redistribution provided that the original author and source are credited.

Editor-in-Chief:

Aflatoonian Abbas M.D.

 OPEN ACCESS

1. Introduction

Endometriosis is one of the most common gynaecological diseases characterized by the growth of endometrial-like tissue outside the uterus that subsequently induces a chronic inflammatory reaction (1, 2). Numerous symptoms are associated with endometriosis, including dysmenorrhea, pelvic pain, dyspareunia, and infertility, as well as reduced quality of life (2–4). The relationship between endometriosis and infertility remains unclear. One of the postulations of decreased fertility in women with endometriosis is the reduction of oocyte quality caused by abnormal folliculogenesis related to peritoneal fluid (PF) inflammation and oxidative stress (5–8).

Folliculogenesis is a dynamic process marked by proliferation and differentiation of granulosa cells and maturation of oocyte. The regulation of ovarian folliculogenesis, determined by a number of growth factors and classic endocrine mechanism, provides an optimal environment to produce fertilizable oocyte (9, 10). There are two important growth factors that contribute to the regulation of ovarian folliculogenesis including the interactions between oocyte and granulosa cells. Those growth factors are Growth Differentiation Factor 9 (GDF-9) produced by oocyte, which is useful for granulosa cell proliferation and differentiation, and Kit Ligand (KitL) secreted by granulosa cells, which induces oocyte maturation (11–14).

In women with endometriosis, ovaries are naturally bathed in PF that is rich in inflammatory mediators. This may cause abnormal folliculogenesis and subsequently result in infertility. The PF of women with endometriosis contains a variety of inflammatory mediators, including Tumor Necrosis Factor α (TNF α), Interleukin-6 (IL-6), and Interleukin-8 (IL-8) (15–17).

Curcumin, which is derived from *Curcuma longa* (turmeric), has a strong potential anti-inflammatory activity. It has been widely used both traditionally

and scientifically to treat various conditions including inflammatory diseases such as rheumatoid arthritis, chronic anterior uveitis, and ulcerative colitis through various intracellular and extracellular molecular pathways (18–20). Treatment with curcumin can reduce implant size and cell proliferation in endometriosis model rat (21), but the effects of curcumin on bovine ovarian growth factors in endometriosis and infertility remain controversial.

The objective of our study was to evaluate whether curcumin could improve growth factors expression in bovine cumulus-oocyte complexes (COC)s by analyzing GDF-9, KitL, and TNF α expressions in culture medium with PF from infertile women with endometriosis.

2. Materials and Methods

In this laboratory experimental study, two growth factors expression of folliculogenesis regulation on bovine COCs cultured in three different types of medium was evaluated. Group 1 (control group) Tissue Culture Medium 199 (TCM199) only, Group 2 (endometriosis group) TCM199 plus PF from infertile women with endometriosis, and Group 3 (endometriosis+curcumin) TCM199 plus PF from infertile women with endometriosis added with curcumin. A total of 21 bovine COCs were cultured in the medium; each group contained 7 COCs.

2.1. Endometriosis peritoneal fluid (PF)

PF samples were collected from 21 women aged between 20 and 40 years with endometriosis undergoing laparoscopy for infertility evaluation at Dr Sutomo Hospital from January to July 2015. The diagnosis of endometriosis was made by visual inspection and peritoneal biopsy according to the American Society for Reproductive Medicine

criteria (22, 23). PF has been collected by aspiration from posterior cul-de-sac during the laparoscopic procedure. The samples were placed in a tube and centrifuged at 600 g for 10 min. The supernatants were stored at -80°C until analysis.

2.2. Curcumin

Curcumin has been obtained from Merck Schuchardt OHG (85662 Hohenbrunn, Germany). Curcumin 0.2 ml/medium (curcumin 20 $\mu\text{g}/\text{ml}$ dissolved into Carboxymethyl Cellulose Sodium/Na-CMC) was added and homogenized in 10 ml of TCM199 medium and PF from infertile women with endometriosis (30 μl). BSA (3%) was added until the pH reached to 7.4–7.8. The solution was then filtered through a 0.22- μm microfilter, and 100 mL of solution was placed in a Petri dish for culture.

2.3. Bovine cumulus-oocyte complex

COCs aspirated from antral follicles with a diameter of 3–8 mm were obtained from bovine ovaries in a slaughterhouse. The ovaries were washed and stored in 0.89% NaCl with penicillin-G (1000 IU/ml) and streptomycin sulfate (0.2 $\mu\text{g}/\text{l}$) at a temperature of 30–35 $^{\circ}\text{C}$. Before follicle aspiration, the diameter of the follicles was measured with a caliper. The COCs were aspirated using an 18-G needle connected to a 5-ml syringe containing 1 ml of phosphate buffered saline (PBS) with 3% bovine serum albumin (BSA) and 50 $\mu\text{g}/\text{ml}$ gentamycin.

The COCs were washed three times successively in PBS medium and one time in TCM199, placed in TCM199 medium with 50 mIU/ml FSH and 50 mIU/ml LH, divided into three groups by placing them into medium groups 1, 2, and 3, and then incubated at a temperature of 38 $^{\circ}\text{C}$ in 5% CO_2 for 24 hr. Subsequently, the COCs of each group

were fixated in a glass flask and subjected to immunohistochemical staining for GDF-9 (Bioss Antibodies Inc USA, catalog no. Bs-4720R), KitL (Abcam USA, catalog no. ab52603), and $\text{TNF}\alpha$ (Bioss Antibodies Inc USA, catalog no. Bs-2081R) expressions.

The three expressions were semi-quantitatively assessed according to the modified Remmele method which is the result of multiplication between the percentage score of immunoreactive cells (positive cells) with the color intensity score generated on the cell (24).

2.4. Ethical consideration

The study has been approved by the Ethical Committee for Health Research at Dr Sutomo Hospital Surabaya (428/Panke. KKE/X/2014). Written informed consent was obtained from participants before starting the study.

2.5. Statistical analysis

Data analysis was performed using SPSS statistical software (Statistical Package for the Social Sciences, version 17.0, SPSS Inc, Chicago, Illinois, USA). Normality of variable was tested with the Kolmogorov–Smirnov test. One-way analysis of variance (ANOVA) test was used to detect significant differences of all variables. $p < 0.05$ was accepted as statistically significant.

3. Results

3.1. GDF-9 expression

The GDF-9 expression in bovine COC was determined by the dark color of immunoreactive cells on the immunohistochemical staining result, and it was different among the three groups (Figures 1A, B, C). The mean expression of GDF-9 in bovine

COC cultured in PF from infertile women with endometriosis+curcumin (2.67 ± 0.98) increased compared to those cultured without curcumin (0.50 ± 0.67) but reduced compared to the control (5.83 ± 1.58). ($p \leq 0.001$) (Figure 2).

3.2. Kit ligand expression

The KitL expression in bovine COC was determined by the dark color of immunoreactive cells on the immunohistochemical staining result, and it was different among the three groups (Figures 1D, E, F). The semi-quantitative results of KitL expression in bovine COC were similar to those of GDF-9 expression. The mean expression of KitL in bovine COC cultured in PF from infertile women with endometriosis (0.33 ± 0.49) reduced compared to the control (3.92 ± 2.02) and those in the endometriosis + curcumin group (2.67 ± 1.23) ($p \leq 0.001$). There was also a significant difference between KitL expression in bovine COC cultured

in PF from infertile women with endometriosis + curcumin and control group ($p = 0.03$) (Figure 2).

3.3. TNF α expression

The semi-quantitative results of TNF α expression in bovine COC cultured in control, endometriosis, and endometriosis+curcumin groups were 0.00 ± 0.00 ; 8.67 ± 3.72 ; and 2.17 ± 1.69 , respectively (25). TNF α expression in bovine COC cultured in PF from infertile women with endometriosis increased compared to those in control group, whereas TNF α expression in bovine COC cultured in PF from infertile women with endometriosis added with curcumin reduced compared to those cultured without curcumin; however, the level increased compared to the control ($p \leq 0.001$) (Figure 2). In this study, the relationship of three variables was also examined using regression analysis and revealed a significant association among TNF α , GDF-9, and KitL expression cultured in PF from infertile women with endometriosis added with curcumin ($p < 0.05$).




Figure 1: The immunoreactive cells (black arrow) of GDF-9 (A,B,C) and KitL (D,E,F) on immunohistochemical staining of bovine COC in the following medium: 1. control, 2. endometriosis, and 3. endometriosis + curcumin ($p < 0.05$). GDF9 = Growth Differentiation Factor-9; KitL = Kit Ligand; COC = cumulus oocyte complex. Magnification 400x; Olympus BX-50. Pentax optio 230 Digital Camera 2.0 megapixel.




Figure 2: The expression of GDF-9, KitL, and TNF α on bovine COC in the following medium: 1. control, 2. endometriosis, and 3. endometriosis + curcumin ($p < 0.05$).

4. Discussion

In this study, GDF-9 and KitL expressions in bovine COC cultured with PF from infertile women with endometriosis were significantly reduced compared to control. It suggests that numerous inflammatory mediators in the PF of infertile women with endometriosis enters the bovine COC and further inhibits oocytes and granulosa cells' activity. This results in abnormal folliculogenesis and subsequently infertility.

The PF of infertile women with endometriosis contains a variety of inflammatory mediators (15, 26) and the PF component plays an essential role in the process of folliculogenesis, ovulation, and fertilization. Communication between oocytes and granulosa cells occurs during folliculogenesis to produce mature and fertilizable oocytes. This process is impaired in infertile women with

endometriosis. We previously reported abnormal GDF-9 and KitL expression in endometriosis (7, 27).

GDF-9 is part of the TGF β family member dominantly secreted by oocyte and plays an important role in ovarian folliculogenesis. Study in GDF-9 deficient female mice has resulted in impaired follicular development after primary follicle growth that causes infertility (28). GDF-9 is a determining factor for follicular growth, maturation and survival through inhibition of the apoptosis process in granulosa cell and follicular atresia (29, 30). In this study, the addition of curcumin to the culture medium of PF from infertile women with endometriosis resulted in more improved GDF-9 expression than those without curcumin. It is possible that the anti-inflammatory effects of curcumin release oocyte suppression. By taking into account the fact that the expression of

GDF-9 was still lower than the control, it indicates that it is necessary to adjust the dosage of curcumin. These findings indicate that the dose of curcumin plays a role in GDF-9 expression.

In our study, KitL expression in bovine COC cultured in PF from infertile women with endometriosis added with curcumin significantly increased from those cultured without curcumin. It indicates that curcumin has a positive effect on KitL expression. KitL is a growth factor produced by granulosa cells in the ovarian follicle. KitL binds to its receptor in the oocyte, c-Kit, and actively functions for signal pathways during folliculogenesis (31). The initiation of primordial follicle growth, and the development of oocyte and follicle are regulated by the KitL/c-Kit system (13, 31). Study in the buffalo ovaries by Mahajan prove that KitL could be used to predict the oocyte maturation as well as oocyte competence (32).

The changes on two growth factors expression, GDF-9, and KitL suggest that curcumin has a repair effect on oocyte-granulosa cell interactions and the regulation of ovarian folliculogenesis. In order to evaluate that curcumin has a repair effect on oocyte-granulosa cell interactions and the regulation of ovarian folliculogenesis, we assessed the effect of curcumin on bovine COC inflammation via $TNF\alpha$ expression. An increase in $TNF\alpha$ expression was noted in the PF from infertile women with endometriosis compared to that from normal women. In addition, we found a correlation between $TNF\alpha$ concentration and the degree of endometriosis severity. In fact, it has been reported that $TNF\alpha$ plays a role in the inflammatory process and angiogenesis, triggers follicular atresia, and impairs oocyte maturation (33–35). The findings of increased $TNF\alpha$ expressions in bovine COC cultured in PF from infertile women with endometriosis suggest that $TNF\alpha$ from PF enters the COC and is active. The addition of curcumin to the culture media results in an improvement as indicated by the reduced $TNF\alpha$ expression.

Curcumin exhibits anti-endometriosis activities by affecting MMP2 and TIMP2 (36). It works as anti-inflammatory medication by inhibiting cyclooxygenase2 and lipoxygenase and suppressing the activity of NFkB and the level of inflammatory cytokines, including $TNF\alpha$, IL-6, IL-8, and $IFN-\gamma$ (18, 19, 20, 36). Based on its mechanism of action, curcumin can suppress an inflammatory process in bovine COC cultured in PF from infertile women with endometriosis. Yet, we found that the decrease in $TNF\alpha$ expression did not reach a normal level. We also found a significant correlation among $TNF\alpha$, GDF9, and KitL expression, suggesting that the inflammatory factor of $TNF\alpha$ may play a role in GDF-9 and KitL interaction. Studies evaluating the dose-response effects of curcumin on the GDF-9-KitL interaction are still needed.

The limitation of our study included the use of only one concentration of curcumin. However, this was the first attempt to evaluate the effects of curcumin on oocyte-granulosa cell interaction and the regulation of ovarian folliculogenesis in the presence of PF from infertile women with endometriosis.

5. Conclusion

Our findings suggest that in the culture of PF from infertile women with endometriosis, curcumin addition improved growth factors expression of bovine COCs through the decrease of inflammation. The increase of GDF-9 and KitL expressions due to the anti-inflammatory effect of curcumin will improve oocyte-granulosa cell interaction and subsequently folliculogenesis. Further studies are warranted.

Acknowledgements

The authors thank Prof. Togas Tulandi from McGill University, Canada, for his assistance in the preparation of the manuscript.

Conflict of Interest

The authors report no conflict of interest.

References

- [1] Kennedy S, Bergqvist A, Chapron C, D'Hooghe T, Dunselman G, Greb R, et al. ESHRE guideline for the diagnosis and treatment of endometriosis. *Hum Reprod* 2005; 20: 2698–2704.
- [2] Dunselman GA, Vermeulen N, Becker C, Calhaz-Jorge C, D'Hooghe T, De Bie B, et al. ESHRE guideline: management of women with endometriosis. *Hum Reprod* 2014; 29: 400–412.
- [3] National Institute for Health and Care Excellence. Endometriosis: diagnosis and management. NICE guideline [NG73]; 2017.
- [4] De Graaff AA, D'Hooghe TM, Dunselman GA, Dirksen CD, Hummelshoj L, Werf EndoCost Consortium, et al. The significant effect of endometriosis on physical, mental and social wellbeing: results from an international cross-sectional survey. *Hum Reprod* 2013; 28: 2677–2685.
- [5] Xu B, Guo N, Zhang XM, Shi W, Tong XH, Iqbal F, et al. Oocyte quality is decreased in women with minimal or mild endometriosis. *Sci Rep* 2015; 5: 10779.
- [6] Du YB, Gao MZ, Shi Y, Sun ZG, Wang J. Endocrine and inflammatory factors and endometriosis-associated infertility in assisted reproduction techniques. *Arch Gynecol Obstet* 2013; 287: 123–130.
- [7] Hendarto H. Pathomechanism of infertility in endometriosis. In: Chaudhury K, Chakravarty B. *Endometriosis-Basic Concepts and Current Research Trends*. Rijeka, Croatia: Intech; 2012; 343–54.
- [8] Malvezzi H, Da Broi MG, Meola J, Rosa E-Silva JC, Ferriani RA, Navarro PA. Peritoneal fluid of women with endometriosis reduces SOD1 in bovine oocytes in vitro maturation. *Cell Tissue Res* 2018; 372: 621–628.
- [9] Onagbesan O, Bruggeman V, Decuypere E. Intra-ovarian growth factors regulating ovarian function in avian species: a review. *Anim Reprod Sci* 2009; 111: 121–140.
- [10] Evron A, Blumenfeld Z, Adashi EY, Kol S. The role of growth factors in ovarian function and development. *The Global Library of Women's Medicine*; 2015.
- [11] Thomas FH, Vanderhyden BC. Oocyte-granulosa cell interactions during mouse follicular development: regulation of kit ligand expression and its role in oocyte growth. *Reprod Biol Endocrinol* 2006; 4: 19.
- [12] Senbon S, Hirao Y, Miyano T. Interactions between the oocyte and surrounding somatic cells in follicular development: lessons from in vitro culture. *J Reprod Dev* 2003; 49: 259–269.
- [13] Celestino JJH, Matos MHT, Saraiva MVA, Figueiredo JR. Regulation of ovarian folliculogenesis by Kit Ligand and the c-Kit system in mammals. *Anim Reprod* 2009; 6: 431–439.
- [14] Sanfins A, Rodrigues P, Albertini DF. GDF-9 and BMP-15 direct the follicle symphony. *J Assist Reprod Genet* 2018; 35: 1741–1750.
- [15] Oral E, Olive DL, Arici A. The peritoneal environment in endometriosis. *Hum Reprod Update* 1996; 2: 385–398.
- [16] Wu MH, Hsiao KY, Tsai SJ. Endometriosis and possible inflammation markers. *Gynecol Minimally Invas Ther* 2015; 4: 61–67.
- [17] Izumi G, Koga K, Takamura M, Makabe T, Satake E, Takeuchi A, et al. Involvement of immune cells in the pathogenesis of endometriosis. *J Obstet Gynaecol Res* 2018; 44: 191–198.
- [18] Jurenka JS. Anti-inflammatory properties of curcumin, a major constituent of *Curcuma longa*: a review of preclinical and clinical research. *Altern Med Rev* 2009; 14: 141–153.
- [19] He Y, Yue Y, Zheng X, Zhang K, Chen S, Du Z. Curcumin, inflammation, and chronic diseases: how are they linked? *Molecules* 2015; 20: 9183–9213.
- [20] Arablou T, Kolehdoz-Mohammadi R. Curcumin and endometriosis: Review on potential roles and molecular mechanisms. *Biomed Pharmacother* 2018; 97: 91–97.
- [21] Kizilay G, Uz YH, Seren G, Ulucam E, Yilmaz A, Cukur Z, et al. In vivo effects of curcumin and deferoxamine in experimental endometriosis. *Adv Clin Exp Med* 2017; 26: 207–213.
- [22] American society for reproductive medicine. Revised american society for reproductive medicine classification of endometriosis: 1996. *Fertil Steril* 1997; 67: 817–821.
- [23] Jorgensen H, Hill AS, Beste MT, Kumar MP, Chiswick E, Fedorcsak P, et al. Peritoneal fluid cytokines related to endometriosis in patients evaluated for infertility. *Fertil Steril* 2017; 107: 1191–1199.
- [24] Nowak M, Madej JA, Dziegiel P. Intensity of COX2 expression in cells of soft tissue fibrosarcomas in dogs as related

- to grade of tumour malignancy. *Bull Vet Inst Pulawy* 2007; 51: 275--279.
- [25] Ardianta W MY, Hendarto H, Widjiati W. Kurkumin Menu-runkan Ekspresi Tumor Necrosis Factor (TNF)- α Kompleks Oosit-Kumulus Sapi pada Kultur dengan Zalir Peritoneum Penderita Infertil Terkait Endometriosis. *Majalah Obstet Ginekol* 2015; 23: 133--139.
- [26] Scholl B, Bersinger NA, Kuhn A, Mueller MD. Correlation between symptoms of pain and peritoneal fluid inflam-matory cytokine concentrations in endometriosis. *Gynecol Endocrinol* 2009; 25: 701--706.
- [27] Hendarto H, Prabowo P, Moeloek FA, Soetjipto S. Growth differentiation factor 9 concentration in the follicular fluid of infertile women with endometriosis. *Fertil Steril* 2010; 94: 758--760.
- [28] Dong J, Albertini DF, Nishimori K, Kumar TR, Lu N, Matzuk MM. Growth differentiation factor-9 is required during early ovarian folliculogenesis. *Nature* 1996; 383: 531--535.
- [29] Orisaka M, Orisaka S, Jiang JY, Craig J, Wang Y, Kotsuji F, et al. Growth differentiation factor 9 is antiapoptotic during follicular development from preantral to early antral stage. *Mol Endocrinol* 2006; 20: 2456--2468.
- [30] Bayne RA, Kinnell HL, Coutts SM, He J, Childs AJ, Anderson RA. GDF9 is transiently expressed in oocytes before follicle formation in the human fetal ovary and is regulated by a novel NOBOX transcript. *PLoS One* 2015; 10: e0119819.
- [31] Carlsson IB, Laitinen MP, Scott JE, Louhio H, Velentzis L, Tuuri T, et al. Kit ligand and c-Kit are expressed during early human ovarian follicular development and their interaction is required for the survival of follicles in long-term culture. *Reproduction* 2006; 131: 641--649.
- [32] Mahajan A, Kumar P, Atole SA, Brahmkshtri BP, Tajane KR, Brahmkshtri BP, et al. Kit Ligand Expression in granulosa cells during follicular culture and associated maturation rate of oocytes in buffalo (*Bubalus bubalis*). *Indian J Anim Res* 2015; 50: 14--18.
- [33] Gogacz M, Bogusiewicz M, Putowski L, Adamiak A, Wertel I, Jakowicki JA, et al. Expression of tumor necrosis factor-alpha (TNF-alpha) on peritoneal fluid mononuclear cells in women with endometriosis. *Ginekol Pol* 2008; 79: 31--35.
- [34] Bedaiwy MA, Falcone T, Sharma RK, Goldberg JM, Attaran M, Nelson DR, et al. Prediction of endometriosis with serum and peritoneal fluid markers: a prospective controlled trial. *Hum Reprod* 2002; 17: 426--431.
- [35] Ma CH, Yan LY, Qiao J, Sha W, Li L, Chen Y, et al. Effects of tumor necrosis factor-alpha on porcine oocyte meiosis progression, spindle organization, and chromosome alignment. *Fertil Steril* 2010; 93: 920--926.
- [36] Jana S, Rudra DS, Paul S, Snehasikta S. Curcumin delays endometriosis development by inhibiting MMP-2 activity. *Indian J Biochem Biophys* 2012; 49: 342--348.