



Letter to Editor

Investigating the mental health status of infertile women

Dear Editor

The stressful experience of infertility is associated with a wide range of psychological damage (1), so infertility affects people's mental health and all aspects of an individual's life (2). Since women in the family are considered to be the main pillars of the community and they are also more vulnerable to illnesses, therefore consideration of their health is also very important (3). The objective of the current letter is investigating the mental health status of infertile women and its related factors as predictors of mental health in infertile women.

Corresponding Author:

Fereshteh Yazdani;
Vali-Asr Ave., Cross of Vali
Asr and Neiaiesh Highway,
Opposite to Rajaei Heart
Hospital, Tehran, Iran.
Email:
Fereshteh_yazdani68@
yahoo.com
Tel: (+98) 9112588048
Postal Code: 1996835119

Received 7 April 2018
Revised 4 July 2018
Accepted 14 November 2018

Production and Hosting by Knowledge E

© Zeinab Hamzehgardeshi
et al. This article is distributed
under the terms of the
[Creative Commons
Attribution License](#), which
permits unrestricted use and
redistribution provided that
the original author and source
are credited.

Editor-in-Chief:
Aflatoonian Abbas M.D.

This is a descriptive study conducted on 100 infertile women referred to the infertility treatment centers in Mazandaran province, North of Iran. The General Health Questionnaire was provided to the infertile women. The questionnaire is a self-reporting questionnaire that is used clinically to track those who are prepared for mental illness (4).

Based on the findings, Total Scale of General Health Questionnaire was 33.18 ± 10.27 that was according to the cut-off point of 22, 85% of the infertile women in this study are in the disorder condition. Also, in terms of subscales: physical complaints was 8.65 ± 3.97 , anxiety and insomnia was 8.69 ± 5.58 , disruption of social function was 12.73 ± 3.34 , and depression was 3.10 ± 3.79 . The most common disorder was related to social disorder subscale and the least common subscale was related to the depression disorder. The subscales of physical symptoms and sleep disturbance and anxiety are ranked almost at one level, and they are classified between the two subscales that were already mentioned.

Therefore, based on our findings and the level of women's mental health, there is a need for a psychologist or midwifery counselor in the infertility treatment centers to improve the mental health of women. In addition, since the mental disease may also affect the outcome of the treatment, attention to the mental health of infertile women is really importance.

Zeinab Hamzehgardeshi^{1,2,3} Ph.D., Fereshteh Yazdani⁴ M.Sc., Forouzan Elyasi⁵ M.D., Mahmood Moosazadeh⁶ Ph.D., Sepideh Peyvandi⁷ M.D., Keshvar Samadaee⁸ M.Sc., Maryam Shahidi^{9,10} Ph.D.

¹Sexual and Reproductive Health Research Center, Mazandaran University of Medical Sciences, Sari, Iran.

OPEN ACCESS

²Department of Reproductive Health and Midwifery, School of Nursing and Midwifery, Mazandaran University of Medical Sciences, Sari, Iran.

³Traditional and Complementary Medicine Research Center, Mazandaran University of Medical Sciences, Sari, Iran.

⁴Student Research Committee, School of Nursing and Midwifery, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

⁵Department of Psychiatry, Sexual and reproductive health research center , Psychiatry and Behavioral Sciences Research Center, Addiction Institute, School of Medicine, Mazandaran University of Medical Sciences, Sari, Iran.

⁶Health Science Research Center, Addiction Institute, Mazandaran University of Medical Sciences, Sari, Iran.

⁷IVF Ward, Mazandaran University of Medical Sciences, Sari, Iran.

⁸Department of Reproductive Health and Midwifery, Tehran Nursing and Midwifery Faculty, Tehran university of medical science, Tehran, Iran.

⁹Department of Medical Physics, Mazandaran Medical University, Mazandaran, Iran.

¹⁰Hazrat_e Maryam Fertility Center, Sari, Iran.

References

- [1] Besharat MA, Hosseinzade Bazargani R. Comparison of mental health and sexual problems of fertile and infertile women. *J Psychiatry Clin Psychol* 2005; 12: 146–153.
- [2] Hamzehgardeshi Z, Yazdani F, Elyasi F, Moosazadeh M, Peyvandi S, Samadaee Gelehkolaee K, et al. The efficacy of group counselling on perceived stress among infertile women undergoing in vitro fertilization treatment: An RCT. *Int J Reprod BioMed* 2019; 17: 57–66.
- [3] Jamilian M, Rafiei M, Jamilian HR, Esmkhani F. [The comparison of general health between fertile and infertile women of Arak City in 2010.] *Arak Med Uni J* 2011; 14: 27–35. (in Persian)
- [4] Latifnejad Roudsari R, Rasolzadeh Bidgoly M, Mousavifar N, Modarres Gharavi M. The effect of collaorative counseling on perceived infertility-related stress in infertile women undergoing IVF. *Women Obstet Infertil Iran* 2011; 14: 22–31.