

Expression of CXC Chemokines Gro/KC and SDF-1a in Rat H4 Hepatoma Cells in Response to Different Stimuli

Gholamhossein Hassanshahi¹, Mohammad Kazemi Arababadi¹, Alan James Dickson²

¹Immunology and Hematology Departments, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran, ²Biochemistry Department, Faculty of Life Sciences, Manchester University, Manchester, UK.

ABSTRACT

Background: It is now well established that several environmental stress factors cause activation of p38 MAP kinase and JNK in various cell types to produce chemokines. **Objective:** To investigate the expression of CXC chemokines Gro/KC and SDF-1 α in rat's H4 hepatoma cells in response to heat shock, hyperosmolarity and oxidative stress. **Methods:** Hepatoma cells were maintained in MEM medium. Cells were subjected to different stresses [(H₂O₂ 0.15% (w/v), mannitol and NaCl (160 mM) and heat shock (42 °C for 20 minutes)]. Cells were harvested and RNA was extracted, purified and the CXC chemokine Gro/KC and SDF-1 α expression was analysed by RT-PCR. cDNA was separated by gel electrophoresis on a 1% (w/v) agarose gel and visualized under a UV transilluminator. **Results:** There was detectable but low expression of both SDF-1 α and Gro/KC in H4 hepatoma cells. Heat shock failed to induce expression of SDF-1 α and Gro/KC in H4 hepatoma cells of rat. Hyperosmolarity also did not stimulate SDF-1 α and Gro/KC expression. In this study we have also shown that oxidative stress did not induce expression of SDF-1 α and Gro/KC. Overall, although detection is possible but regulatory responses were not observed in H4 hepatoma cells. **Conclusion:** Several known injurious conditions cause recruitment of macrophages, neutrophils and other immune cells to the liver. Immune cells are recruited to the hepatic vasculature following local liver injury and subsequent chemokine production. Our results demonstrated that failure to produce chemokines by hepatoma cells may be a way to escape from mechanism of immune surveillance.

Keywords: Hepatoma, SDF-1a, Gro/KC, Chemokine

Corresponding author: Dr. Gholamhosein Hasanshahi, Department of Immunology, Medical School, Rafsanjan University of Medical Sciences, Rafsanjan, Iran. e-mail: ghassanshahi@gmail.com

INTRODUCTION

It is now established that several environmental stress factors including UV radiation, DNA damage, pro-inflammatory cytokines, heat shock, oxidative, and osmolarity stress can cause the activation of p38 MAP kinase and JNK in various cell types (1). The effect of osmotic shock (e.g NaCl) on expression of IL-8 in human monocytes and bronchial epithelial cells has been shown to occur through a p38-dependent pathway (1, 2). The roles of p38, p42/44, and NF- κ B in the expression of IL-8 and Gro/KC in response to hyperosmolarity in human intestinal epithelial cells have been identified (3). Oxidative stress is associated with several liver disorders such as bacterial and viral liver infections; toxin induced liver injury, ethanol consumption, and hepatocellular carcinomas (4). These situations are associated with chronic inflammatory response and increased expression of Gro/KC, MIP-2, and IL-8 in rat and mouse hepatocytes and human Hep-G2 cells (5). H₂O₂ also activates NF- κ B and regulates IL-8 expression in epithelial cells of lung (6). In this research, H4 hepatoma cell line was used as a model to detect CXC chemokines because it is a liver cell line which is reported to be reasonably differentiated and which has previously been well characterised. We investigated the expression of CXC chemokines Gro/KC as an ELR⁺, and SDF-1 α as an ELR⁻ CXC chemokines in rat's H4 hepatoma cells in response to heat shock, hyperosmolarity, and oxidative stress to explore the pattern of expression of these CXC chemokines by hepatoma cells.

MATERIALS AND METHODS

Hepatoma cells were maintained in MEM (minimal essential medium). Cells were subjected to different stress factors [(H₂O₂ 0.15% (w/v), manitol and NaCl (160 mM) and heat shock (42°C for 20 minutes)]. Cells were harvested, RNA was extracted, purified, and expression of the CXC chemokines SDF-1 α and Gro/KC was analysed by RT-PCR. cDNA was separated by gel electrophoresis on a 1% (w/v) agarose gel and visualized under a UV transilluminator. GAPDH mRNA was assessed in parallel as control.

H4 Hepatoma Cell Culture. Rat hepatoma H4-II-C3 cells were obtained from ECACC (European Collection of Animal Cell Culture, Porton Down, Wiltshire, UK). Cells were incubated at 37°C in atmospheric air: 5% CO₂. After 24 hours, medium and any unattached cells were removed and 20 ml of fresh growth medium was added. Cells were then subjected to routine subculture. For the maintenance of subculture, culture medium was removed and the cell sheet was washed with PBS. Cells were detached with gentle agitation and addition of 2 ml sterile trypsin (200mg/ml). Then fresh medium was added to stop trypsin action and the cell suspension was centrifuged at 400g for 5 min at room temperature. The supernatant was discarded, the pellet was suspended in fresh medium, and cells were diluted in medium to generate a density of 3.0-3.5 x 10⁵ cells/ml in T75 ml flasks. Incubation was continued at 37°C and subculture was repeated until cells were 70% confluent (after about 3-5 days of culture).

Reverse Transcription Polymerase Chain Reaction (RT-PCR). To make complementary DNA (cDNA), reverse transcription reactions were performed using the following protocol: 4 μ l 5x strand buffer [125mM tris-HCl pH 8.3, 188 mM KCl, 7.5

mM MgCl₂ 25 mM DTT]; 1 μ l of each dNTP [dATP, dCTP, dGTP, dTTP (stock concentration of 10mM in DEPC-treated water)]; 4 μ l oligo-dT (stock concentration of 125 μ g/ml); 1 μ l RNA (1 μ g/ μ l); 4 μ l DEPC-treated water; 1.5 μ l M-MLV reverse transcriptase enzyme. After addition of M-MLV-reverse transcriptase and mixing, the tube was incubated for 1 hour at 37°C. The resultant product was stored at -20°C for further use in PCR reactions. To amplify DNA species, PCR reaction mixture was prepared by addition of the following reagents to a 0.5 ml micro-centrifuge tube in ice: 10 μ l *Taq* polymerase buffer (10x); 3 μ l MgCl₂ (stock concentration 1.5mM); 2 μ l of each dNTP [(dATP, dCTP, dGTP, dTTP) stock concentration of 10mM]; 2 μ l of each primer pair [(forward and reverse) stock concentration of 25ng/ μ l]; 4 μ l cDNA; and sterile double distilled water to a final volume of 99 μ l. The above mixture was overlaid with 40-60 μ l mineral oil. The PCR thermocycler was adjusted accordingly: 94 °C for 5 minutes, 94 °C for 1 minute (denaturation), 1 minute at appropriate annealing temperature for annealing of different targets (Table 1), and 72°C for 2 minutes (elongation). Denaturation, annealing and elongation procedures were repeated for 30 cycles and were followed by a final reaction at 72°C for 10 minutes (final elongation) and held at 4°C up to 24h. During the last 45 seconds of first stage 1 μ l *Taq* polymerase was added to the mixture. In addition to appropriate annealing and melting temperatures, table 1 indicates the size of products that should be achieved by PCR.

Table 1. Primers used to detect CXC chemokine mRNA

Gene Name	Primers (5' to 3')	AT (°C)	MT(°C)	Fragment Size
GAPDH	F: aagtcggtgtgaacggatt	57	62	835
	R: gagacaacctggtcctcagttag		62	
Gro/KC	F: ttcaactcaagaacatcca	51	56	680
	R: gagcattggttaaagaatataaa		56	
SDF-1 α	F: agtgtgcattgacccgaaatta	60	65	1500
	R: ttacagcacgaaacagttggc		65	

Abbreviations: AT= annealing temperature; F= forward primer; GAPDH= glyceraldehyde-3-phosphate dehydrogenase; MT= melting temperature; R=reverse primer

RESULTS

Results obtained in this report revealed that there were detectable but low expression of both SDF-1 α and Gro/KC in H4 hepatoma cells. Heat shock failed to induce expression of SDF-1 α and Gro/KC in H4 hepatoma cells of rat. Hyperosmolarity also did not stimulate SDF-1 α and Gro/KC expression. In this study we also noted that oxidative stress did not induce expression of SDF-1 α and Gro/KC. Overall, although detection is possible but regulatory responses were not observed in H4 hepatoma cells. (Figures 1-3)




Figure 1. Expression of Gro/KC under heat shock, hyperosmolarity, and oxidative stresses in rat's H4 Hepatoma cells. Expression of CXC chemokine Gro/KC was analyzed by RT-PCR. GAPDH mRNA was assessed in parallel as control. Numbers on right represent the bands of the molecular marker. HC= hepatocytes control; NT= not treated; 1= heat shock treatment for 1h; 2= heat shock treatment for 3h; 3= heat shock treatment for 6h; 4= Manitol treatment for 1h; 5= Manitol treatment for 3h; 6= Manitol treatment for 6h; 7= H₂O₂ treatment for 1h; 8= H₂O₂ treatment for 3h; 9= H₂O₂ treatment for 6h; 10= NaCl treatment for 1h; 11= NaCl treatment for 3h.




Figure 2. Expression of SDF-1 α under heat shock, hyperosmolarity, and oxidative stresses in rat's H4 hepatoma cells. Expression of the CXC chemokine SDF-1 α was analysed by RT-PCR. GAPDH mRNA was assessed in parallel as control. Numbers on right represent the bands of the molecular marker. HC=hepatocytes control; NT=not treated; 1= heat shock treatment for 1h; 2=heat shock treatment for 3h; 3=heat shock treatment for 6h; 4=Manitol treatment for 1h; 5=Manitol treatment for 3h; 6= Manitol treatment for 6h; 7=H₂O₂ treatment for 1h; 8= H₂O₂ treatment for 3h; 9= H₂O₂ treatment for 6h; 10= NaCl treatment for 1h; 11= NaCl treatment for 3h.




Figure 3. Alteration in expression of CXC chemokines SDF-1 α and Gro/KC against GAPDH in rat's hepatoma H4 cells in different conditions and in the presence and absence of different stimuli. HC=hepatocytes control; NT=not treated; S1=heat shock treatment for 1h; S3=heat shock treatment for 3h; S6=heat shock treatment for 6h; M1=Manitol treatment for 1h; M3=Manitol treatment for 3h; M6=Manitol treatment for 6h; H1=H₂O₂ treatment for 1h; H3=H₂O₂ treatment for 3h; H6=H₂O₂ treatment for 6h; N1=NaCl treatment for 1h; N3= NaCl treatment for 3h.

DISCUSSION

Several known injurious conditions such as hepatitis, ischemia/reperfusion, sepsis/endotoxemia, and drugs cause recruitment of macrophages, neutrophils, and other immune cells to the liver parenchyma (7). Neutrophils are recruited to the hepatic vasculature following local liver injury and subsequent CXC chemokine production (8, 9) or following exposure to inflammatory cytokines (10, 7). Although neutrophils and Kupffer cells are widely considered to be the first defence line in liver, the liver immune response is not restricted to these cells, and other immune cells such as T lymphocytes and NK cells are also involved, mostly mediated by chemokines (11). Involvement of CC chemokines such as MCP-1, ELR⁺ CXC chemokines (such as IL-8/MIP-2) and ELR⁻ chemokines (such as IP-10/Mob-1) has been reported (12, 13) in acetaminophen-induced liver injury and modulation of liver regeneration processes (14, 15). So far this report is the only one on expression of chemokines by H4 hepatoma cells. High constitutive levels of SDF-1 α have been observed in the non-inflamed biliary epithelium of the liver (18) in association with CXCR4 expressing lymphocyte recruitment (19, 20). Immunohistochemical studies have revealed decreased levels of SDF-1 α protein in hepatocellular carcinoma compared with other chronic liver diseases such as hepatitis C (21). Expression of Gro/KC in hepatectomized mouse has been shown and it may play an important role in hepatocyte proliferation and liver regeneration (22). Increased Gro/KC expression is associated with some hepatic injuries and diseases including: liver ischemia/reperfusion in mouse and rat (23), infections, paracetamol poisoning, sepsis and LPS injection in mice (24). Results obtained in this report revealed that there was detectable but low expression of both SDF-1 α and Gro/KC in H4 hepatoma cells. In contrast to primary hepatocytes, heat shock did not induce expression of SDF-1 α and Gro/KC in rat's H4 hepatoma cells (25). Hyperosmolarity also failed to stimulate SDF-1 α and Gro/KC expression.

ACKNOWLEDGEMENTS

The authors would like to thank Mrs. Louise Barnes for her help and advice throughout the project. We would also like to thank the members of Lab 2.300, Stopford Building, University of Manchester, for their help and patience. This project is financially supported by Ministry of Health and Medical Education of Islamic Republic of Iran.

REFERENCES

1. Hashimoto S, Matsumoto K, Gon Y, Nakayama T, Takeshita I, Horie T. Hyperosmolarity-induced interleukin-8 expression in human bronchial epithelial cells through p38 mitogen-activated protein kinase. *American Journal of Respiratory Critical Care Medicine*. 1999; 159: 634-640.
2. Shapiro L, Dinarello CA. Osmotic regulation of cytokine synthesis in vitro. *Proceeding of National Academy of Science of USA*. 1995; 92: 12230-12234.
3. Nemeth ZH, Deitch EA, Szabo C, Hasko G. Hyperosmotic stress induces nuclear factor-kappaB activation and interleukin-8 production in human intestinal epithelial cells. *American Journal of Pathology*. 2002; 161: 987-996.
4. Schmelz M, Schmid VJ, Parrish AR. Selective disruption of cadherin/catenin complexes by oxidative stress in precision-cut mouse liver slices. *Toxicological Sciences*. 2001; 61: 389-394.
5. Dong W. et al. Toxic metals stimulate inflammatory cytokines in hepatocytes through oxidative stress mechanisms. *Toxicol Appl Pharmacol*. 1998; 151: 359-366.

6. Luster AD. Chemokines--chemotactic cytokines that mediate inflammation. *New England Journal of Medicine*. 1998; 338: 436-445.
7. Bast A, Wolf G, Oberbaumer I, Walther R. Oxidative and nitrosative stress induces peroxiredoxins in pancreatic beta cells. *Diabetologia*. 2002;45:867-76.
8. Lawson JA, Farhood A, Hopper RD, Bajt ML, Jaeschke H. The hepatic inflammatory response after acetaminophen overdose: role of neutrophils. *Toxicol Sci*. 2000;54:509-16.
9. Maher JJ, Lozier JS, Scott MK. Rat hepatic stellate cells produce cytokine-induced neutrophil chemoattractant in culture and in vivo. *Am J Physiol*. 1998;275:847-53.
10. Jaeschke H, Fisher MA, Lawson JA, Simmons CA, Farhood A, Jones DA. Activation of caspase 3 (CPP32)-like proteases is essential for TNF-alpha-induced hepatic parenchymal cell apoptosis and neutrophil-mediated necrosis in a murine endotoxin shock model. *J Immunol*. 1998;160:3480-6.
11. Bone-Larson CL, Simpson KJ, Colletti LM, Lukacs NW, Chen SC, Lira S et al. The role of chemokines in the immunopathology of the liver. *Immunol Rev*. 2000;177:8-20.
12. Koniaris LG, Zimmers-Koniaris T, Hsiao EC, Chavin K, Sitzmann JV, Farber JM. Cytokine-responsive gene-2/IFN-inducible protein-10 expression in multiple models of liver and bile duct injury suggests a role in tissue regeneration. *J Immunol*. 2001;167:399-406
13. Ishida Y, Kondo T, Ohshima T, Fujiwara H, Iwakura Y, Mukaida N. A pivotal involvement of IFN-gamma in the pathogenesis of acetaminophen-induced acute liver injury. *FASEB J*. 2002;16:1227-36.
14. Hogaboam CM, Bone-Larson CL, Steinhauser ML, Lukacs NW, Colletti LM, Simpson KJ, Strieter RM, Kunkel SL. Novel CXCR2-dependent liver regenerative qualities of ELR-containing CXC chemokines. *FASEB J*. 1999;13:1565-74
15. Ren X, Carpenter A, Hogaboam C, Colletti L. Mitogenic properties of endogenous and pharmacological doses of macrophage inflammatory protein-2 after 70% hepatectomy in the mouse. *Am J Pathol*. 2003;163:563-70.
16. Bone-Larson CL, Hogaboam CM, Evanhoff H, Strieter RM, Kunkel SL. IFN-gamma-inducible protein-10 (CXCL10) is hepatoprotective during acute liver injury through the induction of CXCR2 on hepatocytes. *J Immunol*. 2001;167:7077-83.
17. Kobayashi H, Narumi S, Tamatani T, Lane GJ, Miyano T. Serum IFN-inducible protein-10: a new clinical prognostic predictor of hepatocyte death in biliary atresia. *J Pediatr Surg*. 1999;34:308-11.
18. Coulomb-L'Herminé A, Amara A, Schiff C, Durand-Gasselin I, Foussat A, Delaunay T et al. Stromal cell-derived factor 1 (SDF-1) and antenatal human B cell lymphopoiesis: Expression of SDF-1 by mesothelial cells and biliary ductal plate epithelial cells. *PNAS*. 1999;96: 8585-8590.
19. Terada R, Yamamoto K, Hakoda T, Shimada N, Okano N, Baba N et al. Stromal cell-derived factor-1 from biliary epithelial cells recruits CXCR4-positive cells: implications for inflammatory liver diseases. *Lab Invest*. 2003;83:665-72.
20. Goddard S, Williams A, Morland C, Qin Sh, Gladue R, Hubscher SG et al. Differential expression of chemokines and chemokine receptors shapes the inflammatory response in rejecting human liver transplants. *Transplantation*. 2001;72:1957-1967.
21. Mitra P, De A, Ethier MF, Mimori K, Kodys K, Shibuta K et al. Loss of chemokine SDF-1alpha-mediated CXCR4 signaling and receptor internalization in human hepatoma cell line HepG2. *Cell Signal*. 2001;13:311-9.
22. Su AI, Guidotti LG, Pezacki JP, Chisari FV, Schultz PG. Gene expression during the priming phase of liver regeneration after partial hepatectomy in mice. *Proc Natl Acad Sci U S A*. 2002;99:11181-6.
23. Hisama N, Yamaguchi Y, Ishiko T, Miyanari N, Ichiguchi O, Goto M et al. Kupffer cell production of cytokine-induced neutrophil chemoattractant following ischemia/reperfusion injury in rats. *Hepatology*. 1996;24:1193-8.
24. Salkowski CA, Detore G, Franks A, Falk MC, Vogel SN. Pulmonary and hepatic gene expression following cecal ligation and puncture: monophosphoryl lipid A prophylaxis attenuates sepsis-induced cytokine and chemokine expression and neutrophil infiltration. *Infect Immun*. 1998;66:3569-78.
25. Varley CL, Armitage S, Hassanshahiraviz G, Dickson AJ. Regulation of the C-X-C chemokine, mob-1, gene expression in primary rat hepatocytes. *Cytokine*. 2003;23:64-75.