# Prevalence of *Helicobacter pylori* Infection in Arak, Iran during 2011

### <u>Ali Fani</u><sup>1</sup>, Mohammad Rezaei<sup>2</sup>, Behruz Alizade<sup>1</sup>, Parisa Mirzajani<sup>1</sup>, Soheila Shamsikhani<sup>3</sup>, Mohammad Rafeie<sup>1</sup>, Ali Akbar Malekirad<sup>4</sup>, Saeed Baghi Nejad<sup>5</sup>

<sup>1</sup> Arak University of Medical Sciences, Arak, Iran

<sup>2</sup> Department of Food Safety and Hygiene, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran

<sup>3</sup> Faculty of Midwifery and Nursing, Arak University of Medical Sciences, Arak, Iran

<sup>4</sup> Payame Noor University, Tehran, Iran

<sup>5</sup> Amiralmomenin Hospital, Arak University of Medical Sciences, Arak, Iran

## ABSTRACT

#### **Background:**

*Helicobacter pylori* (*H. pylori*) is a global pathogen with a widely varied prevalence (30%-80%) among different countries and populations. This study aims to determine the prevalence of *H. pylori* in the population of Arak, Iran. *Materials and Methods:* 

This was an epidemiologic cross-sectional population-based study with multistage sampling. The population of Arak according to the 2006 census was 615,722 (493,845 urban and 121,857 rural residents). In this survey, five urban and four rural Health Service Centers were randomly selected. The study population was divided between these centers based on the population of individuals covered by these centers. Subsequently, clustered samples were randomly chosen according to Health Care Unit Family Codes and the Right Hand Rule. After obtaining informed consent and completion of a demographic questionnaire, 5 cc of venous blood was taken from each participant for the *H. pylori* IgG antibody test. In this study IgG antibody against *H. pylori* was measured by ELISA. *Results:* 

Of 1187 participants, 1150 (853 urban and 297 rural) completed the study. Overall, 673 (58.5%) were positive for *H. pylori*. The prevalence of *H. pylori* infection increased with age from 43.9% among those <20 years old to >70% in participants over 50 years old (p=0.002). A total of 57% urban samples and 63.3% rural samples were *H. pylori* positive (p=0.028). There was no relationship between *H. pylori* infection and education level (p=0.37), career (p=0.39) and income, (p=0.29) in either the urban or rural areas [p=0.64 (education level), 0.48 (career), and 0.57 (income)]. *Conclusion:* 

Our study showed a lower prevalence of *H. pylori* compared to some Asian countries, however this prevalence was more common than Western countries. Using disinfected tap water in both urban and rural areas in addition to improved access to general health care might have a role in this relative lower prevalence.

Keywords: Prevalence; Helicobacter pylori; Infection; ELISA; Arak, Iran

#### please cite this paper as:

Fani A, Rezaei M, Alizade B, Mirzajani P, Shamsikhani S, Rafeie Mohammad, Malekirad AA, Baghi Nejad S. Prevalence of *Helicobacter pylori* Infection in Arak, Iran during 2011. *Govaresh* 2014;19:57-62.

#### **Corresponding author:**

Ali Fani, MD Arak University of Medical Sciences, Arak, Iran Tel: +98 918 1617447 E-mail: drfani321@yahoo.com Received: 20 Jan. 2014 Edited: 28 Feb. 2014 Accepted: 29 Feb. 2014

#### **INTRODUCTION**

*Helicobacter pylori* (*H. pylori*) was first described by Warren and Marshall(1). *H. pylori* is a major cause of gastritis and peptic ulcer disease (PUD) and has been implicated in the development of gastric malignancies and common worldwide infections (2-6). Previous sero-epidemiologic studies have estimated that approximately 50% of adults worldwide, 20%-30% in developed countries and up to 80% in developing countries are positive for serum

antibodies against H. pylori WGO 2010(5). The prevalence is declining in most developed countries (4,6,7). Epidemiological studies have shown that infection is mainly acquired in early childhood and the incidence increases with age (8-10). Close contact, low socioeconomic status and poor hygiene are the most important risk factors (5,10,11). It seems that the fecal-oral route is the main basis of the infection (11-13). Primary acquisition in adults or reinfection after successful eradication does occur but is less common with an annual incidence of 0.3%-0.7% in developed countries and 6%-14% in developing countries (5, 11). Since humans are the main known reservoir for infection, it is likely that in developed countries H. *pylori* is acquired predominantly via the oral-oral and fecal-oral routes (11-13). Since the acute acquisition of H. pylori infection usually remains undiagnosed, the incidence has been determined indirectly through epidemiological studies. Some discrepancies between the prevalence and incidence data may be explained by the decreasing rates of infection over time. Thus, adults who are currently infected are more likely to have acquired their disease in childhood rather than during adulthood (2-5,14).

Water has been suggested as a possible source of H. pylori infection (15-24). Studies by Dube in Africa(25), Shi et al (26), and Kelin in Peru(14), Hopkin in Chile (27) have found that the water source may be related to the risk of H. pylori infection. One of the most important factors increases risks of H. pylori infection in children, drinking water from a stream that may be vulnerable to bacterial contamination. Studies have also assessed the relation between H. pylori infection and various hygiene practice indicators in a number of countries. Poor hygiene practices, especially during childhood, appear to be related to a higher sero-prevalence of H. pylori. Some practices include the lack of a water closet or bathroom or no hot water supply in the house during childhood, sharing cups as children, and having mothers who did not use soap when they washed their hands (20,27-29). Although there are different tests for the detection of *H. pylori*, the ELISA method remains a reliable and cost effectiveness test for epidemiologic studies (5,30). Public knowledge and expectations have also increased, with numerous patients requesting information about the chances of acquiring the infection, risk of peptic ulcer, gastric cancer and routes of transmission. In order to provide accurate answers, it is necessary to conduct epidemiological studies specific to areas of interest and update data with improvement of health level. This study intends to determine the prevalence of *H. pylori* infection in the general adult population of Arak, Markazi Province, Iran.

#### **MATERIALS AND METHODS**

This cross-sectional epidemiologic populationbased study was conducted during a five-month period from March to July 2011 to determine the prevalence of H. pylori infection on people over the age of 20 years in Arak, Markazi Province, Iran. The population of Arak according to the 2006 census was 615,702 (493,845 urban and 181,857 rural). This study recruited 1150 participants based upon a sample size that took into consideration the world wide prevalence (4) of 20%-80%, p=50%, d=0.03 and  $\alpha=0.05$  and was estimated at 1150 (853 urban, 297 rural) individuals who represented the general population based on the weighted multistage stratified cluster pattern for sampling. All individuals over the age of 15 years were enrolled with respect to the urban-rural population distribution. The Arak Township Health Service is covered by 11 urban and 8 rural Health Service Centers. In this survey 5 urban and 4 rural Health Service Centers were randomly selected. The study participants were divided between these centers based on the population covered by these centers. Subsequently, the clustered sample was randomly chosen according to the Health Care Unit Family Codes and Right Hand Rule. After obtaining informed consent and completion of the demographic questionnaire that included items such as gender, age, height, weight, education level, family member, career and family income, a total of 5 cc of venous blood was taken from each participant for the H. pylori IgG antibody test. Serum was directly separated and stored at -20°C. In case of the participant refuses, samples were collected from the nearest neighbor. H. pylori status of the samples was determined by the ELISA IgG Antibody Mono-bind Inc. H. pylori Kit (no CA92630, USA) and by a Stat Fax 3200 (USA) ELISA reader that had a sensitivity of 95% and specificity of 98%. Any titer above 20 units was considered positive. Data were analyzed with SPSS version 11.5, the chi-square test, logistic regression and t-test. *p*-values <0.05 were considered statistically significant.

#### **RESULT** =

From 1187 participants, 1150 participants (853 urban and 297 rural) completed the study. Participants' mean age was 39.22±12.7 years. Overall, 673 (58.5%) had evidence of antibodies to *H. pylori* (ELISA IgG- Ab+); 57% urban and 63% rural samples were antibody positive. According to two-tailed analysis, there was no difference

in the positive H. pylori results between urban and rural populations (p=0.379). However in the one-tailed test, this overall difference was meaningful (p=0.038). The prevalence of H. pylori infection increased with age from 43.9% among <20 year-old group to >70% in those over 50 years of age (p=0.002). There was no relationship between *H. pylori* infection and education level (p=0.37), career (p=0.35) and income (p=0.29) for urban residents, or for rural residents in terms of education level (p=0.64), career (p=0.48), and income (p=0.57). Regarding gender, 57% of women and 60.08% of men were H. pylori positive which was not significant (p=0.335). Among males, 59.5% of urban and 62.1% of rural groups were seropositive (p=0.092). In females, 54.7% of urban and 64.8% of rural groups were seropositive (p=0.028). Average BMI in the H. pylori negative group was 25.03±4.55 whereas in the H. pylori positive group it was  $25.94\pm16.14$  (p=0.238). Heartburn symptoms were seen in 11.3% of seronegative and 16.9% of seropositive participants (p=0.207). In Arak, all study participants in both urban and rural areas used chlorinated closed well water.

#### **DISCUSSION**

We used epidemiologic study data to produce the population-based estimation of the prevalence of H. *pylori* infection in Arak and explored the relationships between various demographic factors of infection. There was no significant difference in the prevalence of *H. pylori* between the urban and rural population (p=0.379). The lower rate of infection in the urban population could be related to improvements in public health care services (availability of hot water and soap use). In this study, in terms of gender we observed no difference in the prevalence of H. pylori in both the rural and urban areas. However this prevalence was slightly more common in females from rural areas compared to urban females, which might be related to the family crowding and poorer hygienic conditions in rural areas. Few epidemiologic and community-based studies that determined the *H. pylori* infection in the general population have been performed in Iran. The majority of these studies were undertaken in gastroenterology clinics (16,24,30). Overall, our survey results showed lower prevalence than previous epidemiologic studies conducted in Hormozgan Province (67.1%) and Nahavand 71% (16,19). In the <20 year old age group, we reported a prevalence of 43.9%, which was more common than the prevalence in Yazd that had a 30.6% H. pylori sero-positivity (p=0.045) (18). This difference was significant. In comparison with Ardabil that reported a 47.5% sero-positivity, the difference was

not significant. A study in Shiraz (24) reported 52.7% H. pylori positivity. Although their findings were similar to the findings of the current study, these results might not be comparable because subjects in the Shiraz study were selected from patients with dyspepsia who had higher rates of current infection and were not representative of the general population. Results of these studies have shown that the prevalence of *H. pylori* infection in Iran varies according to ethnicity, cultural and healthcare factors. The overall prevalence of *H. pylori* infection in our study was lower than some published studies from other developing countries (22, 32). In comparison with Asian countries such as Kazakhstan, Bangladesh and India, we have shown that *H. pylori* infection was less common in Iran (Arak) and inversely correlated with the health care system and more common use of closed wells and hot water. Family crowding showed no significant effect on H. pylori prevalence in Iran and Kazakhstan (15). H. pylori infection appeared to be less common in Iran compared to other Middle Eastern countries. An overall estimation of *H. pylori* prevalence in Turkey was 82.5% according to the Urea Breath Test (UBT) method as reported by Sinan (31). Although comparable with some Iranian studies, there was a meaningful difference with the results of the current study (p=0.001). In a review article from Africa(25), the prevalence of H. pylori infection was reported to be 74.8%-92% and depended on the drinking water source (tap water versus well water) which was more common than Iran, particularly Arak. Tkachenko et al. (22) in a 10-year follow-up study in Russia observed a significant drop from 48% in 1995 to 25% in 2005 in H. pylori infection among 15-19 year olds which was concurrent with health care improvements. They concluded that there was an inverse correlation between mothers' and fathers' educational levels and H. pylori sero-positivity. In our study there was no association between the prevalence of H. pylori and any factor tested, including sex, type of dwelling, income, or the number of people living in the home. These results were inversely comparable with the same age group findings in a Russian study (p=0.017) (22). Naja et al. (8), conducted a sero-epidemiologic study in Ontario, Canada among individuals 50 to 80 years of age. In that study a prevalence of 23.1% with a higher prevalence among males was reported. The sample size and methods were similar to the current study, however the age factor differed. Staat et al. (23) studied H. pylori infection in the US population by measuring IgG antibody levels in serum from 2581 subjects aged 6-19 years. Overall, 24.8% of participants had evidence of H. pylori infection which was less common than our

<b>Developing Countries</b>	H. pylori positive		itive	<b>Developed</b> Countries	H. pylori positive		sitive	
Country	Age group <20 years Adults		Adults	Country	Age group <20 years		Adults	
Africa	-		-	North America				
Ethiopia	<6	80%	95%	Canada	<20 7.1	.%	50-80	23.1%
Nigeria	<10	82%	91%	USA and Canada	-		30	)%
Central & South America	-		-	Hong Kong	<20 13.	1%		-
Guatemala	<10	51%	70-90%	Japan	-		20-70	55.4%
Mexico	<10	43%	70-90%	Taiwan	13–15 12	.3%	>25	45.1%
Brazil	10-19	78%	82%	Australasia	<20 15.	.4%	20	)%
Chile	<10	36%	70-90%	Europe				-
Asia	-		-	Eastern Europe			70	)%
Bangladesh	<10	82%	90%	Eastern Europe			30-3	50%
India	<20	87%	88%	Germany	-		48.	8%
Turkey	General po	opulation	82.5%	Iceland			36	5%

 Table 1: World Gastroenterology Organization Global Guidelines for Helicobacter pylori (H. pylori) in developing and developed countries, August 2010.

 Table 2: Summary and prevalence comparison of *H. pylori* in residents <20 years of age from Arak, Iran with studies in Iran and other countries.</th>

			Method				
Country	Editor, year published	Study population (years)	Sample size and method	Negative	Positive	<i>p</i> -value	
Iran, Arak	Present study, 2011	<20	66 ELISA	37 (56.06%)	29 (43.94%)		
Iran, Ardabil	Mikaeili(18), 1999	<20	358 ELISA	188 (52.5%)	170 (47.5%)	0.688	
Iran, Yazd	Mikaeili(18), 1999	<20	353 ELISA	245 (59.4%)	108 (30.6%)	0.045	
Turkey	Apan(20), 2008	<20	529 ELISA	370 (70%)	159 (30%)	0.025	
Russia	Tkachenko(22), 2005	15-19	100 ELISA	75 (75%)	25 (25%)	0.017	
Kazakhstan	Nurgalieva(15), 2002	15-19	55 ELISA	20 (36%)	35 (64%)	0.044	

Table 3: Summary and prevalence comparison of H. pylori in Arak, Iran with studies held in Iran and other countries (general population).

			Method				
Country	Editor, year published	Population selection	Sample size and method	Negative	Positive	<i>p</i> -value	
Iran, Arak	Present study, 2011	General population	1150 ELISA	477 (41.5%)	673 (58.5%)		
Iran, Hormozgan	Hashemi(16), 2006	Adult dyspeptic	1000 RUT1	329 (32.9%)	671 (67.1%)	0.001	
Turkey	Sinan(31), 2009	General population	2382 ELISA		82.5%	0.001	
Iran, Nahavand	Alizadeh(19), 2009	General population	1518 ELISA	449 (29%)	1078 (71%)	0.001	
India	Klein(14), 1991	General population	238 ELISA	50 (17.5%)	188 (79%)	0.001	
Canada, Toronto	Naja(8), 2007	50-80 years	1306 ELISA	922 (70.6%)	384 (29.4%)	0.001	

<sup>1</sup>Rapid urease test

	Age group (years)	١	Urban		n voluo	
		Samples (n)	Seropositive n (%)	Samples (n)	Seropositive n (%)	<i>p</i> -value
	<20	35	14 (40)	31	15 (48.4)	0.824
	21-30	219	153 (61.2)	62	32 (51.6)	0.902
	31-40	250	153 (61.2)	85	58 (69.4)	0.901
	41-50	186	112 (60.2)	55	35 (63.6)	0.901
	51-60	115	66 (57.4)	45	32 (71.1)	0.484
	61-70	40	24 (60)	19	15 (73.7)	0.665
	>70	8	7 (87.5)	0	0	-
	Sum	853	486 (57)	297	187 (63)	0.379
	·					

Table 4: Comparison of H. pylori sero-positive age groups in urban and rural populations of Arak, Iran during 2011.

Table5: The relation	ationship betw	een H. pylori and	family income (	toumans/month/	person)
		1.			

	Low income (<100000)	Moderate income (200000-400000)	High income (>400000)	<i>p</i> -value
H. pylori negative	137 (28.7%)	328 (68.8%)	12 (2.5%)	0.209
H. pylori positive	215 (32%)	453 (67.3%)	5 (0.7%)	
Total	352 (30.6%)	781 (67.9%)	17 (1.5%)	

results although *H. pylori* infection in the US was strongly associated with increasing age and ethnicity (p<0.01). A total of 17.0% of non-Hispanic whites, 40.1% of non-Hispanic blacks and 42% of Mexican Americans were infected. The *H. pylori* prevalence in Arak was more common than non-Hispanic whites and approximated the results of non-Hispanic blacks and Mexican Americans.

There was a lower prevalence of *H. pylori* compared to other studies in the Middle and Far East, excluding

#### REFERENCES

- Robin Warren J, Marshall B. Unidentified curved bacilli on gastric epithelium in active chronic gastritis. *Lancet* 1983;321:1273-5.
- 2. Suerbaum S, Michetti P. Helicobacter pylori infection. *N Engl J Med* 2002;347:1175-86.
- Feldman R. Epidemiologic observations and open questions about disease and infection caused by Helicobacter pylori. Achtman M, Serbaum S Helicobacter pylori: Molecular and Cellular Biology Wymondham: *Horizon Scientific* 2001:29-51.
- Hunt RH, Xiao SD, Megraud F, Leon-Barua R, Bazzoli F, van der Merwe S, et al. Helicobacter pylori in developing countries. World Gastroenterology Organisation Global Guideline. J Gastrointestin Liver Dis 2011;20:299-304.
- 5. Fani A, Fani P, Rafiei M, Shahmirzaei R, Minavand

Japan, India and Africa. Using disinfected well water in both urban and rural areas and improved access to general health care might have had a role in this relative lower prevalence.

#### ACKNOWLEDGMENT

This study was approved and supported by the Research and Ethics Committees of Arak University of Medical Sciences and conducted with regards to the Declaration of Helsinki ethical principles.

K. The impact of helicobacter pylori infection in dyspepsia in low dose Aspirin takers. *Arak Med Univ J* 2010;13:105-12.

- Persson C, Jia Y, Pettersson H, Dillner J, Nyrén O, Ye W. H. pylori seropositivity before age 40 and subsequent risk of stomach cancer: a glimpse of the true relationship? *PloS One* 2011;6:e17404.
- Everhart JE, Kruszon-Moran D, Perez-Perez GI, Tralka TS, McQuillan G. Seroprevalence and ethnic differences in Helicobacter pylori infection among adults in the United States. J Infect Dis 2000;181:1359-63.
- Naja F, Kreiger N, Sullivan T. Helicobacter pylori infection in Ontario: prevalence and risk factors. *Can J Gastroenterol* 2007;21:501-6.
- 9. Logan RP, Walker MM. ABC of the upper

gastrointestinal tract: Epidemiology and diagnosis of Helicobacter pylori infection. *BMJ* 2001;323:920-2.

- Goh K-L, Parasakthi N. The racial cohort phenomenon: seroepidemiology of Helicobacter pylori infection in a multiracial South-East Asian country. *Eur J Gastroenterol Hepatol* 2001;13:177-83.
- Rupnow M, Shachter RD, Owens DK, Parsonnet J. A dynamic transmission model for predicting trends in Helicobacter pylori and associated diseases in the United States. *Emerg Infect Dis* 2000;6:228-37.
- 12. Parsonnet J, Shmuely H, Haggerty T. Fecal and oral shedding of Helicobacter pylori from healthy infected adults. *JAMA* 1999;282:2240-5.
- Parsonnet J. The incidence of Helicobacter pylori infection. *Aliment Pharmacol Ther* 1995;9 Suppl 2:45-51.
- Klein PD, Opekun A, Smith E, Graham D, Gaillour A. Water source as risk factor for Helicobacter pylori infection in Peruvian children. *Lancet* 1991;337:1503-6.
- Nurgalieva ZZ, Malaty HM, Graham DY, Almuchambetova R, Machmudova A, Kapsultanova D, et al. Helicobacter pylori infection in Kazakhstan: effect of water source and household hygiene. *Am J Trop Med Hyg* 2002;67:201-6.
- Hashemi MR, Rahnavardi M, Bikdeli B, Dehghani Zahedani M. H pylori infection among 1000 southern Iranian dyspeptic patients. *World J Gastroenterol* 2006;12:5479-82.
- Mansour-Ghanaei F, Yousefi Mashhour M, Joukar F, Sedigh M, Bagher-Zadeh A, Jafarshad R. Prevalence of Helicobacter pylori infection among children in Rasht, Northern Iran. *Middle East J Dig Dis* 2011;1:84-8.
- Mikaeili J, Malekzadeh R, Ziad Azb, Valizadeh M, Masarat S, Nasseri-Moghaddam S, et al. Prevalence of Helicobacter pylori in two Iranian provinces with high and low incidence of gastric carcinoma. *Arch Iran Med* 2000;3.
- Alizadeh A, Ansari S, Ranjbar M, Shalmani H, Habibi I, Firouzi M, et al. Seroprevalence of Helicobacter pylori in Nahavand: a population-based study. *East Mediterr Health J* 2009;15:129-35.
- Apan TZ, Iseri L, Aksoy A, Güliter S. The Antibody Response to Helicobacter pylori in the Sera from a Rural Population in the Central Anatolia Region of Turkey. J Health Sci 2008;54:671-4.
- Graham DY, Adam E, Reddy GT, Agarwal JP, Agarwal R, Evans Jr DJ, et al. Seroepidemiology ofHelicobacter pylori infection in India. *Dig Dis Sci* 1991;36:1084-8.

- Tkachenko MA, Zhannat NZ, Erman LV, Blashenkova EL, Isachenko SV, Isachenko OB, et al. Dramatic changes in the prevalence of Helicobacter pylori infection during childhood: a 10-year followup study in Russia. J Pediatr Gastroenterol Nutr 2007;45:428-32.
- 23. Staat MA, Kruszon-Moran D, McQuillan GM, Kaslow RA. A population-based serologic survey of Helicobacter pylori infection in children and adolescents in the United States. *J Infect Dis* 1996;174:1120-3.
- 24. Farshad S, Japoni A, Alborzi A, Zarenezhad M, Ranjbar R. Changing prevalence of Helicobacter pylori in south of Iran. *Iran J Clin Infect Dis* 2010;5:65-9.
- 25. Dube C, Tanih N, Clarke A, Mkwetshana N, Green E, Ndip R. Helicobacter pylori infection and transmission in Africa: Household hygiene and water sources are plausible factors exacerbating spread. *African J Biotechnol* 2009;8.
- 26. Shi R, Xu S, Zhang H, Ding Y, Sun G, Huang X, et al. Prevalence and risk factors for Helicobacter pylori infection in Chinese populations. *Helicobacter* 2008;13:157-65.
- Hopkins RJ, Vial PA, Ferreccio C, Ovalle J, Prado P, Sotomayor V, et al. Seroprevalence of Helicobacter pylori in Chile: vegetables may serve as one route of transmission. *J Infect Dis* 1993;168:222-6.
- Forman D, De Backer G, Elder J, Moller H, Damotta L, Roy P, et al. Epidemiology of, and risk factors for, Helicobacter pylori infection among 3194 asymptomatic subjects in 17 populations. The EUROGAST Study Group. *Gut* 1993;34:1672-6.
- Breuer T, Sudhop T, Hoch J, Sauerbruch T, Malfertheiner P. Prevalence of and risk factors for Helicobacter pylori infection in the western part of Germany. *Eur J Gastroenterol Hepatol* 1996;8:47-52.
- Hosseini E, Poursina F, Van de Wiele T, Safaei HG, Adibi P. Helicobacter pylori in Iran: A systematic review on the association of genotypes and gastroduodenal diseases. J Res Med Sci 2012;17:280-92.
- 31. Sinan A, Ozaydin AN. H.Pylori prevalence in Turkey: *A population-based cross-sectional study World Cong Of Health in Turkey Marmara University* 2009:101-2.
- Goodman KJ, Correa P, Aux HJT, DeLany JP, Collazos T. Nutritional factors and Helicobacter pylori infection in Colombian children. J Pediatr Gastroenterol Nutr 1997;25:507-15.