Demographic Features and High Prevalence of Celiac Disease in Patients with Irritable Bowel Syndrome in Khoram Abad, Lorestan

Mosayeb Moradniani¹, Zohre Mirbeik-Sabzevari^{2,*}, Asghar Aaliehpour³, Parastoo Baharvand⁴

¹Assistant Professor, Department of Internal Medicine, Lorestan University of Medical Sciences, Khorramabad, Iran

² Student Research Committee, Lorestan University of Medical Sciences, Khorramabad, Iran

- ³ Assistant Professor, Department of Pathology, Lorestan University of Medical Sciences, Khorramabad, Iran
- ⁴ Assistant Professor, Department of Community Medicine, Lorestan University of Medical sciences, Khorramabad, Iran

ABSTRACT

Background:

Many patients with irritable bowel syndrome (IBS) may have undiagnosed celiac disease (CD). Diagnosis of CD is important because early diagnosis can prevent serious complications. The purpose of this study was to investigate the demographic features and prevalence of CD in patients with IBS in Khoram Abad, Lorestan.

Materials and Methods:

This descriptive cross-sectional study was conducted on 369 patients with IBS who were diagnosed based on Rome III criteria and attended to gastrointestinal clinic of Shohada-y-Ashayer Hospital in KhorramAbad from June 2015 until March 2016. 31 patients did not accept to participate so were excluded from the study. Serological tests were performed and seropositive cases were underwent upper gastrointestinal endoscopy and duodenal biopsy. Data were analyzed using descriptive statistics, t test and Chi-square test through SPSS software version 22.

Results:

The mean age of the patients with CD was 31 ± 12 years and most of them (72.7%) were women. Most of CDs were diarrhead dominant IBS (77.3%). Among 338 patients who completed the study, 25 patients (7.4%) were seropositive, and CD was confirmed in 22 of them (6.5%) according to the Marsh classification. Marsh I was reported in seven cases, Marsh II in four, and Marsh III in 11 cases.

Conclusion:

Considering the high prevalence of CD (6.5%) in patients with IBS, the overlap of the symptoms, as well as the importance of timely diagnosis of CD, it is suggested to screen CD in patients with IBS.

Keywords: Irritable bowel syndrome, Celiac disease, Prevalence

please cite this paper as:

Moradniani M, Mirbeik-Sabzevari Z, Aaliehpour A, Baharvand P. Demographic Features and High Prevalence of Celiac Disease in Patients with Irritable Bowel Syndrome in Khoram Abad, Lorestan. *Govaresh* 2017;22:195-201.

**Corresponding author:*

Zohre Mirbeik-Sabzevari Lorestan University of Medical Sciences, Khorramabad, Iran Telefax: + 98 66 33417701 E-mail: Zm.sabzevari@yahoo.com

Received: 01 May 2017 Edited: 21 Aug. 2017 Accepted: 22 Aug. 2017

INTRODUCTION

Celiac disease (CD) is an intestinal malabsorption syndrome occurring in genetically predisposed individuals who are exposed to gluten-containing diet (1-3). CD symptoms are varied depending on the patient's age, disease duration, and the extent of the disease. In children, symptoms manifest as malnutrition and growth retardation. Adults may have no symptoms or gastrointestinal symptoms (such as diarrhea, abdominal pain, bloating, steatorrhea etc.) or non-gastrointestinal symptoms (such as weight loss, anemia, etc.) (4,5).

Diagnosis of CD is on the basis of clinical symptoms, and positive serology with duodenal pathological findings (6). Non-invasive serological tests including anti-endomysial antibody (anti-EMA, IgA) and anti-tissue transglutaminase antibody (anti-TTG, IgA) with a sensitivity of over 98% and specificity of nearly 100% are used for CD diagnosis (7). Various symptoms of the disease often lead to delays in diagnosis, which may result in serious complications (8,9).

The prevalence of CD is variable in different regions of the world (10). Previous studies have reported its prevalence as about 1% in general population in European countries as well as Iran (11,12). Studies have shown that most of the patients with CD present with symptoms suggesting irritable bowel syndrome (IBS) such as recurrent abdominal discomfort, bloating, and diarrhea, and no alarming signs (13). Therefore, the differentiation between IBS and CD (adult onset) is difficult (14-16) and these patients may be misdiagnosed as having IBS (17,18).

IBS is one of the most common diseases diagnosed by gastroenterologists (19,20). It is a chronic functional gastrointestinal disorder (21), which is characterized by abdominal pain/discomfort and changes in bowel habits and stool consistency, without organic disorder (20,22,23). There is no biochemical test, pathological, or radiological tests for diagnosis of IBS and the diagnosis is only based on the Rome III criteria, and ruling out other diseases with similar symptoms (24,25). Symptoms of CD and IBS are similar (26,27). In fact, many patients diagnosed as having IBS, are more likely to have an undiagnosed CD. The importance of diagnosis of CD is that its early detection can prevent serious complications (19,20). In general, there are contradictory studies on the prevalence of CD in the patients with IBS (28,29). In the study by Cash and colleagues, the prevalence of CD in the US was 7.3% (30). Shahbazkhani and co-workers have reported that the prevalence of CD in patients with IBS was 11.4% (31). On the other hand in the study by Emami and others in Isfahan, no relationship was observed between CD and IBS. In their study, none of the 275 patients with IBS (based on anti-TTG and IgA) had CD (9). Therefore, this study aimed to determine the demographic features and prevalence of CD in patients with IBS in Khoram Abad, Lorestan.

MATERIALS AND METHODS

This descriptive cross-sectional study was conducted in the gastrointestinal clinic of Shohada-y-Ashayer Hospital in Khorram Abad (west of Iran) from July 2015 to March 2016. IBS was diagnosed according to the Rome III criteria.

The Rome III criteria are defined as follows: abdominal pain or discomfort for at least 3 days per month in the past three months, along with two or more of the following characteristics: 1- pain relief with defecation 2- change in the frequency of defecation 3- change in stool consistency, and the onset of symptoms at least 6 months prior to the diagnosis (32).

After a complete physical examination performed by a gastroenterologist, necessary laboratory tests were done for the patients such as complete blood count (CBC), ferritin, calcium, ESR (erythrocyte sedimentation rate), stool examination (to rule out parasitic diseases especially Giardia), occult blood, TSH (thyroid-stimulating hormone), and liver and renal function tests. If the results were normal, the individuals would be enrolled in the study.

Individuals with the following criteria were excluded from the study: significant systemic disease, abdominal surgery (except for caesarean section, appendectomy, hysterectomy), inflammatory bowel disease (ulcerative colitis and Crohn's disease), pregnant and lactating women, family history of colon cancer and inflammatory bowel disease, alarming signs (anemia, blood or leukocyte in stool exam, unintentional weight loss), the onset of symptoms after 50 years of age, nocturnal diarrhea, abdominal mass, and recent use of antibiotics, metformin, or anti-depressants.

Serology

Anti-tissue transglutaminase antibody (anti-TTG, IgA) has a sensitivity of more than 98% and specificity of about 100% for the diagnosis of CD (7). Therefore, this antibody was used for initial screening. Anti-TTG was measured by ELISA method using generic assay kit with a normal upper limit of 20 u/mL.Serum IgA was determined by nephelometry and binding assay kit with a normal IgA range of 70-400 mg/dL. In the case of low serum IgA titer, anti-TTG antibody (IgG) was used by ELISA method with a normal range of less than 20 u/mL.

Prevalence of Celiac Disease in Patients with IBS

Table 1: Demographic data of all subjects							
	No (%)						
Age (mean±SD)	31 ± 9						
Male/female	173 (51.2) / 165 (48.8)						
IBS-diarrhea	232 (68.6)						
IBS-constipation & diarrhea	79 (23.4)						
IBS-constipation	27 (8)						
IBS: Irritable bowel syndrome							





Table 2: Symptoms in patients with irritable bowel syndrome

	Ν	Diarrhea	Constipation	Abdominal pain	Bloating	GERD	Early satiety	Dyspepsia	Nausea
IBS-D	232	232	51	207	97	151	78	91	59
IBS-C	27	4	27	19	22	16	19	5	12
IBS-CD	79	79	79	76	56	36	26	14	6
Total	338	315	157	302	175	203	123	110	77

Endoscopy

In the seropositive cases (anti-TTG [IgA] or anti-TTG [IgG] in IgA deficient cases), a gastroenterologist performed upper endoscopy and six biopsy specimens were taken from the second part of the duodenum (D2) and Bulb.

Pathology

Biopsy samples were studied by an experienced pathologist for diagnosis of CD, according to the newly revised Marsh classification (33).

Among the patients attended to the clinic, 338 patients with the Rome III criteria were selected for this study. This study was approved by the Ethics Committee of Lorestan University of Medical Sciences (lums.rec. 1394, 28). Patients with IBS were included in the study and a written informed consent was obtained from all the patients.

To analyze the data, descriptive statistics (mean, standard deviation, and frequency) and inferential statistics of Chi-square and t tests were used. The level of significance was considered as less than 0.05. Data were analyzed using SPSS software version 22.

RESULT

In this descriptive cross-sectional study, 369 patients with IBS according to the Rome III criteria participated in the study. 31 patients did not accept to participate and were excluded. Finally 338 patients completed the study (figure 1).

Of the 338 studied patients, 173 patients (51.2%) were male and 165 patients (48.8%) were female with the mean age of 31 ± 9 years (with a minimum age of 12 and maximum age of 58 years). 232 patients (68.6%) had diarrhea-dominant IBS. Demographic data of the subjects are shown in table 1. Diarrhea and abdominal pain were the most prevalent symptoms among the patients with IBS, respectively (table 2). 24 subjects (7.1%, 16 female and 8 male) had positive anti-TTG antibody (IgA) (20 u/mL \leq). Of the seven patients (2%) with low serum IgA titer (\leq 70mg/dL) one had positive anti-TTG antibody (IgG) (\geq 20 u/mL).

tuble of Symptoms in patients with cente discuse													
		N	diarrhea	constipation	Abdominal pain	bloating	GERD	Early satiety	dyspepsia	nausea	anemia	steatorrhea	Weight loss
Patients with celiac disease	IBS-diarrhea	17	17	4	15	9	5	9	7	2	6	5	3
	IBS-constipation	2	0	2	1	1	0	1	0	1	1	0	0
	IBS-constipation & diarrhea	3	3	3	2	1	0	0	2	1	1	2	0
Total		22	20	9	18	11	5	10	9	4	8	7	3

Table 3: Symptoms in patients with celiac disease

Table 4: Variables between patients with CD and patients without CD

		Cel			
		No	Yes	P value	
		No (%)	No (%)		
Sex	Male	167(52.8)	6(27.3)	0.02*	
	Female	149(47.2)	16(72.7)		
Age (years)	≤25	94(29.7)	10(45.5)	0.048*	
	25-40	157(49.7)	5(22.7)		
	≥40	65(20.6)	7(31.8)		
Symptoms	Diarrheal dominant	215(68)	17(77.3)	0.536	
	Constipation dominant	25(7.9)	2(9.1)		
	Intermittent Diarrhea- Constipation	76(24.1)	3(13.6)		

* Statistically significant

therefore, 25 seropositive patients underwent upper endoscopy and 22 patients (6.5%) had abnormal pathology according to the Marsh classification. Marsh I was reported in seven cases (31.8%), Marsh II in four (18.2%), and Marsh III in 11 cases(50%). Among the patients with CD, 16 patients (72.7%) were female and 6 patients (27.3%) were male with the mean age of 31 ± 12 years. Among the patients with CD, 17 patients (77.3%) had diarrhea-dominant IBS, 2 patients (9.1%) had constipation-dominant IBS, and 3 patients (13.6%) had intermittent diarrhea and constipation. The most prevalent symptom was diarrhea. Other symptoms are shown in table 3. Variables of age, sex, and clinical symptoms were compared between the two groups of patients with and without CD. No significant relationship was observed between the clinical symptoms of IBS in the patients with CD and others (p = 0.536), while sex and age were significantly different between the two groups (p = 0.02 and 0.048) (table 4). Most of the patients with CD (72.7%) were women and 45.5% of them were younger than 25 years. All the patients with the diagnosis of CD were provided with necessary education about the disease and underwent gluten-free diet. All the patients referred for follow-up and re-visit after two months. At follow-up visit, the patients presented no clinical symptoms and also, in serological tests a significant decrease of antibody titer was observed.

DISCUSSION

The prevalence of CD is estimated as 1% in general population in European studies (18). In a study conducted on the general population in Khuzestan, the prevalence of positive anti-TTG (IgA) antibody was reported to be about 1% (34) and the same results were obtained in a study in Gonbad-e Kavoos (35). Also, in a study in Kerman and Sari, the prevalence of positive anti-TTG (IgA) was reported to be 1 in 104 in the general population (36). While,the prevalence of IBS in the general population is approximately 15-25% (22). Many patients with IBS may be diagnosed as having

Prevalence of Celiac Disease in Patients with IBS

CD (19,20) and considering the similarity between the symptoms of IBS and CD, patients with CD are likely to be misdiagnosed as having IBS. So, the prevalence of CD is expected to be higher in the population suffering from IBS than in the general population.

In the present study out of the 338 patients, 25 had positive serological tests and therefore CD was confirmed in 22 (6.5%) of them according to the pathological findings. In Ahvaz, Shayesteh and colleagues in a study on 465 patients with IBS, showed that of the 47 cases with positive serology (anti-TTG or anti-gliadin or both in some cases), CD was diagnosed in 13 cases with the prevalence of 2.8% (37). In the present study, the prevalence of CD is higher, that can be due to using serum IgA in addition to anti-TTG for IgA deficiency. These methods are more sensitive in the diagnosis of the disease. So, no patient with CD was missed due to IgA deficiency.

Shahbazkhani reported that the prevalence of CD in patients with IBS in Tehran was 11.4% (31), which is higher than our findings and could have been influenced by factors such as sampling site (a referral center in Tehran), as well as different diagnostic criteria used to diagnose IBS (Rome II). In the study of Emamiand colleagues in Isfahan, none of the 275 patients with IBS had positive titer of anti-TTG (IgA), while among 60 cases of duodenal biopsy, 7 CD cases were diagnosed as March 1 to 3a (9) suggesting the lack of sensitivity of serological tests. Also, the prevalence of CD in our study was higher than the results obtained by Ahmadi and co-workers in Kerman (2.79%) (37), and Akhondi Meybodi and colleagues in Yazd (3.2%) (38). The reason for the differences in these results can be attributed to: differences in sample size, different diagnostic criteria for CD and IBS, different sensitivities of diagnostic tests, different prevalence in the general population of studies as well as the effects of genetic and environmental factors. In the study of Jafari and others in Ilam in 2013, the prevalence of CD in patients with IBS was 5.7% (39), which is close to the results of our study. In comparison with the studies performed in other countries, the prevalence of CD in this study was higher than the study done by Sanchez Vargas and colleagues in Mexico (2.5%) (40). Considering the different prevalence of CD and IBS in different regions and populations, different prevalence of IBS in patients is not unexpected. Also the prevalence of CD in patients with IBS in the present study is higher than the study of Sharma and co-workers in India that reported a prevalence of 0.8% (41). The prevalence of CD in patients with IBS in the present study was lower than a study in Bangladesh by Chowdhury (9%) (42), affected by the age, sex, and other factors related to the studied population.

In our study, most of the cases with CD had diarrhea-dominant IBS, which was consistent with the results of many studies (9,18), while other studies do not confirm our findings. For examplein Shavesteh and colleagues study, CD was more likely to be diagnosed in patients with particular clinical manifestations (diarrhea, constipation, or a combination of both) but not statistically significant (43). Most of the patients with CD were female (72.7%), which is consistent with most studies (43-45). The mean age of the patients with CD was 31±12 years, similar to what reported by Shayesteh and co-workers (33 years) (43). While in the study of Shayesteh et al, most of the cases (69%) were reported as March 1 and 2, showing the diagnosis of CD before worsening of the disease (43), which may be influenced by genetic, ethnical, and environmental factors such as the time of first exposure to gluten and the amount of wheat in individuals' diet (3,46).

CONCLUSION

Considering the high prevalence of CD (6.5%) in the patients with IBS and the overlap of the two diseases as well as the importance of timely diagnosis of CD, high-sensitivity serological tests are suggested to be performed for the screening of CD in patients with IBS.

ACKNOWLEDGEMENTS

We would like to acknowledge the Research and Technology Deputy of Lorestan University of Medical Sciences for its financial support.

CONFLICT OF INTEREST

The authors declare no conflict of interests related to this work.

REFERENCES

1. Alavinejad P, Hajiani E, Masjedizadeh R, Hashemi SJ, Faramarzi M, Sebghatollahi V, et al. Epidemiologic and demographic survey of celiac disease in khuzestan province. Middle East J Dig Dis 2014;6:98-103.

- Ciclitira PJ, King AL, Fraser JS. AGA technical review on Celiac Sprue. American Gastroenterological Association. *Gastroenterology* 2001;120:1526-40.
- Rostami Nejad M, Rostami K, Emami M, Zali M, Malekzadeh R. Epidemiology of celiac disease in iran: a review. *Middle East J Dig Dis* 2011;3:5-12.
- Rostami K, Malekzadeh R, Shahbazkhani B, Akbari M, Catassi C. Coeliac disease in Middle Eastern countries: a challenge for the evolutionary history of this complex disorder? *Dig liver Dis* 2004;36:694-7.
- 5. Green PH. The many faces of celiac disease: clinical presentation of celiac disease in the adult population. *Gastroenterology* 2005;128:74-8.
- Lu W, Gwee KA, Siah KT, Kang JY, Lee R, Ngan CC. Prevalence of Anti-deamidated Gliadin Peptide Antibodies in Asian Patients With Irritable Bowel Syndrome. J Neurogastroenterol Motil 2014;20:236-41.
- 7. Farrell RJ, Kelly CP. Celiac sprue. N Engl J Med 2002;346:180-8.
- Lebwohl B, Rubio-Tapia A, Assiri A, Newland C, Guandalini S. Diagnosis of celiac disease. *Gastrointest Endosc Clin N Am* 2012;22:661-77.
- Emami M, Kouhestani S, Gholamrezaei A, Hashemi M, Mahzouni P, Raeisi M, et al. Prevalence of celiac disease in patients with irritable bowel syndrome. *Govaresh* 2008;13:192-7.
- 10. Shamir R, Lerner A, Shinar E, Lahat N, Sobel E, Bar-or R, et al. The use of a single serological marker underestimates the prevalence of celiac disease in Israel: a study of blood donors. *Am J Gastroenterol* 2002;97:2589-94.
- 11. Feighery C. Coeliac disease. BMJ 1999;319:236-9.
- Catassi C, Ratsch IM, Fabiani E, Rossini M, Bordicchia F, Candela F, et al. Coeliac disease in the year 2000: exploring the iceberg. *Lancet* 1994;343:200-3.
- Zipser RD, Patel S, Yahya KZ, Baisch DW, Monarch E. Presentations of adult celiac disease in a nationwide patient support group. *Dig Dis Sci* 2003;48:761-4.
- Frissora CL, Koch KL. Symptom overlap and comorbidity of irritable bowel syndrome with other conditions. *Curr Gastroenterol Rep* 2005;7:264-71.
- 15. Green PHR, Stavropoulos SN, Panagi SG, Goldstein SL, McMahon DJ, Absan H, et al. Characteristics of adult celiac disease in the USA: results of a national survey. *Am*

J Gastroenterol 2001;96:126-31.

- Zobeiri M, Ebrahimi S, Farahvash M, Rezaei M, Abdollahi A. Celiac disease: Serologic prevalence in patients with irritable bowel syndrome. *J Research Med Sci* 2012;17:356-61.
- 17. Morcos A, Dinan T, Quigley EM. Irritable bowel syndrome: role of food in pathogenesis and management. *J Dig Dis* 2009;10:237-46.
- Amiriani T, Besharat S, Roshandel G, Shalizar A. Should we look for celiac disease in irritable bowel syndrome? *Oman Med J* 2011;26:59-60.
- Masoodi M, Sadeghi S, Moosavi A. Celiac disease in patients with irritable bowel syndrome. *Govaresh* 2007;12:200-4.
- 20. Spiegel BM, DeRosa VP, Gralnek IM, Wang V, Dulai GS. Testing for celiac sprue in irritable bowel syndrome with predominant diarrhea: a cost-effectiveness analysis. *Gastroenterology* 2004;126:1721-32.
- 21. Kellow JE. Introduction: a practical evidence-based approach to the diagnosis of the functional gastrointestinal disorders. *Am J Gastroenterol* 2010;105:743-6.
- 22. El-Salhy M, Hatlebakk JG, Gilja OH, Hausken T. Irritable bowel syndrome: recent developments in diagnosis, pathophysiology, and treatment. *Expert Rev Gastroenterol Hepatol* 2014;8:435-43.
- Bellini M, Gambaccini D, Stasi C, Urbano MT, Marchi S, Usai-Satta P. Irritable bowel syndrome: a disease still searching for pathogenesis, diagnosis and therapy. *World J Gastroenterol* 2014;20:8807-20.
- 24. Longstreth GF, Thompson WG, Chey WD, Houghton LA, Mearin F, Spiller RC. Functional bowel disorders. *Gastroenterology* 2006;130:1480-91.
- 25. Dalrymple J, Bullock I. Guidelines: Diagnosis and management of irritable bowel syndrome in adults in primary care: summary of NICE guidance. *BMJ* 2008;336:556-8.
- Bakhshipour A, Nezam SK, Zakeri Z, Gharibi R, Bahari A, Kaykhaei MA. Coeliac disease in irritable bowel syndrome (Rome III) in Southeast Iran. *Arab J Gastroenterol* 2012;13:24-7.
- 27. Tjon JM-L, van Bergen J, Koning F. Celiac disease: how complicated can it get? *Immunogenetics* 2010;62:641-51.
- 28. Sainsbury A, Sanders DS, Ford AC. Prevalence of irritable bowel syndrome-type symptoms in patients with celiac disease: a meta-analysis. *Clin Gastroenterol Hepatol*

Prevalence of Celiac Disease in Patients with IBS

2013;11:359-65 e1.

- Cristofori F, Fontana C, Magistà A, Capriati T, Indrio F, Castellaneta S, et al. Increased prevalence of celiac disease among pediatric patients with irritable bowel syndrome: a 6-year prospective cohort study. *JAMA Pediatr* 2014;168:555-60.
- Cash BD, Rubenstein JH, Young PE, Gentry A, Nojkov B, Lee D, et al. The prevalence of celiac disease among patients with nonconstipated irritable bowel syndrome is similar to controls. *Gastroenterology* 2011;141:1187-93.
- Shahbazkhani B, Forootan M, Merat S, Akbari MR, Nasserimoghadam S, Vahedi H, et al. Coeliac disease presenting with symptoms of irritable bowel syndrome. *Aliment Pharmacol Ther* 2003;18:231-5.
- Drossman D, Corazziari E, Delvaux M, Spiller R, Talley N, Thompson W, et al. Rome III diagnostic criteria for functional gastrointestinal disorders. *Func Gastrointestin Disord* 2006;12:885-915.
- Marsh MN, Johnson MW, Rostami K. Mucosal histopathology in coeliac disease: a rebuttal of Oberhuber's sub-division of Marsh III. *Gastroenterol Hepatol Bed Bench* 2015;8:50-5.
- Masjedizadeh R, Hajiani E, Hashemi J, Shayesteh AA, Moula K, Rajabi T. Celiac disease in South-West of Iran. *World J Gastroenterol* 2006;12:4416-9.
- Khoshnia M, Pourshams A, Mohammadkhani A, Tavangar S, Shahbazkhani B, Malekzadeh R. Celiac disease in gonbad-kavoos. *Govaresh* 2012;10:131-3.
- 36. Akbari MR, Mohammadkhani A, Fakheri H, Javad Zahedi M, Shahbazkhani B, Nouraie M, et al. Screening of the adult population in Iran for coeliac disease: comparison of the tissue-transglutaminase antibody and anti-endomysial antibody tests. *Eur J Gastroenterol Hepatol* 2006;18:1181-6.
- Ahmadi B, Zaherara M. Prevalence of Celiac Disease in Patients with Irritable Bowel Syndrome in Kerman, Iran. J Kerman Univ Med Sci 2015;22:319-27.
- Akhondi Meybodi M, Rabei A, Salehi S. Frequency of Celiac Disease in Irritable Bowel Syndrome Patients with Predominant Diarrhea Referred to Gastroenterology Clinics in Yazd, Iran. J Shahid Sadoughi Univ Med Sci Health Services 2012;12:64-8.
- 39. Jafarihaydarlo A, Mahmoodi A, Yasemi M, Baziar A, Ehsanbakhsh S, Rashidbeygi M. Frequency of Celiac Disease in Patients with Diarrhea Predominant Irritable Bowel Syndrome Referred to Gastroenterology Clinics in Ilam During 2008to 2012. J Ilam Univ Med Sci

2013;21:191-8.

- 40. Sanchez-Vargas LA, Thomas-Dupont P, Torres-Aguilera M, Azamar-Jacome AA, Ramirez-Ceervanes KL, Aedo-Garces MR, et al. Prevalence of celiac disease and related antibodies in patients diagnosed with irritable bowel syndrome according to the Rome III criteria. A case-control study. *Neurogastroenterol Motil* 2016;28:994-1000.
- Sharma H, Verma AK, Das P, Dattagupta S, Ahuja V, Makharia GK. Prevalence of celiac disease in Indian patients with irritable bowel syndrome and uninvestigated dyspepsia. *J Dig Dis* 2015;16:443-8.
- 42. Chowdhury MK, Chakraborty R, Gope S, Rahman MA, Miah AR, Raihan AS, et al. Celiac Disease in Patients Fulfilling the Rome III Criteria for Irritable Bowel Syndrome Attending Gastroenterology Department of A Tertiary Care Hospital in Bangladesh. *Mymensingh Med* J 2016;25:102-8.
- 43. Shayesteh AA, Hajiani E, Hashemi SJ, Masjedizadeh A, Latifi SM, Shayesteh M. Prevalence of celiac disease in Iranian patients with irritable bowel syndrome: a crosssectional study. *J Dig Dis* 2014;15:12-7.
- 44. Yap TW, Chan WK, Leow AH, Azmi AN, Loke MF, Vadivelu J, et al. Prevalence of serum celiac antibodies in a multiracial Asian population--a first study in the young Asian adult population of Malaysia. *PLoS One* 2015;10:e0121908.
- 45. Gomez JC, Selvaggio GS, Viola M, Pizarro B, la Motta G, de Barrio S, et al. Prevalence of celiac disease in Argentina: screening of an adult population in the La Plata area. *Am J Gastroenterol* 2001;96:2700-4.
- 46. Norris JM, Barriga K, Hoffenberg EJ, Taki I, Miao D, Haas JE, et al. Risk of celiac disease autoimmunity and timing of gluten introduction in the diet of infants at increased risk of disease. *JAMA* 2005;293:2343-51.