

# *In Vitro* Lipid Peroxidation and Antimicrobial Activity of *Mucuna pruriens* Seeds

YERRA RAJESHWAR, MALAYA GUPTA and UPAL KANTI MAZUMDER

Division of Pharmaceutical Chemistry and Pharmacology, Department of Pharmaceutical Technology, Jadavpur University, Kolkata, India.

Received May 5, 2005; Revised May 25, 2005; Accepted June 2, 2005

This paper is available online at <http://ijpt.iuims.ac.ir>

## ABSTRACT

The present investigation is aimed to carry out the *in vitro* lipid peroxidation and antimicrobial activities of the methanol extract of *Mucuna pruriens* (MEMP) (Family: Fabaceae) seeds. Lipid peroxidation was monitored by the change in optical density of the prepared concentrations (10-320 µg/ml) and the % inhibition was calculated. Ascorbate/FeSO<sub>4</sub>-induced peroxidation was inhibited by standard antioxidants such as quercetin, L-ascorbic acid and MEMP. Moreover, the % inhibition of the methanol extract increased with increase in concentration. IC<sub>50</sub> value for the MEMP, L-ascorbic acid and quercetin for lipid peroxidation was found to be 217.25 µg/ml, 41 µg/ml and 19.75 µg/ml respectively. The antimicrobial activity of MEMP was determined by disc diffusion method with various Gram positive and Gram-negative microorganisms. MEMP showed broad-spectrum antimicrobial activity against all the tested microorganisms except *Staphylococcus aureus* ML 152 and *Vibrae cholera* 14035. The results obtained in the present study indicate that MEMP can be a potential source of natural antioxidant and antimicrobial agent.

**Keywords:** *Mucuna pruriens* seeds, *In vitro* lipid peroxidation, Antimicrobial activity

There has been growing interest in the investigation of the natural products from plants for the discovery of new antimicrobial and antioxidant agents as well as an alternative route for the substitution of synthetic chemicals, side effects of which are always in question. For this, the essential oils and the extracts of many plants have been prepared and screened for their antimicrobial and antioxidant activities leading to the accumulation of a large number of reports in the literature concerning the above mentioned properties of plants [1-5]. Because of the side effects and the resistance that pathogenic microorganisms build against antibiotics, much recent attention has been paid to extracts and biologically active compounds isolated from plant species used in herbal medicine [6]. Plant based antimicrobials represent a vast untapped source for medicines and further exploration of plant antimicrobials needs to occur. Antimicrobials of plant origin have enormous therapeutic potential. They are effective in the treatment of infectious diseases while simultaneously mitigating many of the side effects that are often associated with synthetic antimicrobials [7].

*Mucuna pruriens* is a twinning herb found all over tropical parts of India. It, an Indian indigenous leguminous plant, is well known for producing itch. This property is attributed to the trichomes (hair) present on the

Pods. It has been established that this unique property is accounted by the presence of 5-hydroxy tryptamine (5-HT) in the hair. [8]. It is also likely that histamine and kinin like substances may also be responsible [9]. Some reports show that anti-histaminics afford protection against the itch [9]. It has been reported to be anti-diabetic [10]. *Mucuna pruriens* seeds are herbaceous forage and food legumes that have for a long time found widespread usage as rotation crops for management of various pests and pathogens, as well as in soil improvement and weed control [13, 14]. Seeds of *Mucuna pruriens* are known to produce the unusual non-protein amino acid 3-(3,4-dihydroxyphenyl)-L-alanine (L-DOPA), a potent neurotransmitter precursor that is, at least in part, believed to be responsible for the toxicity of *Mucuna* seed [15]. L-DOPA, a potentially neurotoxic agent used in the treatment of Parkinson's Disease, is found in relatively large amounts in *Mucuna pruriens* seeds [16, 17] to the point where the seeds have been suggested as a medical source of L-DOPA [16] and even in the treatment of Parkinson's Disease [18].

Previously, we reported the antiepileptic and anti-neoplastic activity of methanol extract of *Mucuna pruriens* root [19] from our laboratory. Our recent findings revealed that the methanol extract of *Mucuna pruriens* seeds showed significant *in vitro* antioxidant activity

[20]. In this work, we have tested the *in vitro* lipid peroxidation and antimicrobial activity (against Gram positive and Gram negative bacteria) of the methanol extract of *Mucuna pruriens*.

## MATERIALS AND METHODS

### Chemicals

Quercetin, L-ascorbic acid and TBA were purchased from Sigma Chemical Co. (St. Louis, MO, USA). All other chemicals and reagent used were of analytical grade.

### Plant Extract

The seeds of *Mucuna pruriens* (MP) were purchased from the United Chemicals and Allied Products, Kolkata, India. They were identified by the Botanical Survey of India (BSI), Kolkata, India. For the extract, the seeds were dried in shade and powdered in a mechanical grinder. The powder of MP seeds was initially defatted with petroleum benzene (60-80°C) followed by 1000 ml of methanol by using a Soxhlet extractor for 72 h at a temperature not exceeding the boiling point of the solvent [21]. The extract was filtered using Whatman filter paper (No. 1) and then concentrated in vacuum and dried. The extract thus obtained was directly used in the assay of lipid peroxidation and antimicrobial activity.

Previously isolated classes of compounds

The phytochemical study revealed that the methanol extract of *Mucuna pruriens* (MEMP) seeds contained alkaloids, flavones, saponins, aminoacids, and fatty acids [22].

### Bacterial Strains Employed

Microorganisms (*Staphylococcus aureus* 8531, *Staphylococcus aureus* ML 174, *Staphylococcus aureus* ML 152, *Bacillus pumillus* 8241, *Bacillus cereus*, *Escherichia coli* 51, *Escherichia coli* 54B, *Vibrea cholera* 14035, *Vibrea cholera* 1353, and *Vibrea cholera* 226101) were obtained from the stock culture of Central Drugs Laboratory, Kolkata; Indian Institute of Chemical Biology, Kolkata and Mycology and Plant Pathology Laboratory, Calcutta University, Kolkata, India.

### Lipid Peroxidation

Lipid peroxidation induced by  $Fe^{2+}$ -ascorbate system in rat liver homogenate by the method of Bishayee and Balasubramaniyam [23] was estimated as thiobarbituric acid reacting substances (TBARS) by the method of Ohkawa *et al.* [24]. The reaction mixture contained rat liver homogenate 0.1 ml (25% w/v) in Tris-HCl buffer (20 mM, pH 7.0); KCl (30 mM);  $FeSO_4(NH_4)_2SO_4 \cdot 7H_2O$  (0.06 mM); and various concentrations of *Mucuna pruriens* extract in a final volume of 0.5ml. The reaction mixture was incubated at 37°C for 1 h. After the incubation period, 0.4ml was removed and treated with 0.2ml sodium dodecyl sulphate (SDS) (8.1%); 1.5 ml thiobarbituric acid (TBA) (0.8%); and 1.5 ml acetic acid (20%, pH 3.5). The total volume was made up to 4.0 ml with distilled water and then kept in a




Fig 1. Effect of methanol extract of *Mucuna pruriens* on lipid peroxidation.

water bath at 95 to 100°C for 1 h. After cooling, 1.0 ml of distilled water and 5.0 ml of n-butanol and pyridine mixture (15:1 v/v) were added to the reaction mixture, shaken vigorously and centrifuged at 4000 rpm for 10 min. The butanol-pyridine layer was removed and its absorbance at 532 nm was measured to quantify TBARS. Inhibition of lipid peroxidation was determined by comparing the optical density (OD) of treatments with that of the control. Quercetin and L-ascorbic acid were used as standard.

### Determination of Antimicrobial Activity

Antimicrobial activity was measured using the standard method of diffusion disc plates on agar [25]. 0.1 ml of each culture of bacteria was spread on agar plate surfaces. For antibacterial assays, all bacterial strains were grown in Mueller Hinton Broth medium (Merck) for 24 h at 37°C. The concentration of bacterial suspensions was adjusted to  $10^8$  colony forming units ( $10^8$  cfu/ml) in Mueller Hinton Agar. Paper discs (6 mm in diameter) were impregnated on the agar to load 10 µl of each sample. The impregnated disks were placed on the medium suitably spaced apart and the plates were incubated at 5°C for 1 h to permit good diffusion and then transferred to an incubator at 37°C for 24 h. The results were recorded by measuring the zones of growth inhibition surrounding the disc. Clear inhibition zones around the discs indicated the presence of antimicrobial activity. All data on antimicrobial activity are the average of triplicate analyses. In order to determine the antibacterial effect of the MEMP, chloramphenicol (10 µg/ml/disc) were used as positive control. Inhibition diameters were measured after incubation for 24 h at 37°C.

## RESULTS

### Effect of MEMP on Lipid Peroxidation

The effect of MEMP and commercially available antioxidants namely quercetin and L-ascorbic acid on the *in vitro* inhibition of lipid peroxidation is shown in Fig 1. The generation of lipid peroxidase by  $Fe^{2+}$ -ascorbate

**Table 1.** Effect of MEMP on selected antimicrobial species.

Microorganism	10% DMSO/ml/disc	MEMP		Chloramphenicol (10 µg/ml/disc)
		500 µg/ml/disc	750 µg/ml/disc	
<i>Staphylococcus aureus</i> 8531	6	8	13	22
<i>Staphylococcus aureus</i> ML 174	6	6	10	21
<i>Staphylococcus aureus</i> ML 152	6	6	8	18
<i>Bacillus pumillus</i> 8241	7	14	17	22
<i>Bacillus cereus</i>	6	6	11	13
<i>Escherichia coli</i> 51	7	6	13	18
<i>Escherichia coli</i> 54B	6	11	18	22
<i>Vibrea cholera</i> 14035	6	7	9	23
<i>Vibrea cholera</i> 1353	6	15	19	15
<i>Vibrea cholera</i> 226101	6	19	23	26

All the values were the mean of three experiments.

The values given are the diameter of zone of inhibition (mm) including disk diameter of 6 mm.

in rat liver homogenate seems to be inhibited by MEMP with IC<sub>50</sub> value of 217.25 µg/ml. A similar effect was produced by L-ascorbic acid (IC<sub>50</sub>=41 µg/ml) and quercetin (IC<sub>50</sub>=19.75 µg/ml), indicating that the effect of MEMP on the inhibition of lipid peroxide production is significant ( $p < 0.05$ ). The inhibition percentage of lipid peroxidation in the presence of extract was found to be 61.41% of the corresponding controls. The values for L-ascorbic acid and quercetin were found to be 72.11% and 84.09%, respectively at 320 µg/ml.

#### Effect of MEMP on Antimicrobial Activity

The data presented in Table 1 indicate that the methanol extract of *Mucuna pruriens* (MEMP) inhibit the growth of some of the tested microorganisms (Gram positive and Gram negative) to various degrees. The MEMP at a concentration of 500 µg/ml and 750 µg/ml exhibited significant ( $p < 0.05$ ) antimicrobial effect against all the tested microorganisms. The extract showed strong antibacterial activity against *Bacillus pumillus* 8241, *Escherichia. Coli* 5B and *Vibrae Cholera* 1353 and 226101. However, their activity against *Staphylococcus aureus* ML 152 and *Vibrae cholera* 14035 was found to be significantly ( $p > 0.05$ ) less than the control. The antimicrobial activity was compared with the standard Chloramphenicol at a concentration of 10 µg/ml.

#### Statistical Analysis

All treatments were performed in triplicate and each data point in the results is the mean of two or three replicate tests. All experiments were repeated at least once. The statistical significance of a treatment effect was evaluated by student's *t*-test and the values were expressed as mean ± SEM. Probability limit was set at  $p < 0.05$ .

#### Discussion

Unsaturated lipids in liver tissue are very susceptible to peroxidation when they are exposed to reactive oxygen species (ROS). In the present investigation we have incubated the liver tissue in presence of a ROS generating system, ascorbate/FeSO<sub>4</sub>, and examined the effect on tissue homogenate by measuring the optical density (OD) at 532 nm. The results of the investigations revealed that MEMP had potent lipid peroxidation inhibition activity.

The antimicrobial activity of the MEMP was studied by the disc diffusion method against various microorganisms. Disc diffusion methods are used extensively to investigate the antibacterial activity of natural substances and plant extracts. These assays are based on the use of discs as reservoirs containing solutions of the substances to be examined. In the case of solutions with a low activity, however, a large concentration or volume is needed. Because of the limited capacity of discs, holes or cylinders are preferably used [26]. MEMP showed a broad spectrum of activity against all the bacterial strains as shown in Table 1. Chloramphenicol (10 µg/ml/disc) was used as a positive control.

On the basis of the results obtained in the present study, we conclude that the methanol extract of *Mucuna pruriens* had significant *in vitro* lipid peroxidation and antimicrobial activity. The components responsible for the inhibition of lipid peroxidation of MEMP are currently unclear. Further studies are needed to isolate the active components, responsible for the lipid peroxidation and antimicrobial activities.

#### ACKNOWLEDGMENTS

One of the authors, Yerra Rajeshwar is grateful to All India Council for Technical Education and Training (AICTE), New Delhi, India, for providing financial support for this investigation.

#### REFERENCES

1. Dapkevicius A, Venskutonis R, van Beek TA, Linssen JPH. Antioxidant activity of extracts obtained by different isolation procedures from some aromatic herbs grown in Lithuania. *J Sci Food Agric* 1998;**77**:140–6.
2. Sokmen A, Jones BM, Erturk M. The *in vitro* antibacterial activities of Turkish medicinal plants. *J Ethnopharmacol* 1999;**67**:79–86.
3. Dorman HJD, Deans SG. Antimicrobial agents from plants: antibacterial activity of plant volatile oils. *J Appl Microbiol* 2000;**88**:308–16.
4. Jantova S, Nagy L, Ruzekova L, Grancai D. Antibacterial activity of plant extracts from the families Fabaceae, Olaceae, Philadelphaceae, Rosaceae and Staphyleaceae. *Phyther Res* 2000;**14**:601–3.
5. Dang MN, Takacsova M, Nguyen DV, Kristianova K. Antioxidant activity of essential oils from various species. *Nahrung* 2001;**45**:64–6.

6. Essawi T, Srour M. Screening of some Palestinian medicinal plants for antibacterial activity. *J Ethnopharmacol* 2000;**70**:343-9.
7. Iwu MW, Duncan AR, Okunji CO. New antimicrobials of plant origin. In: Janick, J. (Ed.), Perspectives on New Crops and New Uses. ASHS Press, Alexandria, VA. 1999; 457-62.
8. Armstrong D' Arcy RML, Keela CA Maikhana M. Observations on chemical excitants of cutaneous pain in man. *J Physiology* 1953;**120**:326-51.
9. Broadbent JL. Observation on itching produced by Cowhage and on the part played by histamine as a mediator of the itch sensation *Brit J Pharmacol Chemother* 1953;**8**:263-70.
10. Dhawan BN, Dubey MP, Mehrotra BN, Rastogi RP, Tandon JS. Screening of Indian plants for biological activity. Part 9. *Ind J Expt Biol* 1980;**18**:594-06.
11. Aguiyi JC, Uguro MO, Gesa AA. Mechanism of action of the aqueous seed extract of *Mucuna pruriens* on the guinea pig ileum. *Phytother Res* 1997;**11**:328-9.
12. Guerranti R, Aguiyi JC, Leoncini R, Pagani R, Cinci G, Marinello E. Characterization of the factor responsible for the anti-snake activity of *Mucuna pruriens* seeds. *Journal of Preventive Medicine and Hygiene* 1999;**1(40)**:25-8.
13. Buckles D. Velvet bean: a new plant with a history. *Economic Botany* 1995;**49**:13-25.
14. Duke JA. Handbook of legumes of world economic importance. New York: Plenum Press 1981.
15. Lorenzetti E, Maclsaac S, Arnason JT, Awang DVC, Buckles D. The phytochemistry, toxicology and food potential of velvet bean (*Mucuna Adans* spp., Fabaceae). In D. Buckles, O. Osiname, M. Galiba, & G. Galiano, Cover crops of West Africa: contributing to sustainable agriculture. IDRC, Ottawa, Canada & IITA, Ibadan, Nigeria. 1998; pp. 57.
16. Bell EA, Janzen DH. Medical and ecological considerations of L-DOPA and 5-HT in seeds. *Nature* 1971;**229**:136-7.
17. Daxenbichler ME, VanEtten CH, Hallinan EA, Earle FR, Barclay AS. Seeds as sources of L-DOPA. *J Med Chem* 1971;**14**:463-5.
18. Hussain G, Manyam BV. *Mucuna pruriens* proves more effective than L-DOPA in Parkinson's Disease animal model. *Phytother Res* 1997;**11**:419-23.
19. Gupta M, Mazumder UK, Chakraborti S, Rath N, Bhawal SR. Antiepileptic and anticancer activity of some indigenous plants. *Indian J Physiol Allied Sci* 1997;**51(2)**:53-6.
20. Rajeshwar Y, Senthil Kumar GP, Gupta M, Mazumder UK. Studies on *in vitro* antioxidant activities of methanol extract of *Mucuna pruriens* (Fabaceae) seeds. *European Bulletin of Drug Research* 2005;**13(1)**:31-9.
21. Lin J, Opoku AR, Geheeb-Keller M, Hutchings AD, Terblanche SE, Jager AK, van Staden J. Preliminary screening of some traditional Zulu medicinal plants for anti-inflammatory and antimicrobial activities. *J Ethnopharmacol* 1999;**68**:267-74.
22. Ghosal S, Singh S, Bhattacharya SK. "Alkaloids of *Mucuna pruriens*. Chemistry and pharmacology." *Planta Medica* 1971;**19**:279.
23. Bishayee S, Balasubramaniam AS. *J Neurochem* 1971;**18**:909-20.
24. Ohkawa H, Onishi N, Yagi K. Assay of lipid peroxidation in animal tissue by thiobarbituric acid reaction. *Anal Biochem* 1979;**95**:351-8.
25. Abou-Zeid, Abou-Zeid A, Shehata, Youssef. *Indian J Pharmacol* 1969;**31(3)**:72.
26. Bartner A, Pfeiffer KP & Batner H. Applicability of disc diffusion methods required by the pharmacopoeias for testing antibacterial activity of natural compounds. *Pharmazie* 1994;**49**:512-6.

---

**Address correspondence to:** Prof. Upal Kanti Mazumder, Department of Pharmaceutical Technology, Jadavpur University, Kolkata, India. E-mail: [vrajeshwar@yahoo.co.in](mailto:vrajeshwar@yahoo.co.in)

---