

1 RESEARCH ARTICLE

2 Antibacterial Activity and Cytotoxicity Screening of 3 Sumatran Kaduk (*Piper sarmentosum* Roxb.)

4 E. ATIAX, F. AHMAD, H. M. SIRAT and D. ARBAIN

5 For author affiliations, see end of text.

6 Received December 15, 2009; Revised July 3, 2010; Accepted September 28, 2010

7 This paper is available online at <http://ijpt.iums.ac.ir>8 **ABSTRACT**

9 Phytochemical investigations of *Piper sarmentosum* Roxb., yielded four compounds; three amides,
10 identified as 3-(3',4',5'-trimethoxyphenylpropanoyl) pyrrolidine, 3-(4'-methoxyphenylpropanoyl) pyrrole, *N*-
11 (3-phenylpropanoyl) pyrrole and a sterol namely β -sitosterol. 3-(4'-Methoxyphenylpropanoyl) pyrrole was
12 found for the first time in this *Piper* species. All chemical constituents were tested for their antibacterial
13 activity using disk diffusion method and cytotoxicity screening using sul-forhodamine B (SRB) assay. All
14 of the compounds were found only active towards gram-positive bacteria except 3-(4'-
15 methoxyphenylpropanoyl) pyrrole with no activity against both gram-positive and gram-negative bacteria.
16 Meanwhile, the cytotoxicity screening using SRB assay indicated that none of these compounds was
17 active as an anticancer agent.

18 **Keywords:** *Piperaceae*, *P. sarmentosum*, *Amides*, *Antibacterial*, *Cytotoxicity*

19 The study of medicinal plants opened the door to the 47 them for a long time in a variety of medicinal capacities
20 development of purified and defined chemical 48 [5].
21 compounds as dose-controlled medicines. Natural 49 *Piper sarmentosum* Roxb. is one of the *Piper* genus
22 compounds can become central players in the treatment 50 which is known as kaduk, sirih duduk or mengkadak in
23 of disease and in the understanding of disease 51 Indonesia. Traditionally, it was used as a remedy for
24 mechanisms. Compounds that emerged from the study 52 tooth-ache and for fungoid dermatitis on the feet, for
25 of ethnobotanic extracts became important as medicines 53 treatment of coughs, influenza and rheumatism [6]. A
26 and were enabling as pharmacologic tools in the 54 decoction of the leaves is drunk to treat malarial fever
27 elucidation of disease mechanisms [1]. Piperaceae 55 [7] and the crushed leaves are mixed with water and
28 family has provided many past and present civilizations 56 used for bathing to treat kidney stones and difficulty in
29 with a source of diverse medicines and food grade spice 57 urination [8]. Previous chemical constituents on this
30 [2]. This plant is distributed pantropically. The earliest 58 plant have resulted in the isolation of a number of
31 classification of the Piperaceae family recognized 59 compounds [9-13]. We now describe the isolation of an
32 between 7 to 15 genera and five of them such as *Piper*, 60 additional amide from the aerial part of this plant which
33 *Peperomia*, *Lepianthes*, *Macropiper*, and *Trianopiper* 61 was collected from Sumatra, Indonesia and also its
34 are only accepted as the principle genera of Piperaceae. 62 antibacterial and cytotoxicity activities.

35 This genus contains over 1000 species in the world [3].
36 This plant can be recognized by three main features: 63
37 articulate stem, asymmetrical or cordate leaves, and
38 axillary spikes of little round berry-like fruits [4].
39 According to Jaramilo [3], Asian tropic has 340 species
40 of *Piper*, including Sumatra tropical rainforests. This
41 species takes the form of shrubs, herbs, lianas, and
42 mostly woody climbers. They are common in the warm,
43 humid region and in the lowland of wet forests. The
44 leaves are typically aromatic or have pungent smell.
45 This genus consists of a large family of plants
46 indigenous to the tropic and native people have used

64 **MATERIALS AND METHODS**65 *General Experimental Procedures*

66 Mps. (uncorr.) were determined using the Leica
67 Gallen III apparatus. IR spectra were recorded on a
68 Perkin-Elmer 1650 FTIR spectrophotometer. NMR
69 spectra were recorded on a Bruker Avance 300
70 Spectrometer, ^1H NMR spectra were measured at 300
71 MHz and ^{13}C NMR spectra were measured at 75 MHz.
72 Deuterated solvent of chloroform (CDCl_3) was used as




Fig 1. Chemical structure of the isolated compounds of *P. sarmentosum*

72 solvents. Mass spectra data were obtained from Kent 108 concentrated and further purified by recrystallization
73 Mass Spectrometry Services, United Kingdom. CC: 109 from hexane to yield 3-(3',4',5'-trimethoxy-
74 silica gel (Merck 70-230 mesh and 230-400 mesh). 110 phenylpropanoyl)pyrrolidine (Fig 1- Compound 2)
75 Spots on TLC were visualized by UV (254 and 365 nm) 111 (135.6 mg, $8.5 \times 10^{-3}\%$) as colourless crystalline solids
76 and vanillin-sulphuric acid reagent. Streptomycin 112 with mp 158-159°C (lit. [15] 156-157°C).
77 sulphate standard was purchased from Oxoid 113. The crude EtOAc extract (10 g) was fractionated by
78 (Hampshire, UK). 114 VLC using hexane, mixture of hexane-CH₂Cl₂ and
115 CH₂Cl₂ by step gradient polarity technique to yield
116 seven fractions (PSEA-PSE G). Fraction PSEB (1.033

79 Plant Material

80 The aerial parts of *P. sarmentosum* were collected 117 g) was purified by CC over SiO₂ (70-230 mesh, 75 g)
81 from Desa Sariak, Sungai Pua, about 11 km from 118 with hexane-CH₂Cl₂ (50: 50) as eluent to give four
82 Bukittinggi, West Sumatra, Indonesia in 2005. The 119 fractions. Fractions 25-124 were combined and
83 sample (EM-01/1205) was identified by Mr. Rusdi 120 concentrated to yield 3-(4'-methoxyphenylpropanoyl)
84 Tamin and Ms. Nurainas and specimen was deposited at 121 pyrrole (Fig 1- Compound 3) (18.9 mg, 0.0012%) as
85 the Andalas Herbarium (ANDA), University of 122 yellow crystalline solids with mp 83-84°C (lit. [16] 86-
86 Andalas, Padang, Indonesia. 123 87°C).

87 Extraction and Isolation

88 The powdered of aerial plant parts of *P.* 126 (10:90) as eluent to afford 102 fractions (PSED1-PSE
89 *sarmentosum* (1.6 kg) was soxhlet-extracted 127 D9). Fraction PSE D4 (83.6 mg) was further
90 successively with 3.5 L of each hexane and ethyl acetate 128 rechromatographed by CC over SiO₂ with CH₂Cl₂ as
91 for 18 hours. The solvent of each extract was evaporated 129 eluent to yield N-(3-phenylpropanoyl) pyrrole (Fig 1-
92 *in vacuo* to afford the crude hexane, PSH (28.9 g, 1.81 130 Compound 4) (34.5 mg, 0.0022%) as colourless liquid.

93%) and ethyl acetate, PSE (24.9 g, 1.55 %). The crude 131 **Antibacterial Assay (disk diffusion method)**

94 PSH extract (10 g) was fractionated by VLC over silica 132 The chemical constituents from *P. sarmentosum*
95 gel (230-400 mesh, 250 g) and eluted with gradient 133 were tested against gram-negative; *Escherichia coli*,
96 solvent system of hexane, hexane- CH₂Cl₂ and CH₂Cl₂ 134 *Pseudomonas aeruginosa* and gram positive bacteria;
97 to afford 10 fractions (PSHA-PSHJ). 135 *Bacillus subtilis*, and *Staphylococcus aureus*. Agar
98 Fraction PSHD and PSHF were combined (421.5 136 cultures of the test microorganisms were prepared
99 mg) and subjected to CC over SiO₂ (70-230 mesh, 40 g) 137 according to Mackeen *et al.* [17]. Samples were
100 to yield nine fractions. The seventh fraction was 138 dissolved in MeOH (1 mL). The test samples (10 µL)
101 concentrated and recrystallized from CH₂Cl₂-hexane to 139 were loaded onto each Whatman filter paper disks (0.6
102 give *β*-sitosterol (Fig 1- Compound 1) (95.1 mg, 140 mm) and evenly placed on the agar surface previously
103 0.0059%) as white crystalline needles with melting 141 inoculated with the suspensions of microorganism to be
104 point (mp) 133-134°C (lit. [14] 138-139°C). 142 tested. Standard disk of streptomycin sulphate (10
105 Fraction PSH I (248.5 mg) was chromatographed 143 µg/disk) was used as the positive control and DMSO
106 over SiO₂ (30 g) CC with CH₂Cl₂ (100%) as eluent to 144 was used as the negative control. The plates were
107 give 156 fractions. The combined fractions 17-39 was

Table 1. Antibacterial activity of the isolated compounds of *P. sarmentosum*

Compounds	Zone of Inhibition (mm)			
	Gram-Positive Bacteria		Gram-Negative Bacteria	
	<i>B. s</i>	<i>S. a</i>	<i>P. a</i>	<i>E. c</i>
(1)	9.7 ± 0.52	10.3 ± 0.82	-	-
(2)	12.3 ± 0.52	-	-	-
(3)	-	-	-	-
(4)	-	10.2 ± 0.41	-	-
SS	19.3 ± 1.21	21.3 ± 0.81	19.1 ± 0.54	18.8 ± 0.74

Data represent mean ± standard deviation of three independent experiments performed in duplicate. (-): no activity; *B.s*: *Bacillus subtilis*; *S.a*: *Staphylococcus aureus*; *E.c*: *Escherichia coli*; *P.a*: *Pseudomonas aeruginosa*.

Table 2. MIC and MBC value of the isolated compounds of *P. sarmentosum*

Compounds	MIC (µg/mL)				MBC (µg/mL)			
	Gram-Positive bacteria		Gram-Negative Bacteria		Gram-Positive Bacteria		Gram-Negative Bacteria	
	<i>B. s</i>	<i>S. a</i>	<i>P. a</i>	<i>E. c</i>	<i>B. s</i>	<i>S. a</i>	<i>P. a</i>	<i>E. c</i>
(1)	-	500	-	-	-	500	-	-
(2)	500	-	-	-	1000	-	-	-
(3)	-	-	-	-	-	-	-	-
(4)	-	125	-	-	-	125	-	-
SS	3.91	3.91	3.91	3.91	7.81	7.81	7.81	7.81

MIC: Minimal inhibitory concentration, MBC: minimal bacterial concentration, *B.*: *Bacillus subtilis*; *S.a*: *Staphylococcus aureus*; *E.c*: *Escherichia coli*; *P.a*: *Pseudomonas aeruginosa*.

145 inverted and incubated for 18 hours at 37°C. Clear
146 inhibition zones around the discs indicated the presence
147 of antimicrobial activity.

148 The positive results then followed by the
149 determination of Minimum Inhibitory Concentration
150 (MIC) by the micro-titer broth dilution method [18].
151 This test was performed in a sterile 96-well micro titer
152 plates. The test samples (1 mg) were dissolved in
153 methanol to obtain 1000 µg/mL stock solution. Each
154 methanolic stock samples (10 µL) was transferred to
155 micro titer plate well in duplicate at row A. A number of
156 wells were reserved in each plate for positive and
157 negative controls. Sterile broth (100 µL) was added to
158 each micro-titer plate well from row B to row H. Then,
159 the suspensions of microorganisms (200 µL) were added
160 to the samples at row A. Mixture from row A (100 µL)
161 was transferred to each micro titer plate well in order to
162 obtain a twofold serial dilution of stock samples
163 (concentration of 500 µg/mL to 3.9 µg/mL) plates were
164 then incubated for 18 hours at 37°C. Bacterial growth
165 was indicated by the presence of turbidity and a pellet at
166 the bottom of the well. The lowest concentrations,
167 which did not show any growth of tested
168 microorganisms after macroscopic evaluation were
169 determined as MIC values.

170 The MIC values were confirmed by the
171 determination of Minimal Bactericidal Concentration
172 (MBC) values according to method developed by Arias
173 *et al.* [19]. All wells in the MIC study, which did not
174 show any growth of bacteria after incubation period
175 were first diluted in fresh nutrient broth (1:4) and then
176 sub-cultured onto the surface of freshly prepared
177 nutrient agar plates (Ø,15 mm). The plates were
178 incubated for 18 hours at 37°C. The MBC were
179 recorded as the lowest concentration of the sample that
180 did not permit any visible bacteria colony growth on the
181 appropriate agar plate after the incubation period.

182 Cytotoxicity Screening

183 The cytotoxicity screening was carried out according
184 to sul-forhodamine B (SRB) method described by
185 Houghton *et al.* [20]. This method relies on the uptake
186 of the negatively charge pink aminoxanthine dye,
187 sulphorhodamine B by basic amino acids in the cells.
188 The greater the number of cells, the greater the amount
189 of dye is taken up and, after fixing, when the cells are
190 lysed, the released dye will give a more intense colour
191 and greater absorbance.

192 The screening of cytotoxicity test for isolated
193 compounds from this *Piper* species against four
194 cancerous cell lines i.e. human breast carcinoma cell
195 lines (MCF-7 and MDA-MB-231), human intestine
196 epithelial cell line (HT-29) and human ovarian
197 carcinoma cell line (SKOV-3), was carried out by Mr.
198 Cheah Yew Hong from the Institute for Medical
199 Research (IMR), Malaysia.

200 Statistical Analysis

201 Statistical analyses were performed using Sigma plot
202 8.0. Data is presented as means standard error of
203 triplicate samples.

RESULTS

205 Two chemical constituents have been isolated from
206 the crude hexane extract of *P. sarmentosum* identified
207 as β -sitosterol (Compound 1) and 3-(3',4',5'-
208 trimethoxyphenylpropanoyl)-pyrrolidine (Compound 2)
209 and two amides have also been isolated from the crude
210 EtOAc extract namely as 3-(4'-
211 methoxyphenylpropanoyl)pyrrole (Compound 3) and *N*-
212 (3-phenylpropanoyl)pyrrole (Compound 4).

213 The antibacterial activity using disk diffusion
214 method, followed by the determination of MIC and
215 MBC were presented in Table 1 and Table 2. The

Table 3. Percentage of cells survival on cytotoxicity assay of isolated compounds of *P. sarmentosum* by SRB assay

Compounds	Percentage of cell survival (%) at 20 µg/ml of samples			
	Cell lines			
	MCF-7	SKOV3	HT-29	MDA-MB-231
(1)	100.16	128.59	102.54	110.43
(2)	97.39	110.26	133.71	113.79
(3)	93.67	105.23	102.49	100.46
(4)	69.00	122.31	91.53	94.28

SRB = sulphorhodamin B, MCF-7 and MDA-MB-231 (human breast cancer cell lines), SKOV3 (human ovarian carcinoma cell lines), HT-29 (human colon/intestinal carcinoma cell lines).

216 isolated compounds were also screened for their
217 cytotoxic assay using SRB assay. Their activities are
218 given in Table 3.

219 DISCUSSION

220 The isolated compounds were identified based on
221 the physical, chemical and spectroscopic properties and
222 comparison with data of the literatures. This is the first
223 reported of the isolation of 3-(4'-
224 methoxyphenylpropanoyl) pyrrole from *P.*
225 *sarmentosum*. It was reported previously from *Piper*
226 *lolot* C.DC., from Vietnam [16]. Occurrence of
227 chemical constituents of a plant species depends on
228 several factors, such as location or environment that will
229 probably give variation in constituents. Geographical
230 distribution, seasons, different plant parts and
231 morphology, climate as well as ecological conditions
232 may also influence the biosynthesis of the secondary
233 metabolites of the plants. This is may be the reason why
234 the chemical constituents of this species are different
235 from the same species which were reported previously.

236 As shown in Tables 1 and 2, all isolated compounds
237 were found active towards Gram positive bacteria
238 except 3-(4'-methoxyphenylpropanoyl)pyrrole that
239 shown no activity against both Gram negative and
240 positive bacteria. 3-(3',4',5'-Trimethoxyphenyl-
241 propanoyl)pyrrolidine showed significant activity
242 against *B. subtilis* (MIC 500 µg/ml, MBC 1000 µg/ml)
243 followed by β -sitosterol (MIC and MBC 500 µg/ml) and
244 *N*-(3-phenylpropanoyl)pyrrole (MIC and MBC 125
245 µg/ml) against *S. aureus*. But, activity of these
246 compounds is not as good as the activity of positive
247 control streptomycin sulphate (MIC 3.91 µg/ml and
248 MBC 7.81 µg/ml). All isolated compounds exhibited no
249 activity towards Gram negative bacteria.

250 In the toxicity screening using SRB assay showed
251 that all isolated compounds have the percentage of the
252 cell survival was higher than 50%. Thus, indicated that
253 none of these compounds was active as anticancer agent
254 (Table 3).

255 As the conclusion, geographical distribution,
256 location or environment, seasons, different plant parts,
257 climate as well as ecological conditions may influence
258 the biosynthesis of the secondary metabolites of the
259 plants species which portray the variation in chemical
260 constituents. Investigations on the methanolic extracts
261 of these *Piper* species should be carried out. Different
262 models of biological activities should be performed on
263 the crude extracts and pure compounds to verify the
264 mode of action of the active candidates.

265 REFERENCES

2661. Rishton JM. Natural Products as a Robust Source of New Drugs
267 and Drug Leads: Past Successes and Present Day Issues. *Am J*
268 *Cardiol* 2008; 101: 43D-49D.
2692. Scott IA, Jensen HR, Philogene BJR, Arnason JTA. Review of
270 *Piper* spp. (Piperaceae) Phytochemistry, Insecticidal Activity
271 and Modes of Action. *Phytochem Rev* 2008; 7:65-75.
2723. Jaramillo MA, Manos PS. Phylogeny and Pattern of Floral
273 Diversity in the Genus *Piper* (Piperaceae). *Am J Bot* 2001;
274 88:706-16.
2754. Wiart C. *Medicinal Plants of Asia and the Pacific*. Boca Raton,
276 London, New York: Taylor & Francis. 2006.
2775. Parmar VS, Jain SC, Bisht KS, Jain R, Taneja P, Jha A, Tyagi
278 OD, Prasad AK, Wengel J, Olsen CE, Boll PM. Phytochemistry
279 of the Genus *Piper*. *Phytochemistry* 1997; 46:597-503.
2806. Rahman NNNA, Furuta T, Kojima S, Takane K, Mohd MA.
281 Anti-malarial Activity of Extracts of Malaysian Medicinal
282 Plants. *J Ethnopharmacol* 1999; 64:249-54.
2837. Ong HC, Nordiana M. Malay Ethno-medico Botany in Machang,
284 Kelantan, Malaysia. *Fitoterapia* 1999; 70: 502-13.
2858. Ong HC, Norzalina J. Malay Herbal Medicine in Gemenech,
286 Negeri Sembilan, Malaysia. *Fitoterapia* 1999; 70:10-4.
2879. Masuda T, Inazumi A, Yamada Y, Padolina WG, Kikuzaki H,
288 Nakatani N. Antimicrobial Phenylpropanoids from *Piper*
289 *sarmentosum*. *Phytochemistry* 1991; 30:3227-8.
29010. Likhithitayawuid K, Ruangrasi N. Structural Elucidation and
291 Synthesis of New Components Isolated from *Piper sarmentosum*
292 (Piperaceae). *Tetrahedron* 1987; 43:3689-94.
29311. Stohr JR, Xiao PG, Bauer R. Isobutylamides and a new
294 methylbutylamide from *Piper sarmentosum*. *Planta Med* 1999;
295 65:175-7.
29612. Rukachaisirikul T, Siriwanakit P, Sukcharoenphol K,
297 Wongvein C, Rattanaweang P, Wongwattanavuch P.
298 Suksamrarn A. Chemical Constituents and Bioactivity of *Piper*
299 *sarmentosum*. *J Ethnopharmacol* 2004; 93:173-6.
30013. Tuntiwachwuttikul P, Phansa P, Pootaengon Y, and Taylor WC.
301 Chemical Constituents of the Roots of *Piper sarmentosum*.
302 *Chem Pharm Bull* 2006; 54:149-51.
30314. Gunston FD. *Fatty Acid and Lipid Chemistry*. London, Glasgow,
304 New York: Blacky Academic & Professional 1996.
30515. Achenbach H, Fietz W, Woerth J, Waibel R, Portecop J.
306 Constituents of Tropical Medicinal Plants IXX GC/MS-
307 investigations of the Constituents of *Piper amalago*-30, New
308 Amides of the Piperine-type. *Planta Med* 1986; 1:12-7.
30916. Luger P, Weber M, Dung NX, Luu VT, Rang DD, Tuong DT,
310 Ngoc PH. The Crystal Structure of 3-(4'-
311 Methoxyphenylpropanoyl)pyrrole of *Piper lolot* C.DC., from
312 Vietnam. *Crystal Res Tehmol* 2002; 37:627-33.
31317. Mackeen MM, Ali AM, El-Sharkawy SH, Manap MY, Salleh
314 KM, Lajis NH, Kawazu K. Antimicrobial and Cytotoxic
315 Properties of Some Malaysian Traditional Vegetables. *Int J*
316 *Pharmacog* 1997; 35:174-8.
31718. Zavala MA, Perez GS, Perez GRM. Antimicrobial Screening of
318 Some Medicinal Plants. *Phytother Res* 1997; 11:368-71.
31919. Arias ME, Gomez JD, Cudmani NM, Isla MI. Antibacterial
320 Activity of Ethanolic and Aqueous Extracts of *Acacia aroma*
321 Gill. *Ex Hook et Arn. Life Sciences*, 2004; 75:191-202.

32220. Houghton P, Fang R, Techatanawat I, Steventon G, Hylands PJ, 331 F. Ahmad, Department of Chemistry, Faculty of Science, Universiti
323 Lee CC. The sulphorhodamine (SRB) Assay and other 332 Teknologi Malaysia, 81310 Skudai, Johor Bahru, Malahysia. E-
324 Approaches of Testing Plant Extracts and Derived Compounds 333 mail: farediah@kimia.fs.utm.my
325 for Activities Related to Reputed Anticancer Activity. *Cancer*
326 *Methods* 2007; 42:377-87. 334 H. M. Sirat, Department of Chemistry, Faculty of Science, Universiti
327 **CURRENT AUTHOR ADDRESSES** 335 Teknologi Malaysia, 81310 Skudai, Johor Bahru, Malahysia. E-
328 E. Atiax, Department of Chemistry, Faculty of Science, Universiti 336 mail: hasnah@kimia.fs.utm.my
329 Teknologi Malaysia, 81310 Skudai, Johor Bahru, Malahysia. E-337 D. Arbain, Faculty of Pharmacy, University of Andalas, Padang, West
330 mail: emrizal_atiax@yahoo.com (Corresponding author) 338 Sumatra, Indonesia. E-mail: arbain_d@yahoo.com
339

UNCORRECTED PROOF