Original Article

Assessment of the structural validity of the domestic violence healthcare providers' survey questionnaire using a Nigerian sample

Ime A John, ^{a,*} Stephen Lawoko ^a

KEV WODDS

^a Division of Social Medicine, Department of Public Health Sciences, Karolinska Institutet. SE-17176 Stockholm, Sweden.

Abstract: Background: There has been increased advocacy to involve healthcare providers in the prevention

KEI WORDS	of intimate partner violence (IPV) through screening for it in healthcare. Yet, only one in ten
	providers screen for IPV, suggesting barriers. Understanding the readiness of healthcare providers
Domostio violonoo	to screen for IPV is therefore paramount. The Domestic Violence Healthcare Provider Survey Scales
Domestic violence-	(DVHPSS) is a previously validated, comprehensive scale to study readiness of healthcare providers
survey	to screen for IPV. However, an understanding of its usefulness in the Sub-Saharan African context
Structural validitv	remains elusive. The current study undertook to examine the structural validity of the DVHPSS in
II	Nigeria.
Healincare-	Methods: Exploratory factor analysis and Cronbach's Alpha were run to reveal the factorial
providers	structure and reliability of the instrument/subscales respectively. Established thresholds were used to
Nigeria	determine significant factor loadings and alphas coefficient.
	Results: A six factor model emerged, with 2 factors similar to the original scale, another two
	differing slightly and a further two factors resulting from a splitting up of the original combination of
	victim/provider safety to having distinct victim and provider safety subscales.
	Conclusions: With slight modifications, the DVHPSS can be use to study IPV screening among
	Nigerian healthcare professionals. Introducing screening protocols could promote better
	understanding of crucial questions that were lost in the analysis.
Accepted 2010-03-31	© 2010 KUMS, All right reserved

*Corresponding Author at:

Dr. Ime A John: Department of Public Health Sciences, Division of Social Medicine, Karolinska Institutet, SE 171 76 Stockholm, Sweden, Mobil: +46 762408518 Fax: +46 8 307351 E-mail: <u>ime.john@ki.se</u> (I.A. John).

© 2010 KUMS, All right reserved

Introduction

ntimate partner violence (IPV) against women remains a public health concern worldwide despite significant law reforms aimed at curbing abuse in various countries.¹⁻³ This is indicative of the fact that secondary prevention measures on their own may not be reaching the desired goal. Resources thus should be redirected towards primary prevention. Healthcare professionals can play an important role in this regard through screening for IPV among their patients. Over the past decade, a number of instruments to assist healthcare providers in screening for IPV have been developed, particularly in Europe and America.⁴⁻¹⁰ Despite these developments, barely 10% of health care providers screen for IPV in those settings evidencing barriers to screening for IPV in healthcare.^{11,12} Barriers to screening could evolve from the provider or from the client. An assessment of providers' readiness to screen for IPV as well as Clients readiness to be screened for IPV thus seems paramount before effective

A number of instruments have emerged in the past decade to assess providers' readiness to screen for IPV.^{13,14} Among the most comprehensive of them is the Domestic Violence Healthcare Provider Survey Scales (DVHPSS).¹⁵ The scale measures healthcare professionals' readiness to screen in terms of their perceived knowledge/ efficacy in screening, conflicting professional roles, availability of social support networks to which IPV victims can be referred, whether inquiries into IPV may pose safety challenges for patient/care providers and providers' general attitudes towards screening for IPV. The DVHPSS has been validated in the western context but to the best of our knowledge, is not yet in use in the Sub-Saharan African context. Thus, knowledge of the readiness of healthcare providers to screen for IPV in the Sub-Saharan African context, as well as of their screening behavior per se remains elusive. This study sets the foundation to fill this knowledge gap by validating the DVHPSS for use in Nigeria.

Specifically, this study will assess the structural validity of the DVHPSS in terms of its factorial structure and sub-scale reliability.

Methods

Study settings, design and participants

This study was carried out at the Amino Kano Teaching Hospital, in Kano, Nigeria, which is the largest multidepartmental federal health institution in Kano state. The staff and patients have a multi-ethnic background and speak English, the official language in the country. In general, the staff members at this hospital have not undergone any specific training in screening for IPV. All Health Care Providers having regular contact with patients (i.e. n = 430) were informed of the study by department heads and invited to participate. Self-administered questionnaires were sent to the eligible participants of which two hundred and seventy four (response rate of 64%) returned the completed questionnaire. Voluntary participation was emphasized and informed consent given. The participants included Psychiatrists, Obstetricians and Gynecologists, Pediatricians, Physicians, Laboratory Scientists, Opticians, Nurses and Midwives. Only those laboratory employees who sometimes meet patients were included. The professions with less than 5 participants (i.e. opticians and laboratory Assistants) were grouped under "others". Table1 shows some demographic and occupational characteristics of the participants.

Ethical consideration

This study received ethical approval from the Nigerian Institute of Medical Research, Lagos, Nigeria and the authorities of Aminu Kano Teaching Hospital, Kano. The aims and relevance of the study were further emphasized in a separate document accompanying the questionnaires. Questionnaires were delivered to all the clinical and laboratory departments within the hospital. Only laboratory employees who sometimes meet patients were included.

Table	1:	Characteristics	of	Participants
	••		•••	

		N	%
Profession			
	Doctor	156	58
	Nurse	61	22.7
	Midwife	12	4.4
	Social worker	29	10.8
	Others	11	4.1
Gender			
	Male	147	56.5
	Female	113	43.5
Age (years)			
	21-30	116	45.1
	31-40	105	40.9
	41-60	36	14.0
Marital status			
	Married	145	55.1
	Single	108	41.1
	Divorced	6	2.3
	Separated	4	1.5
Religion			
	Muslim	114	42.7
	Catholic	47	17.6
	Protestant	76	28.5
	Others	30	11.2
Ethnicity			
	Hausa	84	31.5
	lbo	70	26.2
	Yoruba	28	10.5
	Others	85	31.8
Department			
	Medicine	41	15
	Surgery	41	15
	Pediatric	44	16.1
	Obstetrics/Gynecology	36	13.2
	General practice	81	29.7
	Others	30	11

N = absolute number

% = percentage of total within the group

Voluntary participation was emphasized, privacy guaranteed and informed consent given. Participants dropped off the filled questionnaires at a special collection point centrally located at the hospital.

Instrument measures

The Domestic Violence Health Care Provider Survey Scale measures healthcare providers' readiness to screen for IPV as well as actual screening activity.¹⁵ The instrument has been previously validated with promising results in developed countries. This paper attempts to assess its structural validity in the Nigerian context. The questionnaire consists of the following 5 subscales:

The perceived self-efficacy subscale (4 items) which assesses providers' own perceived efficacy in inquiring about IPV (details in table 2).

The system support sub-scale (4 items) which assesses healthcare providers' access to support networks for referral/management of IPV victims (details in table 2).

The professional roles resistant/fear of offending clients sub-scale (6 items) which assesses providers' opinions on whether inquiries about IPV may conflict with ethical issues governing their communication with clients (details in table 2).

The blame victim sub-scale (7 items) which assesses providers' attitudes towards victims (details in table 2).

The victim/provider safety sub-scale (10 items) which assesses provider perception on whether inquiring about IPV from batteres would further jeopardize victim/care provider safety (details in table 2).

All items require the respondent to take a position on specific statements. The response alternatives to each statement range from 1 (strongly disagree) to 5 (strongly agree).

Some of the items (statements) are phrased in a way such that their scores need to be reversed to match with other items in the same scale before any further analysis can be done.

Statistical analysis

Exploratory factor analysis using principal component method was performed to test underlying factors and their stability as expressed in the factor loadings. Varimax rotation was applied to limit the number of high loadings under the same factor. This would enhance clearer identification of items emerging under each subscale. Criteria for the number of resulting significant factors was based on Kaiser Criterion and confirmed with screen plots.¹⁵⁻¹⁷ Items with factor loading of at least 0.30 were considered significant; this is based on criteria for significant correlation.¹⁸ The contribution of each factor in explaining the total variation in the item pool was reported. Significant factors (i.e. those having a highest loading of over 0.30) were tested for internal consistency using Cronbach's Alpha. Each item was then scrutinized further to assess whether the removal of that item would improve the alpha coefficient. If removal of an item implied improved alpha, that item would be removed and the reliability test rerun without that item. The process would continue until a point of saturation was reached (i.e. removal of additional items would not improve alpha). Alpha coefficients of at least 0.70 were considered significant, a threshold adequate for research purposes.^{19,20} The resulting items/scales following the reliability test were then re-examined in a new factor analysis. If any of the highest item loadings was less than 0.30, the process described above (i.e. series of factor analyses and reliability test) would continue until all remaining items loaded at least 0.30, the a priori set threshold. Where double loadings were evident, the item was assigned to the factor under which it loaded highest. Bivariate correlations were run to investigate the factor distinctiveness of the final factor solution.^{19, 20}

Prior to the above analyses, certain procedures were carried out to clean data. First, only participants who had responded to all items were included in the analyses above as failure to do so may introduce erroneous estimates. Second, items were checked for normality using the skewness statistic and its confidence interval. Skewness statistic of magnitude zero is an indication of perfect symmetry. All analyses were run with SPSS version 16.

Results

Data cleansing

Only participants who had responded to all items of the original scales were assessed in the factor analyses and reliability tests. This resulted in a total of 162 participants (59% of total respondents to the questionnaire). The results of the normality test revealed that the confidence intervals for the skewness statistic for each of the 35 items included, or was close to, zero, suggesting minor or no violation of normality assumptions.

Initial factorial structure with all items

When all items of the original instrument were included in the factor analysis, 6 factors emerged based on the Kaiser Criterion (Eigen values > 1) (table 2). The victim/provider safety subscale of the original instrument emerged as two separate subscales in our data, reflecting victim safety and provider safety respectively. The majority of the items of the

Note: Factors loading over 0.30 appear in bold. The extraction method used was Principal component Analysis and rotation method: Varimax rotation

www.SID.ir

Archive of SID

78 Injury & Violence

able 2: Rotated factor loadings for Domestic Violence Healmcare Providers surve	y scales re	estricted to	o o raciors			
Components	1	2	3	4	5	6
Professional Role Resistance/Fear of offending the Patients						
I am afraid of offending patients if I ask about their abusive behavior	0.62	-0.04	0.08	0.02	0.08	-0.04
I am afraid of offending the patient if I ask about DV	0.69	0.08	0.02	0.00	-0.10	-0.24
Asking patients about DV is an invasion of their privacy	0.78	0.08	0.16	-0.12	0.06	-0.07
It is demeaning to patients to question them about abuse	0.72	0.18	0.20	-0.03	0.08	0.13
If I ask non-abused patients about DV, they will get very angry	0.34	0.13	0.11	-0.04	0.01	-0.05
It is not my place to interfere with how a couple chooses to resolve conflicts	0.57	0.04	0.10	-0.10	-0.13	0.07
When challenged, batterers frequently direct their anger toward health care providers	0.49	-0.13	0.34	0.09	0.11	0.22
If patients do not reveal abuse to me, then they feel it is none of my business	0.14	0.17	0.07	-0.08	-0.04	-0.22
Blame Victim						
A victim must be getting something out of the abusive relationship, or else he/she would leave.	0.11	0.68	-0.06	0.15	0.06	0.06
People are only victims if they choose to be.	0.06	0.45	0.18	0.18	-0.10	0.07
When it comes to domestic violence victimization, it usually "takes two to tango."	0.00	0.66	0.10	0.06	0.17	0.16
I have patients whose personalities cause them to be abused.	0.02	0.59	0.26	0.02	-0.11	0.15
Women who choose to step out of traditional roles are a major cause of DV.	-0.06	0.74	0.14	-0.00	0.03	-0.08
The victim's passive-dependent personality often leads to abuse.	0.17	0.46	0.14	0.20	-0.04	0.08
The victim has often done something to bring about violence in the relationship	0.17	0.62	0.16	0.13	0.34	-0.18
Victim safety						
I think that investigating the underlying cause of a patient's injury is not part of medical care	0.21	0.02	0.42	-0.04	-0.15	0.11
I feel it is best to avoid dealing with the batterer out of fear and concern for the victim's safety	0.20	0.12	0.71	-0.05	0.14	-0.04
There is no way to ask batterers about their behaviors without putting the victims in more danger	0.11	0.18	0.81	-0.03	-0.01	0.01
I am afraid if I talk to the batterer, I will increase risk for the victim	0.10	0.15	0.74	0.00	0.00	-0.05
I have ready access to information detailing management of DV	0.13	0.26	0.46	0.37	0.06	0.23
Perceived Self-efficacy						
There are strategies I can use to encourage batterers to seek help.	-0.09	0.09	-0.21	0.74	0.11	0.26
There are strategies I can use to help victims of DV change their situation.	0.03	0.12	-0.06	0.79	0.28	0.05
I feel confident that I can make appropriate referrals for batterers.	-0.07	0.13	0.14	0.64	0.39	0.01
I feel confident that I can make the appropriate referrals for abused patients.	-0.27	0.06	0.21	0.59	0.12	0.05
There're ways I can ask batterers about their behavior that will minimize risk to the potential victim	0.19	0.17	0.16	0.49	-0.01	0.42
I don't have the time to ask about DV in my practice	0.03	0.07	-0.00	-0.08	-0.12	-0.15
System support						
I have ready access to medical social workers or community advocates to assist in the management of DV.	0.12	0.11	0.17	0.15	0.72	0.08
I feel that medical social work personnel can help manage DV patients.	-0.07	0.08	-0.07	0.31	0.50	0.36
I have ready access to mental health services should our patients need referrals.	-0.01	0.06	-0.11	0.16	0.72	0.12
I feel that the mental health services at my clinic or agency can meet the needs to DV victims in cases where they are needed.	-0.04	-0.01	0.08	0.21	0.67	0.22
Provider safety						
I feel there are ways of asking about battering behavior without placing myself at risk	-0.02	-0.07	-0.02	0.08	0.17	0.69
I feel I can effectively discuss issues of battering and abuse with a battering patient	-0.16	0.05	-0.03	0.28	0.15	0.77
I feel I can discuss issues of battering and abuse with a battering patient without further endangering the victim I can reluvant to ack hatterers about their abusive behavior out of concern for my	0.06	0.22	0.03	0.01	0.20	0.70
personal safety.	0.33	-0.02	0.33	-0.00	0.23	-0.13
Inere is not enough security at my work place to sately permit discussion of DV with batterers	0.06	0.12	0.26	0.04	0.02	0.09
Eigenvalues	5.98	5.06	2.22	1.88	1.67	1.51
% of Variance	17.09	14.47	6.35	5.37	4.77	4.31

Table 2: Rotated factor loadings for Domestic violence Healthcare Providers survey scales restricted to 6 factors

with Kaiser Normalization (eigenvalues >1).

original DVHPSS exhibited significant factor loading according to the a priori decided threshold of 0.30 (table 2).

Factor 1 emerged as professional role resistant/fear of offending the patients subscale with all but two items of the original scale loading significantly under this factor. Moreover, one item that in the original instrument belonged to the provider safety scale (i.e. "when challenged, batterers frequently direct their anger towards healthcare providers) now loaded significantly under factor 1. This factor included a total of 7 items and explained 17% of the variation in the total item pool.

Factor 2 emerged as Blame victim subscale with all 7 items of the original scale loading significantly under the same. The factor explained about 14.5% of the variation in the total item pool.

Factor 3 reflected a victim safety subscale component, though two additional items from the originally perceived self-efficacy and professionally resistant subscales respectively (i.e. "I have ready access to information detailing management of domestic violence" and "I think that investigating the underlying cause of a patient's injury is not part of medical care") now loaded significantly under this factor. This factor included a total of 5 items and explained 6.5% of the variation in the total item pool.

Factor 4 emerged as a perceived self-efficacy subscale with all but two of the items in the original subscale loading significantly under this factor. This factor included a total of 5 items and explained about 5.4% of the variation in the total item pool.

Factor 5 emerged as system support subscale with all 4 items of the original scale loading significantly under this factor. The factor explained about 4.8% of the variation in the total item pool.

Factor 6 reflected a Provider safety subscale with three of the original six items in the victim/provider safety scale loading significantly under this factor. The factor explained 4.3% of the variation in the total item pool.

Internal reliability

Only items with highest loadings of 0.30 and above were subjected to a reliability test. Thus, 4 items of the total 35 were dropped (i.e. "I don't have time to ask about Domestic Violence in my practice", "If patients do not reveal abuse to me, then they feel it is none of my business", "There is not enough security at my work place to safely permit discussion of domestic violence with batterers" and "I am reluctant to ask batterers about their abusive behavior out of concern for my own safety").

As indicated in table 3, the internal reliability of the 7

items of the professional role resistant/fear of offending the patients subscale (meeting above criteria) was 0.80. Removal of additional items from this scale would only reduce the internal reliability as expressed in the column "Cronbach's alpha if item removed" of table 3. In addition, the internal reliability of the 7 items of the Blame victim subscale (meeting above criteria) was 0.77. Removal of additional items from this scale would only reduce the internal reliability. The internal reliability of the 5 items of the new victim safety subscale was 0.73. However, removal of two of these items (i.e. "I have ready access to information detailing management of domestic violence" and "I think that investigating the underlying cause of a patient's injury is not part of medical care") was bound to improve alpha. Thus, when these items were removed alpha increased to 0.78. The internal reliability of the 5 items of the perceived self-efficacy subscale was 0.77. Removal of additional items from this scale would only reduce the internal reliability. Likewise, the internal reliability of the 4 items of the system support subscale was 0.73. Removal of additional items from this scale would only reduce the internal reliability. Finally, the internal reliability of the 3 items of the provider safety subscale was 0.72. Removal of additional items from this scale would only reduce the internal reliability.

In total therefore, the reliability test dropped an additional two items.

Final factorial structure

The items emerging from the reliability test (i.e. 29 items) were now subjected to a new factor analysis. As indicated in table 4, six factors emerged based on Kaiser Criterion (Eigen values > 1) and all items now loaded significantly (i.e. highest factor loadings of at least 0.30) under respective factor. The explanatory power of each factor in explaining the total variation in the item pool is also reported in table 4.

Inter-factor correlation

As indicated by the bivariate correlations in table 5, significant correlations ranging in magnitude between 0.17-0.53 were found between most factors.

Discussion

The factorial structure and internal reliability of the Domestic Violence Healthcare Provider Survey Scales (DVHPSS) in a sample of Nigerian healthcare providers was tested and the instrument to a small extent refined following a priori defined criteria for inclusion. The analysis was thus designed so as to maintain only items an factors that met these criteria. The

Table 3: Internal Reliability scores for items of Domestic violence Healthcare Providers survey scales

Scales items	No of scale items	Cronbach's Alpha	Cronbach's Alpha*
Factor 1: Professional Role Resistance/Fear of offending the Patients scale	7	0.80	
Asking patients about DV is an invasion of their privacy			0.78
It is demeaning to patients to question them about abuse			0.76
If I ask non-abused patients about DV, they will get very angry			0.77
I am afraid of offending the patient if I ask about DV			0.79
I am afraid of offending patients if I ask about their abusive behavior			0.77
It is not my place to interfere with how a couple chooses to resolve conflicts			0.76
When challenged, batterers frequently direct their anger toward health care providers			0.80
Factor 2: Blame Victim scale	7	0.77	
A victim must be getting something out of the abusive relationship, or else he/she would leave.			0.74
People are only victims if they choose to be.			0.75
When it comes to domestic violence victimization, it usually "takes two to tango."			0.73
I have patients whose personalities cause them to be abused.			0.75
Women who choose to step out of traditional roles are a major cause of DV.			0.74
The victim's passive-dependent personality often leads to abuse.			0.76
The victim has often done something to bring about violence in the relationship			0.75
Factor 3: Victim safety scale	5	0.73	
There is no way to ask batterers about their behavior without putting the victims in more danger			0.58
I am afraid if I talk to the batterer, I will increase risk for the victim			0.72
I feel it is best to avoid dealing with the batterer out of fear and concern for the victim's safety			0.78
I have ready access to information detailing management of DV			0.78
I think that investigating the underlying cause of a patient's injury is not part of medical care			0.78
Factor 4: Perceived Self-efficacy scale	5	0.77	
There are strategies I can use to encourage batterers to seek help.			0.72
There are strategies I can use to help victims of DV change their situation.			0.68
I feel confident that I can make appropriate referrals for batterers.			0.72
I feel confident that I can make the appropriate referrals for abused patients.			0.75
There're ways I can ask batterers about their behavior that will minimize risk to the potential victim			0.76
Factor 5: System support scale	4	0.73	
I have ready access to medical social workers or community advocates to assist in the management of DV.			0.68
I feel that medical social work personnel can help manage DV patients.			0.69
I have ready access to mental health services should our patients need referrals.			0.65
I feel that the mental health services at my clinic or agency can meet the needs to DV victims in cases where they are needed.			0.67
Factor 6: Providers safety scale	3	0.72	
I feel I can effectively discuss issues of battering and abuse with a battering patient.			0.54
I feel there are ways of asking about battering behavior without placing myself at risk			0.64
I feel I can discuss issues of battering and abuse with a battering patient without further endangering the victim			0.71
The transformed The transforme			

DV= Domestic Violence

results in many regards mirrored that of the original instrument. The factors "blame the victim" and "system support" in the original questionnaire was confirmed in our data without any exceptions (i.e. all items in the original subscales met inclusion criteria in our data). Though the other factors did not differ substantially from the original scales, some important observations warranting acknowledgement were made.

First, the victim/provider subscale that formed a distinct sub-scale in the original instrument was split up in our data into two separate scales i.e. victim safety and provider safety subscales. This suggests that healthcare providers in the setting

Table 4. Emerging factor loadings for Domestic violence Healthcare Providers survey	scales					
Components	1	2	3	4	5	6
Professional Role Resistance/Fear of offending the Patients						
I am afraid of offending patients if I ask about their abusive behavior	0.63					
I am afraid of offending the patient if I ask about DV	0.70					
Asking patients about DV is an invasion of their privacy	0.79					
It is demeaning to patients to question them about abuse	0.73					
If I ask non-abused patients about DV, they will get very angry	0.41					
It is not my place to interfere with how a couple chooses to resolve conflicts	0.61					
When challenged, batterers frequently direct their anger toward health care	0.45					
providers Plana Viatin						
A victim must be getting something out of the gbusive relationship, or else he/she						
would leave.		0.70				
People are only victims if they choose to be.		0.64				
When it comes to domestic violence victimization, it usually "takes two to tango."		0.74				
I have patients whose personalities cause them to be abused.		0.62				
Women who choose to step out of traditional roles are a major cause of DV.		0.53				
The victim's passive-dependent personality often leads to abuse.		0.50				
The victim has often done something to bring about violence in the relationship		0.41				
System support						
I have ready access to medical social workers or community advocates to assist in the management of DV.			0.75			
I feel that medical social work personnel can help manage DV patients.			0.56			
I have ready access to mental health services should our patients need referrals.			0.64			
I feel that the mental health services at my clinic or agency can meet the needs to DV victims in cases where they are needed.			0.67			
Perceived Self-efficacy						
There are strategies I can use to encourage batterers to seek help.				0.69		
There are strategies I can use to help victims of DV change their situation.				0.76		
I feel confident that I can make appropriate referrals for batterers.				0.67		
I feel confident that I can make the appropriate referrals for abused patients.				0.59		
There're ways I can ask batterers about their behavior that will minimize risk to the potential victim				0.43		
Victim safety						
I feel it is best to avoid dealing with the batterer out of fear and concern for the victim's safety					0.72	
There is no way to ask batterers about their behavior without putting the victims in more danger					0.82	
I am afraid if I talk to the batterer, I will increase risk for the victim					0.76	
Provider safety						
I feel there are ways of asking about battering behavior without placing myself at risk						0.68
I feel I can effectively discuss issues of battering and abuse with a battering patient						0.77
I feel I can discuss issues of battering and abuse with a battering patient without further endangering the victim						0.77
Eigenvalues	5.47	4.61	2.20	1.58	1.52	1.29
% of Variance	18.24	15.37	7.32	5.29	5.06	4.31

studied recognize victim and provider safety as two distinct aspects of screening for IPV unlike their peers in the developed countries.¹⁵ In addition, some aspects related to victim and provider safety (i.e. "There is not enough security at my work place to safely permit discussion of domestic violence with batterers" and "I am reluctant to ask batterers about their abusive behaviour out of concern for my own safety") were dropped out subsequently in the analysis as they did not meet the a priori conditions set. Noteworthy here is that these factors were concerned with how to deal with the perpetrators and not the victims. A plausible explanation as to why these issues loaded low could be that respondents may

Safety

Safety

Self-

Table 5. Bivariate Pearson Correlations of Domestic Violence healthcare Providers Survey Scales								
	Professio nal Role	Blame Victim	System Support	Victim Safety	Self- efficacy	Provider Safety		
Professiona I Role	1.000							
Blame Victim	0.257**							
System Support	0.006	0.198*						
Victim	0.406**	0.382**	0.060					

0.320** 0.528** efficacy Provide 0.049 0.171* 0.424** 0.021 0.431** 1.000

0.075

**Correlation is significant at 0.01 level (2-tailed)

* Correlation is significant at 0.05 level (2-tailed)

-0.081

not have identified with these questions. Indeed, it may be difficult to identify batterers among patients, precluding an assessment of batterer-related security risk. Moreover, in the Nigerian context men rarely accompany their partners/children to hospital making access to the potential batterers difficult, particularly in reproductive health and pediatric units.

Second, the perceived self-efficacy and professional role resistance scales lost three items that had to do with lack of time for IPV inquiry, access to information to manage IPV, and challenging whether IPV is part of the medical practice. The reason why these items loaded low is difficult to explain and deserves further scrutiny in future work. However, a plausible explanation, at least for the later two, is that IPV inquiry is not yet an integral part of healthcare practice at this facility. An understanding of what access to information on how to manage IPV implied or the role of IPV inquiry in medical practice might have varied, leading to random response to these items. Introducing protocols on IPV inquiry at this facility in the future thus could help align providers understanding of such issues.

To assess factor distinctiveness, bivariate correlations were established between the emerging factors. These correlations, though statistically significant, ranged from low to moderate, indicating that though inter-related, the factors represent rather distinct aspects of provider readiness to screen.

The current study attempted to assess the usefulness of an already existing instrument (developed in the western world) in the Nigerian context and found the instrument applicable based on its structural validity. However, the item pool used in this study may not be exhaustive of challenges to screening for IPV in the Nigerian context. Qualitative studies could

reveal additional challenges to screening specific to the Nigerian context. Indeed dissimilar cultural values could imply unique sets of challenges to screening in each culture. For example, recent studies on violence indicate that in many Sub-Saharan countries including Nigeria, wife beating remains an acceptable societal norm.²¹ How such norms may pose specific challenges to screening both from a patient and provider perspective remain to be studied. In addition, other aspects of the validity of the instrument (e.g. concurrent and discriminate validity) may help further validate the usefulness of this instrument in the Nigerian context. These aspects are currently being studied separately.

In conclusion, the factorial stability and internal reliability of the DVHPSS scale in the Nigerian sample has to a large extent been confirmed and can therefore be used to scrutinise readiness to screen for IPV among Nigerian healthcare professionals. However, the Nigerian healthcare providers made an important distinction between victim and provider safety unlike peers in the developed countries.¹⁵ In addition, batterer-related aspects of screening did not figure as significant factors in this analysis, suggesting that cultural specific factors could play a role. Finally, issues pertaining as to whether IPV should be considered a part of medical practice or access to information on IPV management did not specifically load under any factor, suggesting that the question may have been variedly understood. Clarification of what screening implies (e.g. using protocols) may be useful to align healthcare providers' understanding of these items. Further investigation of the other aspects of validity is warranted to understand occupational, demographic and cultural aspects that could impede or promote screening activity in this context.

Acknowledgement

The authors would wish to thank the Management and Staff of Aminu Kano Teaching Hospital, Kano for their cooperation during this study.

Funding: The funding source for I.A.J's PhD studies had no role in the study design, data collection, analysis, interpretation and in writing the report.

Competing interest: None declared.

Ethical approval: Ethical approval for this study was received from the Nigerian Institute of Medical Research, Lagos, Nigeria and the authorities of Aminu Kano Teaching Hospital, Kano. Nigeria.

References

- 1. Johnson M, Elliott BA. Domestic violence among family practice patients in midsized and rural communities. J Fam Pract. 1997; 44: 391-400.
- Coker AL, Smith PH, Bethea L, King MR, McKeown RE. Physical health consequences of physical and psychological intimate partner violence. Arch Fam Med. 2000; 9: 451–7.
- 3. Hiese L, Ellsberg M & Gottemoeller M. Ending violence against women. Population Reports, Population Information Program, Series L, No. 11. Baltimore: Johns Hopkins University School of Public Health, 1999.
- 4. Feldhaus KM, Kozio-Mclain J, Amsbury HL, Norton IM, Lowenstein SR, Abbott JT. Accuracy of 3 brief screening questions for detecting partner violence in the emergency department. J Am Med Assoc. 1997; 277:1357–61.
- 5. McFarlane J, Parker B, Soeken K, Bullock L: Assessing for abuse during pregnancy. Severity and frequency of injuries and associated entry into prenatal care. J Am Med Assoc. 1992; 267:3176–8.
- 6. Sohal H, Eldridge S, Feder G. The sensitivity and specificity of four questions (HARK) to identify intimate partner violence: a diagnostic accuracy study in general practice. BMC Fam Pract. 2007; Aug 29;8:49.
- 7. Brown JB, Lent B, Brett PJ, Sas G, Pederson LL. Development of the Woman Abuse Screening Tool for Use in Family Practice. Fam Med. 1996 Jun; 28(6):422-8.
- 8. Sherin KM, Sinacore JM, Li XQ Li, Zitter RE, Shakil A. HITS: A Short Domestic Violence Screening Tool for Use in a Family Practice Setting. Fam Med. 1998; 30:508–12.
- 9. Weiss SJ, Ernst A, Cham E, Nick TG. Development of a screen for ongoing intimate partner violence. Violence Vict. 2003; 18:131-41.
- 10. Smith PH, Earp JA, DeVellis R. Measuring battering: Development of the Women's Experience with Battering (WEB) Scale. Women's Health. 1995; 1: 273-88.
- 11. Roelens K, Verstraelen H, Van Egmond K & Temmerman M. A knowledge, attitudes and practice survey among obstetrician-gynaecologists on intimate partner violence in Flanders, Belgium. BMC Public Health, 2006; 6: 238.
- 12. Erikson MJ, Hill TD, Siegal RM. Barriers to domestic Violence screening in the Pediatric Setting. Pediatrics. 2001;108: 98-102.
- 13. Short LM, Alpert E, Harris JM, Surprenant ZJ.A tool for measuring physician readiness to manage intimate partner violence. Am J Prev Med. 2006;30:173-80.
- Rodríguez M.A, Bauer, HM, McLoughlin E, Grumbach K. Screening and Intervention for intimate Partner Abuse: Practices and attitudes of Primary care Physicians. J Am Med Assoc. 1999: 282: 468–74.
- 15. Maiuro RD, Vitaliano PP, Sugg NK, Thompson DC, Rivara FP. Thompson RS. Development of a health care provider survey for domestic violence: psychometric properties. Am J Prev Med. 2000; 19: 245–52.
- 16. Carroll JB. Biquartimin Criterion for rotation to Oblique simple structure in factor analysis. Science. 1957; 126: 1114–15.
- 17. Field AE, Aneja P, Rosner B. The validity of self-reported weight change among adolescents and young adults. Obesity (Silver Spring). 2007; 15: 2357–67.
- 18. Cohen J. Statistical Power Analysis for Behavioral Sciences, 2nd ed. Hillsdale: Lawrence Erlbaum Associates, 1988.
- 19. Streiner DL, Norman GR. Health measurement scales a Practical Guide to their Development and use. New York: Oxford University Press, Inc, 1989.
- 20. Nunnaly J. Psychometric theory. New York: McGraw-Hill.1978.
- 21. Uthman OA, Moradi T, Lawoko S. The independent contribution of individual-, neighbourhood-, and country-level socioeconomic position on attitudes towards intimate partner violence against women in sub-Saharan Africa: A multilevel model of direct and moderating effects. Soc Sci Med. 2009; 68:1801–09.