Journal of Plant Physiology and Breeding

2016, 6(2): 9-18 ISSN: 2008-5168



# In Vitro Propagation of Damask Rose Using the Temporary Immersion System

# Ali Bosh<sup>1</sup>, Ahmad Moieni<sup>2\*</sup>, Hamid Dehghani<sup>2</sup> and Zahra Movahedi<sup>3</sup>

Received: May 8, 2016 Accepted: December 16, 2016

<sup>1</sup>M.Sc., Department of Plant Breeding and Biotechnology, Tarbiat Modares University, Tehran, Iran

<sup>2</sup>Associate Prof., Department of Plant Breeding and Biotechnology, Faculty of Agriculture, Tarbiat Modares University, Tehran, Iran

<sup>3</sup>Assistant Prof., Department of Agronomy and Plant Breeding, Faculty of Agriculture, Malayer University, Malayer, Iran

\*Corresponding author; Email: moieni\_a@modares.ac.ir

### Abstract

For centuries, Damask rose has been one of the most popular species of the rose family. Damask rose can be propagated using vegetative methods such as micropropagation. In this research, the use of temporary immersion system was reported for the first time for Damask rose micropropagation. For this reason, a traditional temporary immersion system was designed to facilitate its micropropagation. To optimize this system, the effects of the BAP concentration, immersion time, explant size, number of explants per temporary immersion unit were investigated in the independent experiments using completely randomized design. Temporary immersion system was also compared with the solid and liquid media. The highest shoot number per explant was obtained from a temporary immersion system using an immersion time of 3 min every 150 min (11.8 shoots), a modified MS medium with 6 mg l<sup>-1</sup> BAP (12.33 shoots) and explants with shoot length of 4 cm (12.45 shoots). Results indicated that the temporary immersion system was significantly better than solid and liquid media for the shoot number per explant (10.66, 2.56 and 5.66, respectively).

Keywords: BAP; Explant; Immersion time; Micropropagation; Rosa damascena Mill

### Introduction

For centuries, rose has been regarded as the most popular ornamental plant in the world. Among the numerous species of the Rosaceae family, Rosa damascena Mill., known as Damask rose, is considered as one of the oldest and most valuable varieties. Its aroma and essential oils have therapeutic, sedative, antidepressant, antispasmodic, antioxidizing (Achuthan et al. 2003), antiseptic, antibacterial, antimicrobial (Basim and Basim 2003; Ozkan et al. 2004) and anti-HIV effects (Mahmood et al. 1996). In addition to having astringent agents and playing a role in altering blood cholesterol, it accounts for diverse world wide applications in the manufacture of various products.

Cutting, layering and grafting (Horn 1992) are the common vegetative methods of propagating R. damascena Mill. In some regions, the use of root suckers is the traditional and the most common rose propagation method. However, since it depends on season, sucker production in Damask rose has very low frequency (Mahmoudi Noodezh et al. 2012). Moreover, the methods of clonal propagation can't be fully relied on to attain healthy disease-free plants, thus resulting in slow rates of multiplication and subsequent reduction of production. Hence, the first widely accepted method of propagation of this plant in biotechnology was micropropagation as a popular large-scale of clonal propagation, through which Damask rose could be propagated to produce a

large number of plants within a small physical space and thus save time (Skirvin et al. 1990; Mahmoudi Noodezh et al. 2012) besides ensuring a disease-free plant production. Different features of tissue culture propagation systems for R. damascena have been described by several researchers. Supplementing the media with GA<sub>3</sub> (Kumar et al. 2000), 2.5-3.0 mg 1-1 BA in combination with low rate of IBA (Jabbarzadeh and Khosh-Khui 2005) and 2 mg l<sup>-1</sup> BAP, 0.1 mg l<sup>-</sup> <sup>1</sup>NAA and 100 mg l<sup>-1</sup> phloroglucinol (Salekjalali, 2012) could improve in vitro multiplication of Damask rose. In another research the use of a liquid modified MS medium (with eliminated Cl<sup>-</sup> and reduced NH4<sup>+</sup> ions) caused the best growth of newly proliferated shoots and no aging was occurred (Nikbakht et al. 2005).

Notably, the development of the area under cultivation of Damask rose in some regions is necessary to produce rosewater and essential rose oils, which serve as the highly valuable commercial resource (Mahmoudi Noodezh et al. 2012). Since new large fields are needed annually to produce important crop species with high costs, conventional micropropagation has been limited To reduce commercially. the costs of micropropagation, automation via a bioreactor application has been proven to be an effective method (Aitken-Christie et al. 1995; Levin et al. 1998). Thus, the use of temporary immersion bioreactors (TIBs), which generally reduce the problems of hyperhydricity, poor quality of propagules and transplantation on a solid medium in the elongation and/or rooting stage usually confronted in the permanent liquid cultures, is simple and cost-effective. As a result, several

commercial micropropagation laboratories have started the employment of the aforementioned systems for their productions. Furthermore, a superior mass balance, which resulted based on the use of TIBs, was associated with a higher proliferation rate, improved labor efficiency and reduced cost compared to the conventional micropropagation conducted on a semisolid medium (Teisson and Alvard 1995; Escalona et al. 1999; Etienne et al. 2006). In the TIBs, cultures are temporarily immersed in the medium at specified intervals with a preset duration. A typical design involves the utilization of two plastic or glass vessels: one holds the liquid medium and the other the explants (Adelberg and Simpson 2002). Several studies have been carried out to investigate the use of automated temporary immersion systems to study the effects of environmental conditions, such as nutrient supply and internal atmosphere composition in the culture vessel, on the physiology of some recipient species (Martre et al. 2001; Escalona et al. 2003). The use of temporary immersion system has been reported in other plants such as hybrid chestnut (Vidal et al. 2015), teak (Quiala et al. 2012), pistachio (Akdemir et al.

2014) and peach palm (Steinmacher et al. 2011).

The use of temporary immersion system in Damask

rose micropropagation is yet to be studied.

Therefore, in this research, a traditional temporary

immersion system was designed and thereafter the

effects of BAP concentration, immersion time,

explant length and number of explant per temporary immersion unit were investigated on the

shoot number per explant.

#### **Materials and Methods**

For the micropropagation of Damask rose, a temporary immersion system (T.I.S.), the BIT (Twin Flask System) type (Escalona 1999), was designed and used. The different parts of this system include silicone tubes, air filters, programmable timers, air compressors, solenoid valves and glass vessels. The system applied here included two glass containers (volume, 600 ml), one of which was used for the growing of *in vitro* plantlets while the other served as a liquid medium reservoir (150 ml). The silicone tubes are used as the channel for the transfer of compressed

air between the compressor and the vessels and also the transfer of culture medium between the vessels. The airflow was sterilized when passing through the hydrophobic filters of 0.45  $\mu$ m and the air pressured by an air compressor pushed the medium from one container to the other, so that the explants could be immersed and the shoots could be regenerated. Then, to withdraw the medium from the container of culture, the airflow was reversed. Meanwhile, the programmable timers controlled the immersion period frequency and length (Figure 1).



Figure 1. Temporary immersion system (1: programmable timer, 2: air compressor, 3: solenoid valve, 4: the first glass container for growing plant materials, 5: the second glass container was used as a liquid medium reservoir, 6: air filter)

The explants used in this study were prepared from a single Damask rose (cultivar Azaran) plant grown in a Damask rose garden. The height of the shrub was between 100 and 150 cm, and this shrub had 10 to 20 one-year stems, with each stem having approximately 20 to 25 axillary buds. The nodal explants (about 1.5 cm in length, with one axillary bud) from the mid-stem region of one-year stems (0.4–0.6 cm in diameter) were chosen as explants.

Initially, the leaves subtending each bud were cut at the base of the petiole, the plant materials (stems with 10 cm in length with a maximum of four buds) were then washed with liquid detergent (common dishwashing liquid) diluted in water and

were placed under running tap water for 30 min. The nodal sections were then initially treated with 5.25% sodium hypochlorite (w/v) for 10 min, followed by rinsing with sterile distilled water. Afterward, the buds were rinsed with 70% ethanol for 30 s, followed by rinsing in the sterile distilled water and a further disinfection was done in 5.25% sodium hypochlorite solution (w/v) with five drops of Tween-20 added for 30 min, and this solution was gently stirred with a magnetic stirrer. Finally, the explants were rinsed three times with sterile distilled water (Mahmoudi Noodezh et al. 2012). The base medium was the modified MS medium, A<sub>19</sub> (amount of NH<sub>4</sub>NO<sub>3</sub> and CaCl<sub>2</sub> increased 1.5 times. iron supplementation (FeEDDHA) increased about 20%) supplemented with 4 mg  $l^{-1}$ of BAP, 0.25 mg l<sup>-1</sup> IAA, 50 mg l<sup>-1</sup> ascorbic acid, 50 mg l<sup>-1</sup> citric acid, 3% sucrose, 6 g l<sup>-1</sup> agar-agar and 2 g l<sup>-1</sup>phytagel) (Mahmoudi Noodezh et al. 2012). Glass bottles with autoclave-resistant plastic caps (5.5 cm in diameter, 8 cm in height and 250 ml in volume) containing 50 ml of modified MS medium (pH 5.8) were autoclaved for 20 min (121 °C and 1.2 bars). The cultures with three nodal explants per bottle were incubated at 24±1°C in a controlled growth chamber under a 16-h photoperiod with a light intensity of 3000 lux. After 15 days, for studying the effects of the BAP concentration  $(2, 4, 6 \text{ and } 8 \text{ mg1}^{-1})$ , immersion time (3 min every 90, 150 and 300 min) and the number of explants per temporary immersion unit (3, 4 and 5 explants), in three independent experiments, the cultivated explants together with their regenerated shoot (mean length of the shoots about 1.5-2 cm) were selected and transferred to the temporary immersion system. In another experiment, effect of

the size of the regenerated shoot on each explant was investigated. Therefore, the explants with different shoot size (1, 2, 3 and 4 cm in height) were selected and transferred to the temporary immersion system. Also temporary immersion system was compared with the liquid and solid A<sub>19</sub> media in the same glass bottles. Each experiment was performed as a completely randomized design layout with three replications (each replication consisted of three explants per temporary immersion unit). After 30 days the mean shoot number per explant was recorded. According to our research on previous Damask rose micropropagation (Mahmoudi Noodezh et al. 2012)  $\frac{1}{2}$  A<sub>19</sub> medium supplemented with 2 mg l<sup>-1</sup> of IBA was used for rooting some of the best regenerated shoots in the temporary immersion system (a separated rooting experiment has not been performed). Then the plantlets were transplanted to small plastic pots containing perlite and maintained in a controlled growth chamber at 25±1 °C and high Relative Humidity (RH) under a 16 hr photoperiod. RH was gradually reduced and after about 20 days the plants were transferred to large plastic pots containing perlite and cocopeat (1: 5 v/v) and transferred to a glass greenhouse at 27± 2 °C.

Preliminary statistical analyses such as normality test (Kolmogorov-Smirnov test) and homogeneity of variances (Levene's test) were conducted. After analysis of variance, the means of treatments were compared using Duncan's multiple range procedure. All statistical analyses were carried out using SPSS version 14.

### **Results and Discussion**

The results of analysis of variance indicated a significant difference between the different concentrations of BAP at 1% probability level of for the mean shoot number per explant. The mean comparisons (Figure 2) showed that the use of 6 mg 1<sup>-1</sup> BAP produced the highest mean shoot number per explant (12.33 shoots). There was no significant difference between 4, 6 and 8 mg l<sup>-1</sup> BAP, but the use of 4 mg l<sup>-1</sup> BAP was more suitable due to the production of shoots with better quality and less vitrified tissues. In the solid medium (control) containing these BAP concentrations, the shoots did not grow well enough and these concentrations apparently exhibited poisonous effects. In the temporary immersion system, a high concentration of BAP (6 mg 1-1) did not show poisonous effect due to the temporary contact between the liquid medium and plant material, while there was a permanent contact in the continuous liquid or solid media.

In the second experiment, analysis of variance showed a significant difference between the different immersion times at the 1% probability level for mean shoot number per explant. The comparison of means (Figure 3) indicated that the immersion-treatment of 3 min every 150 min produced the highest mean shoot number per explant (11.86 shoots), while the control (solid medium) had the lowest mean shoot number per explant (2.67 shoots). Every plant species owns a specific immersion time for *in vitro* growth and development, which is a key factor for the use of temporary immersion system in managing the uptake of nutrients of and expression hyperhydricity. A variety of immersion times have been considerably used by different researchers and this is probably due to the presence of a large variety of species, micropropagation processes and temporary immersion systems (Hvoslef-Eide and Preil 2005). Based on immersion time, efficient tuberization of potato and somatic embryo production in coffee and rubber would take a long (1 h every 6 h) and very short (1 min every 12 h) times, respectively (Etienne et al. 1997a, 1997 b), while for the propagation of grapevine shoots, very frequent immersions (30 s every 30 s) prove to be of a high efficiency (Harris and Mason 1983). Immersion frequencies were further reported to be important in the proliferation of service berry shoots. Additionally, hyperhydricity was seen to occur with immersions of 5 min every 30 min, but not 5 min every 60 min (Krueger et al. 1991).

The analysis of variance of the third experiment showed also a significant difference between the treatments for mean shoot number per explant. The mean comparison (Figure 4) showed that the use of explants with shoot length of 4 cm in the temporary immersion system produced the highest mean shoot number per explant (12.45 shoots) and explants with shoot length of 1 cm had the lowest mean shoot number per explant (8.33 shoots). The comparison of means showed that the explants with more length produced more shoots in the temporary immersion system. The explant length was directly correlated with the number of

# Journal of Plant Physiology and Breeding

ISSN: 2008-5168

2016, 6(2): 9-18





Figure 2. Mean comparison of BAP concentrations in Damask rose micropropagation using temporary immersion system in terms of the mean shoot number per explant



Figure 3. Mean comparison of the different immersion times for the mean shoot number per explant in Damask rose micropropagation using temporary immersion system



Figure 4. Mean comparison for the explants with different shoot lengths in terms of the mean shoot number per explant in Damask rose micropropagation using temporary immersion system

regenerated shoots. The longer and stronger explants had the higher number of lateral buds, as well as bigger lateral buds. In a study, it was shown that initial explants have a positive effect on growth and reproduction rates in cacao (Traore *et al.* 2003).

In the fourth experiment, there was no significant difference among different number of explants (3, 4 and 5 explants) per temporary immersion unit. This can be an important advantage for this propagation system, since more explant number will produce more regenerated shoots.

The analysis of variance for the comparison of the temporary immersion system with the solid and liquid media showed that there was a significant difference between the treatments. The immersion system (Figure 5) produced the highest mean shoot number per explant (10.66 shoots), while the solid medium had the lowest mean shoot number per explant (2.56 shoots).

In general, these results showed that the temporary immersion system is a suitable method for Damask rose micropropagation. In this method, the highest mean shoot number per explant was obtained using 6 mg l<sup>-1</sup> BAP (12.33 shoots per explant), immersion time of 3 min in every 150 min (11.8 shoots per explant) and explants with shoot length of 4 cm (12.45 shoots per explant). Compared agar-based to cultures. plant micropropagation in a temporary immersion system provides several advantages, including better control of plant tissue-culture medium contact, aeration, medium circulation and culture scaling-up and optimal supply of nutrients and growth regulators. In conclusion, these results showed that the temporary immersion system is a potential method used in the micropropagation of Damask rose, which resulted in significant improvements regeneration of shoot and



Figure 5. Mean comparison of the temporary immersion, solid and liquid media for the mean shoot number per explant in Damask rose micropropagation



Figure 6) a: *In vitro* shoot proliferation of Damask rose in the temporary immersion system, b: The shoots regenerated from one explant in the temporary immersion system

proliferation. The optimum micropropagation in the temporary immersion system will depend on a better understanding of plant responses to every specific microenvironment and on specific culture manipulation to control the morphogenesis of regenerated plantlets in this propagation method.

## References

- Achuthan CR, Babu BH and Padikkala, 2003. Antioxidant and hepatoprotective effects *of Rosa damascena*. Pharmaceutical Biology 41: 357-361
- Adelberg JW and Simpson EP, 2002. Intermittent immersion vessel apparatus and process for plant propagation. Internl. S/N: PCT/US01/06586
- Aitken-Christie J, Kozai T and Takayama S, 1995. Automation in plant tissue culture. General introduction and overview. In: Aitken-Christie *et al.* (Eds). Automation and Environment Control in Plant Tissue Culture. Pp. 1-18. Kluwer Acad. Publ., Dordrecht, the Netherlands.
- Akdemir H, Su<sup>°</sup>zerer V, Onay A, Tilkat E, Ersali Y and Ciftci YO, 2014. Micropropagation of the pistachio and its rootstocks by temporary immersion system. Plant Cell Tissue and Organ Culture 117: 65–76.
- Basim E and Basim H, 2003. Antibacterial activity of Rosa damascena essential oil. Fitoterapia 74: 394-396
- Escalona M, Lorenzo JC, González B, Daquinta M, González JL, Desjardins Y and Borroto CG, 1999. Pineapple (*Ananas comosus* L. Merr) micropropagation in temporary immersion systems. Plant Cell Reports 18: 743-748.
- Escalona M, Samson G, Borroto C and Desjardins Y, 2003. Physiology of effects of temporary immersion bioreactors on micropropagated pineapple plantlets. *In Vitro* Cellular and Developmental Biology-Plant 39: 651-656.
- Etienne H, Bertrand B, Anthony F, Côte F and Berthouly M, 1997a. L'embryogenèse somatique: un outil pour l'amélioration génétique du caféier. In: 17<sup>th</sup> International Scientific Colloquium on Coffee. 21-25 June, Nairobi, Kenya. Pp. 457–465. ASIC Publishers, Paris, France.
- Etienne H, Dechamp E, Etienne BD and Bertrand B, 2006. Bioreactors in coffee micropropagation. Brazilian Journal of Plant Physiology 18 (1): 45-54.
- Etienne H, Lartaud M, Michaux–Ferrière N, Carron MP, Berthouly M and Teisson C, 1997b. Improvement of somatic embryogenesis in *Hevea brasiliensis* (Müll. Arg.) using the temporary immersion technique. *In Vitro* Cellular and Developmental Biology-Plant 33: 81–87.
- Harris RE and Mason EB, 1983. Two machines for *in vitro* propagation of plants in liquid media. Canadian Journal of Plant Science 63: 311–316.
- Horn WAH, 1992. Micropropagation of rose. In: Bajaj YPS (Ed). Biotechnology in Agriculture and Forestry. Vol. 20. High-tech and Micropropagation IV. Pp. 320-342. Springer, Germany.
- Hvoslef-Eide AK and Preil W, 2005. Liquid Culture Systems for In Vitro Plant Propagation. Springer, Dordrecht.
- Jabbarzadeh Z and Khosh-Khui M, 2005. Factors affecting tissue culture of Damask rose (*Rosa damascene* Mill.). Scientia Horticulturae 105: 475-482.
- Krueger S, Robacker C and Simonton W, 1991. Culture of *Amelanchier* × *grandiflora*in a programmable micropropagation apparatus. Plant Cell Tissue and Organ Culture 27: 219–226.
- Kumar A, Sood A, Palni UT, Gupta AK and Palni LMS, 2001. Micropropagation of *Rosa damascene* Mill. from mature bushes using thidiazuron. The Journal of Horticultural Science and Biotechnology 76: 30-34.
- Levin R, Stav R, Alper Y and Watad AA, 1998. A technique for repeated non-axenic subculture of plant tissues in a bioreactor on liquid medium containing sucrose. Plant Tissue Culture Biotechnology 3: 41–45.
- Mahmood N, Piacente S, Pizza C, Burke A, Khan A and Hay A, 1996. The anti-HIV activity and mechanisms of action of pure compounds isolated from *Rosa damascene*. Biochemical and Biophysical Research Communications 229: 73–79.
- Mahmoudi Noodezeh H, Moieni A and Baghizadeh F, 2012. *In vitro* propagation of the Damask rose (*Rosa damascena* Mill.). *In vitro* Cellular and Development Biology-Plant 48: 530-538.
- Martre P, Lacan D, Just D and Teisson C, 2001. Physiological effects of temporary immersion on *Hevea* brasiliensis (Müll. Arg.) callus. Plant Cell Tissue and Organ Culture 67: 25–35.
- Nikbakht A, Kafi M, Mirmasomi M and Babalar M, 2005. Micropropagation of Damask Rose (*Rosa damascene* Mill.) cvs. Azaran and Ghamsar. International Journal of Agriculture and Biology 7: 535-538.
- Ozkan G, Sagdic O, Baydar NG and Baydar H, 2004. Antioxidant and antibacterial activities of *Rosa damascene* flower extracts. Food Science and Technology International 10: 277-281.

18	Bosh <i>et al</i> .	2016, 6(2): 9-18

- Quiala E, Can<sup>°</sup>al MJ, Meijo'n M, Rodriguez R, Chave'z M, Valledon L, Feria M and Barbo'n R, 2012. Morphological and physiological responses of proliferating shoots of teak to temporary immersion and BA treatments. Plant Cell Tissue and Organ Culture 109: 223–234.
- Salekjalali M, 2012. Phloroglucinol, BAP and NAA enhance axillary shoot proliferation and other growth indicators in *in vitro* culture of Damask Rose (*Rosa damascene* Mill.). American-Eurasian Journal of Agricultural and Environmental Sciences 12 (7): 960 -966.
- Skirvin RM, Chu MC and Young HJ, 1990. Rose. In: Ammirato PV, Evans D, Sharp WR and Bajaj YPS (Eds). Handbook of Plant Cell Culture. Vol. 5. Ornamental species. Pp. 716-743. McGraw Hill, New York, USA.
- Steinmacher DA, Guerra MP, Saare-Surminski K and Lieberei R, 2011. A temporary immersion system improves *in vitro* regeneration of peach palm through secondary somatic embryogenesis. Annals of Botany 108: 1463–1475.
- Teisson C and Alvard D, 1995. A new concept of plant in vitro cultivation in liquid medium: temporary immersion. In: Terzi M, Cella R and Falavigna A (Eds). Current Issues in Plant Molecular and Cellular Biology. Pp. 105-109. Kluwer Academic Publisher.
- Traoré A, Maximova SN and Guiltinan MJ, 2003. Micropropagation of *Theobroma cacao* L. using somatic embryo-derived plants. *In vitro* Cellular and Development Biology-Plant 409: 1-7.
- Vidal N, Correa B, Rial E, Regueira M, Sa'nchez C and Cuenca B, 2015. Comparison of temporary and continuous immersion systems for micropropagation of axillary shoots of chestnut and willow. Acta Horticulturae 1083: 227–233.