**Open Access** 

# Approximate symmetry reduction to the perturbed coupled KdV equations derived from two-layer fluids

Xin Li and Suping Qian<sup>\*</sup>

#### Abstract

**Purpose:** The purpose of this paper is to research the solution of the perturbed coupled Korteweg-de Vries (KdV) equations.

**Methods:** The results are obtained through researching a system of corresponding partial differential equations which approximates the perturbed coupled KdV equations and the symmetry method.

**Results:** The symmetry structure and the corresponding general approximate symmetry reduction are derived. Then, using the symmetry reduction, the infinite series solutions are obtained.

**Conclusions:** The approximate symmetry reduction approach can be used to search for similar results of other perturbed nonlinear differential equations with small parameters.

Keywords: Coupled KdV equations, Approximate symmetry reduction, Series reduction solutions

#### Background

It is well known that nonlinear dynamical systems emerge in various fields, such as fluid mechanics, plasma physics, biology, hydrodynamics and optical fibers. The nonlinear phenomena which occurred in such fields are often related to some nonlinear wave equations. In general, one often chooses to exactly solve the related unperturbed partial differential systems as the beginning point of research. However, many nonlinear partial differential equations (PDEs) are, of course, approximations in their derivation, i.e. higher order terms, and dissipation may have been neglected. In order to better understand such phenomena as well as further apply them in practical scientific research, it is important to study the properties of the perturbed PDEs. The approximate symmetry perturbation approach [1] is a powerful method for such substitutions and has been applied to many classical PDEs [2-7]. Maybe in theory it is not difficult, however, the concrete study shows that it, indeed, is not a trivial work to construct the different-order perturbation solutions and even

\*Correspondence: qsp@cslg.edu.cn

School of Mathematics and Statistics, Changshu Institute of Technology, Changshu, Jiangsu, 215500, China

give the general formulae. During the computations, one must make appropriate choices not only on the type of the solution functions but also on the parameters which occurred during the computations, such as some integral constants, to render the solutions physically meaningful and have correct asymptotic forms. Next, we take the coupled Korteweg-de Vries (KdV) equations [8,9] as an example, which reads

$$u_t + u_{xxx} + 2uu_x + (uv)_x = 0, (1a)$$

$$v_t + v_{xxx} + 2vv_x + (uv)_x = 0.$$
 (1b)

The model equation (Equation 1) is important to describe many kinds of physical systems and atmosphere [10], especially, for a two-layer fluid system. To describe the real two-layer fluid system, we have to introduce some types of physical effects, say, the viscosity. In this letter, we consider the above coupled KdV equations

$$u_t + u_{xxx} + 2uu_x + (uv)_x + \epsilon u_{xx} = 0,$$
 (2a)

$$v_t + v_{xxx} + 2vv_x + (uv)_x + \epsilon v_{xx} = 0.$$
 (2b)

## Springer

#### in terms of approximate symmetry reduction method.

© 2012 Li and Qian; licensee Springer. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/2.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

#### **Results and discussion**

Substituting Equation 14 into Equation 15 and eliminating  $u_{jt}$  and  $v_{jt}$   $(j = 0, 1, \cdots)$  in terms of Equation 13 result in determining equations by vanishing all coefficients of different partial derivatives of  $u_j$  and  $v_j$  for the unknown functions X, T,  $U_j$  and  $V_j$   $(j = 0, 1, \cdots)$ , which have the solution

$$X = ax + x_0, \ T = 3at + t_0, \ U_0 = (-2a - b)v_0 + cu_0,$$
  

$$V_0 = bv_0 + (-c - 2a)u_0, \ U_1 = c_1u_0 + (a + c)u_1$$
  

$$+ d_1v_0 - (2a + b)v_1, \ V_1 = -c_1u_0 - (c + 2a)u_1$$
  

$$- d_1v_0 + (a + b)v_1, \ U_2 = c_2u_0 + c_1u_1$$
  

$$+ (c + 2a)u_2 + d_2v_0 + d_1v_1 - (2a + b)v_2,$$
  

$$V_2 = -c_2u_0 - c_1u_1 - (c + 2a)u_2 - d_2v_0 - d_1v_1$$
  

$$+ (2a + b)v_2, \ \cdots \ \cdots ,$$
  

$$U_j = (c + ja)u_j - (2a + b)v_j + \sum_{i=0}^{j-1} (c_{j-i}u_i + d_{j-i}v_i),$$
  

$$V_j = -(c + 2a)u_j + (b + ja)v_j$$
  

$$- \sum_{i=0}^{j-1} (c_{j-i}u_i + d_{j-i}v_i) \cdots,$$
  
(3)

where *a*, *b*, *c*,  $x_0$ ,  $t_0$ ,  $c_j$  and  $d_j$  are arbitrary constants. The sum items in the general formula of  $U_j$  and  $V_j$  are, respectively, the turn of  $u_j$  and  $v_j$ , so we just suppose  $c_j = 0$  and  $d_j = 0$ . Subsequently, solving the characteristic equations

$$\frac{dx}{X} = \frac{dt}{T} = \frac{du_0}{U_0} = \frac{du_1}{U_1} = \frac{du_2}{U_2} = \dots = \frac{du_j}{U_j} = \dots$$
(4)

leads to the similarity solutions to Equation 2 which can be distinguished in the following two subcases:

**Case 1.** When  $a \neq 0$ , and to be convenient, we let  $x_0 = 0$  and  $t_0 = 0$ , the similarity solutions are

$$\begin{split} u_0 &= \frac{(2a+b)Q_0(\xi)}{t^{\frac{2}{3}}} + P_0(\xi)t^{\frac{b+c+2*a}{3a}},\\ v_0 &= \frac{(c+2a)Q_0(\xi)}{t^{\frac{2}{3}}} - P_0(\xi)t^{\frac{b+c+2a}{3a}},\\ u_1 &= t^{\frac{b+c+3a}{3a}}P_1(\xi) + \frac{(2a+b)Q_1(\xi)}{t^{\frac{1}{3}}},\\ v_1 &= -t^{\frac{b+c+3a}{3a}}P_1(\xi) + \frac{(2a+c)Q_1(\xi)}{t^{\frac{1}{3}}},\\ u_2 &= t^{\frac{b+c+4a}{3a}}P_2(\xi) + (2a+b)Q_2(\xi), \end{split}$$

$$\begin{aligned}
\nu_{2} &= -t^{\frac{b+c+4a}{3a}} P_{2}(\xi) + (2a+c)Q_{2}(\xi), \\
u_{3} &= t^{\frac{b+c+5a}{3a}} P_{3}(\xi) + (2a+b)Q_{3}(\xi)t^{\frac{1}{3}}, \\
\nu_{3} &= -t^{\frac{b+c+5a}{3a}} P_{3}(\xi) + (2a+c)Q_{3}(\xi)t^{\frac{1}{3}}, \\
\dots &\dots \\
u_{j} &= t^{\frac{b+c+(j+2)a}{3a}} P_{j}(\xi) + (2a+b)Q_{j}(\xi)t^{\frac{j-2}{3}}, \\
\nu_{j} &= -t^{\frac{b+c+(j+2)a}{3a}} P_{j}(\xi) + (2a+c)Q_{j}(\xi)t^{\frac{j-2}{3}}, \\
\dots &\dots &\dots \end{aligned}$$
(5)

with the similarity variable  $\xi = \frac{x}{t^{\frac{1}{3}}}$ . Hence, the perturbation series solution to Equation 2 is

$$u = \sum_{j=0}^{\infty} \epsilon^{j} \left[ t^{\frac{b+c+(j+2)a}{3a}} P_{j}(\xi) + (2a+b)Q_{j}(\xi)t^{\frac{j-2}{3}} \right],$$
(6a)
$$v = \sum_{j=0}^{\infty} \epsilon^{j} \left[ -t^{\frac{b+c+(j+2)a}{3a}} P_{j}(\xi) + (2a+c)Q_{j}(\xi)t^{\frac{j-2}{3}} \right].$$
(6b)

The similarity reduction equations related to similarity solution equations are

$$O(\epsilon^{0}): P_{0\xi\xi\xi} = A(P_{0}Q_{0})_{\xi} + \frac{\xi P_{0\xi}}{3} - \frac{(b+c+2a)P_{0}}{3a},$$
(7a)

$$Q_{0\xi\xi\xi} = BQ_0Q_{0\xi} + \frac{\xi Q_{0\xi}}{3} + \frac{2}{3}Q_0,$$
 (7b)

$$O(\epsilon^{1}): P_{1\xi\xi\xi} = A \sum_{i=0}^{1} (Q_{i}P_{1-i})_{\xi} + \frac{\xi P_{1\xi}}{3} + \frac{(c+b+3a)P_{1}}{3a} - P_{0\xi\xi}, \quad (7c)$$

$$Q_{1\xi\xi\xi} = B \sum_{i=0}^{1} \frac{1}{2} (Q_i Q_{1-i})_{\xi} + \frac{\xi Q_{1\xi}}{3} + \frac{\xi Q_1}{3} - Q_{0\xi\xi},$$
(7d)

$$O(\epsilon^{2}): P_{2\xi\xi\xi} = A \sum_{i=0}^{2} (Q_{i}P_{2-i})_{\xi} + \frac{\xi P_{2\xi}}{3} + \frac{(c+b+4a)P_{2}}{3a} - P_{1\xi\xi},$$
(7e)

$$Q_{2\xi\xi\xi} = B \sum_{i=0}^{2} \frac{1}{2} (Q_i Q_{2-i})_{\xi} + \frac{\xi Q_{2\xi}}{3} - Q_{1\xi\xi},$$
(7f)

$$O(\epsilon^{3}): P_{3\xi\xi\xi} = A \sum_{i=0}^{3} (Q_{i}P_{3-i})_{\xi} + \frac{\xi P_{3\xi}}{3} + \frac{(c+b+5a)P_{3}}{3a} - P_{2\xi\xi}, \qquad (7g)$$

... ... ...,

$$Q_{3\xi\xi\xi} = B \sum_{i=0}^{\infty} \frac{1}{2} (Q_i Q_{3-i})_{\xi} + \frac{\varsigma Q_{3\xi}}{3} - \frac{1}{3} Q_3 - Q_{2\xi\xi}, \qquad (7h)$$

$$O(\epsilon^{j}): P_{j\xi\xi} = A \sum_{i=0}^{j} (Q_{i}P_{j-i})_{\xi} + \frac{\xi P_{j\xi}}{3} + \frac{(c+b+(j+2)a)P_{j}}{3a} - P_{j-1\xi\xi}, \quad (7i)$$
$$Q_{j\xi\xi\xi} = B \sum_{i=0}^{j} \frac{1}{2} (Q_{i}Q_{j-i})_{\xi} + \frac{\xi Q_{j\xi}}{3} - \frac{j-2}{3} Q_{j} - Q_{j-1\xi\xi}, \quad (7j)$$
$$\dots \dots \dots ,$$

where A = -b - c - 4a and B = -2b - 2c - 8a are integral constants, and  $P_{-1} = Q_{-1} = 0$ .

**Case 2. (1).** a = 0,  $b + c \neq 0$ : We let  $t_0 = 1$ , then the similarity solutions are

$$u_{j} = e^{(b+c)t}F_{j}(\xi) + bG_{j}(\xi), v_{j} = -e^{(b+c)t}F_{j}(\xi) + cG_{j}(\xi)$$
(8)

with the similarity variable  $\xi = x - x_0 t$ , where  $F_j$  and  $G_j$  (j = 0, 1, ...) are arbitrary functions of  $\xi$ . Accordingly, the similarity reduction equations related to similarity solution equations are

$$O(\epsilon^{0}): F_{0\xi\xi\xi} = -(b+c)(F_{0}G_{0})_{\xi} + x_{0}F_{0\xi} - (b+c)F_{0},$$
(9a)

$$G_{0\xi\xi\xi} = -2(b+c)G_0G_{0\xi} + x_0G_{0\xi}$$
 (9b)

$$O(\epsilon^{1}): F_{1\xi\xi\xi} = -(b+c)(F_{0}G_{1} + F_{1}G_{0})_{\xi} + x_{0}F_{1\xi} - (b+c)F_{1} - F_{0\xi\xi},$$
(9c)  
$$G_{1\xi\xi\xi} = -2(b+c)(G_{0}G_{1})_{\xi} + x_{0}G_{1\xi} - G_{0\xi\xi},$$

$$G_{1\xi\xi\xi} = -2(\delta + \ell)(G_0G_1)_{\xi} + x_0G_{1\xi} - G_{0\xi\xi},$$
(9d)

$$O(\epsilon^{2}): F_{2\xi\xi\xi} = -(b+c) \sum_{i=0}^{2} (F_{i}G_{2-i})_{\xi} + x_{0}F_{2\xi}$$
$$-(b+c)F_{2} - F_{1\xi\xi}, \qquad (9e)$$

$$G_{2\xi\xi\xi} = -2(b+c)(G_0G_2)_{\xi}$$

$$-2(b+c)G_1G_{1\xi} + x_0G_{2\xi} - G_{1\xi\xi},$$
(9f)
$$O(\epsilon^3) : F_{3\xi\xi\xi} = -(b+c)\sum_{i=0}^3 (F_iG_{3-i})_{\xi} + x_0F_{3\xi}$$

$$-(b+c)F_3 - F_{2\xi\xi},$$
(9g)
$$G_{3\xi\xi\xi} = -2(b+c)(G_1G_2 + G_0G_3)_{\xi}$$

$$+ x_0G_{3\xi} - G_{2\xi\xi},$$
(9h)

$$O(\epsilon^{j}): F_{j\xi\xi\xi} = -(b+c) \sum_{i=0}^{j} (F_{i}G_{j-i})_{\xi} + x_{0}F_{j\xi}$$
  
-  $(b+c)F_{j} - F_{j-1\xi\xi},$  (9i)  
$$G_{j\xi\xi\xi} = -(b+c) \sum_{i=0}^{j} (G_{i}G_{j-i})_{\xi} + x_{0}G_{j\xi}$$
  
-  $G_{j-1\xi\xi},$  (9j)

... ... ....

with  $F_{-1} = 0$  and  $G_{-1} = 0$ . (2). a = 0, b + c = 0: In this case, the similarity solutions are

$$u_j = F_j(\xi), \ v_j = G_j(\xi)$$
 (10)

with the similarity variable  $\xi = x - mt$ , where *m* is an arbitrary constant, and here,  $F_j$  and  $G_j$  ( $j = 0, 1, \dots$ .) are also arbitrary functions of  $\xi$ . The reduction equations related to similarity solution equations are

$$O(\epsilon^0): F_{0\xi\xi\xi} = mF_{0\xi} - (2F_0 + G_0)F_{0\xi} - F_0G_{0\xi},$$
(11a)

$$G_{0\xi\xi\xi} = mG_{0\xi} - (F_0 + 2G_0)G_{0\xi} - F_{0\xi}G_0,$$
(11b)

$$O(\epsilon^{1}): F_{1\xi\xi\xi} = mF_{1\xi} - (2F_{0} + G_{0})F_{1\xi}$$
  
- (2F\_{1} + G\_{1})F\_{0\xi} - F\_{0\xi\xi} - F\_{0}G\_{1\xi}  
- F\_{1}G\_{0\xi}, (11c)  
$$G_{1\xi\xi\xi} = mG_{1\xi} - (F_{0} + 2G_{0})G_{1\xi}$$
  
- (F\_{1} + 2G\_{1})G\_{0\xi} - G\_{0\xi\xi}  
- F\_{1\xi}G\_{0} - F\_{0\xi}G\_{1}, (11d)

$$O(\epsilon^{2}): F_{2\xi\xi\xi} = mF_{2\xi} - F_{1\xi\xi}$$
$$-\sum_{i=0}^{2} [(F_{2-i} + G_{2-i})G_{i}]_{\xi}, \quad (11e)$$
www.SID.ir

$$G_{2\xi\xi\xi} = mG_{2\xi} - G_{1\xi\xi} - \sum_{i=0}^{2} [(F_i + G_i)G_{2-i}]_{\xi}, \quad (11f)$$

$$O(\epsilon^{3}): F_{3\xi\xi\xi} = mF_{3\xi} - F_{2\xi\xi}$$
  
-  $\sum_{i=0}^{3} [(F_{3-i} + G_{3-i})G_{i}]_{\xi}$ , (11g)  
 $G_{3\xi\xi\xi} = mG_{3\xi} - G_{2\xi\xi}$   
-  $\sum_{i=0}^{3} [(F_{i} + G_{i})G_{3-i}]_{\xi}$ , (11h)

$$O(\epsilon^{j}): F_{j\xi\xi\xi} = mF_{j\xi} - F_{j-1\xi\xi} - \sum_{i=0}^{j} [(F_{j-i} + G_{j-i})G_{i}]_{\xi}, \quad (11i) G_{j\xi\xi\xi} = mG_{j\xi} - G_{j-1\xi\xi} - \sum_{i=0}^{j} [(F_{i} + G_{i})G_{j-i}]_{\xi}, \quad (11j)$$

with  $F_{-1} = 0$  and  $G_{-1} = 0$ .

#### Conclusions

In summary, by applying the approximate symmetry reduction approach to perturbed coupled KdV equations which is an important physical model to describe many kinds of physical problems especially related to the twolayer fluids with unavoidable viscosity, we have unearthed that the similarity reduction solutions and similarity equations of different orders are coincident in their forms. Therefore, we summarize the infinite series solutions and general formulae of the similarity equations.

Nonetheless, we have not mentioned the convergence of infinite series solutions because of its difficulty. Actually, the convergence radius of the series is dependent on the concrete exact solutions. However, it is still very difficult to get exact solutions for infinitely many reduction equations. More details on this topic should be studied further. The approximate symmetry reduction approach can be used to search for similar results of other perturbed nonlinear differential equations with small parameters, and it is worthwhile to summarize a general principle for the perturbed nonlinear differential equations holding analogous results.

#### Methods

### Approximate symmetry reduction approach to the coupled KdV equations

With the perturbation theory, solutions of perturbed partial differential equations can be expressed as a series containing a small parameter. Specifically, for Equation 2, the solution can be supposed to be of the general form

$$u = \sum_{j=0}^{\infty} \epsilon^{j} u_{j}, v = \sum_{j=0}^{\infty} \epsilon^{j} v_{j},$$
(12)

with  $u_j$  being functions of x and t. Substituting Equation 12 into Equation 2 and vanishing the coefficients of  $\epsilon$ , we obtain a system of partial differential equations

$$O(\epsilon^0): \ u_{0t} + u_{0xxx} + 2u_0u_{0x} + (u_0v_0)_x = 0,$$
 (13a)

$$v_{0t} + v_{0xxx} + 2v_0v_{0x} + (u_0v_0)_x = 0,$$
 (13b)

$$D(\epsilon^{1}): u_{1t} + u_{1xxx} + u_{0xx} + (u_{0}v_{1})_{x} + (u_{1}v_{0})_{x} + 2(u_{0}u_{1})_{x} = 0,$$
(13c)

$$\nu_{1t} + \nu_{1xxx} + \nu_{0xx} + (\nu_0 u_1)_x + (\nu_1 u_0)_x + 2(\nu_0 \nu_1)_x = 0,$$
(13d)

$$O(\epsilon^{2}): u_{2t} + u_{2xxx} + \left(\sum_{i=0}^{2} u_{i}u_{2-i}\right)_{x} + \left(\sum_{i=0}^{2} u_{i}v_{2-i}\right)_{x} + u_{1xx} = 0,$$
(13e)

$$2t + v_{2xxx} + \left(\sum_{i=0}^{2} v_i v_{2-i}\right)_x + \left(\sum_{i=0}^{2} u_i v_{2-i}\right)_x + v_{1xx} = 0,$$
(13f)

v

$$O(\epsilon^{j}): u_{jt} + u_{jxxx} + \left(\sum_{i=0}^{j} u_{i}u_{j-i}\right)_{x} + \left(\sum_{i=0}^{j} u_{i}v_{j-i}\right)_{x} + u_{j-1xx} = 0,$$
(13g)

$$v_{jt} + v_{jxxx} + \left(\sum_{i=0}^{j} v_i v_{j-i}\right)_x + \left(\sum_{i=0}^{j} u_i v_{j-i}\right)_x + v_{j-1xx} = 0,$$
(13h)

where we just let  $u_{-1} = 0$  and  $v_{-1} = 0$ . The next crucial step is to study the symmetry reduction of the above system. To that end, we search for the Lie point symmetries in the form

$$\sigma_j = X u_{jx} + T u_{jt} - U_j, \ (j = 0, 1, \cdots),$$
 (14a)

$$\sigma'_{j} = X v_{jx} + T v_{jt} - V_{j}, \ (j = 0, 1, \cdots),$$
(14b)

www.SID.ir

where *X*, *T*,  $U_j$  and  $V_j$  are functions with respect to *x*, *t*,  $u_j$  and  $v_j$  ( $j = 0, 1, \cdots$ ). The linearized equations for Equation 2 are

$$\sigma_{0t} + \sigma_{0xxx} + (\nu_0 \sigma_0)_x + (u_0 \sigma_0')_x + 2 (\sigma_0 u_0)_x = 0,$$

$$\sigma_{0t}' + \sigma_{0xxx}' + (\nu_0 \sigma_0)_x + (u_0 \sigma_0')_x + 2(\sigma_0' \nu_0)_x = 0,$$
(15b)

$$\sigma_{1t} + \sigma_{1xxx} + \sigma_{0xx} + (\nu_0 \sigma_1 + \nu_1 \sigma_0)_x + (u_0 \sigma_1' + u_1 \sigma_0')_x + 2 (u_0 \sigma_1 + u_1 \sigma_0)_x = 0,$$
(15c)

$$\sigma_{1t}' + \sigma_{1xxx}' + \sigma_{0xx}' + (\nu_0 \sigma_1 + \nu_1 \sigma_0)_x + (u_0 \sigma_1' + u_1 \sigma_0')_x + 2 (\nu_0 \sigma_1' + \nu_1 \sigma_0')_x = 0,$$
(15d)

$$\sigma_{2t} + \sigma_{2xxx} + \sigma_{1xx} + \left(\sum_{i=0}^{2} v_i \sigma_{2-i}\right)_x + \left(\sum_{i=0}^{2} u_i \sigma'_{2-i}\right)_x$$

$$+2\left(\sum_{i=0}^{\infty}u_i\sigma_{2-i}\right)_x=0,$$
(15e)

$$\sigma_{2t}' + \sigma_{2xxx}' + \sigma_{1xx}' + \left(\sum_{i=0}^{2} v_i \sigma_{2-i}\right)_x + \left(\sum_{i=0}^{2} u_i \sigma_{2-i}'\right)_x + 2\left(\sum_{i=0}^{2} v_i \sigma_{2-i}'\right)_x = 0,$$
(15f)

... ... ...

$$\sigma_{jt} + \sigma_{jxxx} + \sigma_{j-1xx} + \left(\sum_{i=0}^{j} v_i \sigma_{j-i}\right)_x + \left(\sum_{i=0}^{j} u_i \sigma_{j-i}'\right)_x + 2\left(\sum_{i=0}^{j} u_i \sigma_{j-i}\right)_x = 0, \quad (15g)$$
$$\sigma_{jt}' + \sigma_{jxxx}' + \sigma_{j-1xx}' + \left(\sum_{i=0}^{j} v_i \sigma_{j-i}\right)_x + \left(\sum_{i=0}^{j} u_i \sigma_{j-i}'\right)_x$$

$$+2\left(\sum_{i=0}^{\infty}\nu_i\sigma'_{j-i}\right)_x = 0, \tag{15h}$$

... ... ...,

which means that Equation 2 is invariant under the transformations  $u_j \rightarrow u_j + \epsilon' \sigma_j$  and  $v_j \rightarrow v_j + \epsilon' \sigma'_j$   $(j = 0, 1, \dots)$  with an infinitesimal parameter  $\epsilon'$ .

#### **Competing interests**

The authors declare that they have no competing interests.

#### Authors' contributions

XL drafted the manuscript. SP conceived the study and participated in its design and coordination. Both authors read and approved the final manuscript.

#### Authors' information

XL, with a Master's degree, is a teacher while SP, with a Doctor's degree, is a professor in Changshu Institute of Technology.

#### Acknowledgements

The work was supported by the Special Funds of the National Natural Science Foundation of China (Grant No. 11141003). The authors are indebted to Prof. SY Lou for his helpful discussions.

#### Received: 6 April 2012 Accepted: 7 June 2012 Published: 18 July 2012

#### References

(15a)

- Fushchich, WI, Shtelen, WM: On approximate symmetry and approximate solutions of the nonlinear wave equation with a small parameter. J. Phys. A: Math. Gen. 22, L887–890 (1989)
- Gao, G: A theory of interaction between dissipation and dispersion of turbulence. Sci. Sinica (Ser A). 28, 616–627 (1985)
- Malfliet, W, Hereman W: The tanh method: II. Perturbation technique for conservative systems. Phys. Scr. 54, <u>56</u>9–575 (1996)
- Marchant, TR, Smyth, NF: Pulse evolution for marangoni-bénard convection. Math. Comput. Model. 28, 45–58 (1998)
- Allen, MA, Rowlands, G: A solitary-wave solution to a perturbed KdV equation. J. Plasma Phys. 64, 475–480 (2000)
- Liu, XZ: Approximate similarity reduction for perturbed Kaup-Kupershmidt equation via Lie symmetry method and direct method. Commun. Theor. Phys. 54, 797–802 (2010)
- Liu, P, Li, ZL, Lou, SY: A class of coupled nonlinear Schrodinger equations: Painlevé property, exact solutions, and application to atmospheric gravity waves. Appl. Math. Mech. **31**, 1383–1404 (2010)
- Lou, SY, Tong, B, Hu, HC, Tang, XY: Coupled KdV equations derived from two-layer fluids. J. Phys. A: Math. Gen. 39, 513–527 (2006)
- Qian, SP, Tian, LX: Modification of the Clarkson-Kruscal's direct method for a coupled system. Chin. Phys. Lett. 24, 2720–2723 (2007)
- 10. Dong, ZZ, Huang, F, Chen, Y: Symmetry reductions and exact solutions of the two-layer model in atmosphere. Z. Naturforch. **66a**, 75–86 (2011)

#### doi:10.1186/2251-7456-6-13

**Cite this article as:** Li and Qian: Approximate symmetry reduction to the perturbed coupled KdV equations derived from two-layer fluids. *Mathematical Sciences* 2012 **6**:13.

## Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- ► Open access: articles freely available online
- ► High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at > springeropen.com