

ORIGINAL RESEARCH

Open Access

# On some structures of soft topology

Bashir Ahmad<sup>1,2</sup> and Sabir Hussain<sup>3,4\*</sup>

## Abstract

In this paper, we define soft exterior and study its basic properties. We establish several important results relating soft interior, soft exterior, soft closure, and soft boundary in soft topological spaces. Moreover, we characterize soft open sets, soft closed sets, and soft clopen sets via soft boundary. All these findings will provide a base to researchers who want to work in the field of soft topology and will help to establish a general framework for applications in practical fields.

**Keywords:** Soft set, Soft topology, Soft open(closed) set, Soft nbd, Soft interior(exterior), Soft closure(boundary)

## Introduction

In 1999, Molodtsov [1] initiated the theory of soft sets as a new mathematical tool to deal with uncertainties while modelling the problems in engineering, physics, computer science, economics, social sciences, and medical sciences. In [2], Molodtsov et al. successfully applied soft sets in directions such as smoothness of functions, game theory, operations research, Riemann integration, Perron integration, probability, and theory of measurement. Maji et al. [3,4] gave the first practical application of soft sets in decision-making problems. In 2003, Maji et al. [4] defined and studied several basic notions of the soft set theory. In 2005, Pei and Miao [5] and Chen [6] improved the work of Maji et al. [3,4].

Many researchers have contributed towards the algebraic structure of the soft set theory [7-19]. The application of the soft set theory in algebraic structures was introduced by Aktas and Cagman [8]. They established the basic notions of soft groups as a generalization of the idea of fuzzy groups. Jun [9] investigated BCK/BCI algebras and studied their applications in ideal theory. Feng et al. [20] worked on soft semirings, soft ideals, and idealistic soft semirings. Ali et al. [21] and Shabir and Irfan Ali [22] studied semigroups and soft ideals over a semigroup which characterized generalized fuzzy ideals and fuzzy ideals with thresholds of a semigroup.

Recently, in 2011, Shabir and Naz [23] and Cagman et al. [24] initiated the study of soft topology and soft topological spaces independently. Shabir and Naz [23] defined soft topology on the collection  $\tau$  of soft sets over  $X$ . Consequently, they defined basic notions of soft topological spaces such as soft open and closed sets, soft subspace, soft closure, soft nbd of a point, soft  $T_i$  spaces,  $i = 1, 2, 3, 4$ , soft regular spaces, and soft normal spaces and established their several properties. In [25], we presented further several important properties of notions defined and studied in [23]. On the other hand, Cagman et al. [24] introduced a soft topology on a soft set and defined a soft topological space. They defined basic notions and concepts of soft topological spaces such as soft open and closed sets, soft interior, soft closure, soft basis, soft nbd of a point, soft limit point of a soft set, soft complement, soft difference, and soft boundary and established several properties of these soft notions. The work in both papers is appreciable.

The notion of soft topology by Cagman et al. [24] is more general than that by Shabir and Naz [23]. Therefore, we continue investigating the work of Cagman et al. [24] and follow their notations and mathematical formalism. In this paper, first, we define and study soft exterior. We characterize soft open sets, soft closed sets, and soft clopen sets in terms of soft boundary. We establish several interesting properties of soft interior, soft exterior, soft closure, and soft boundary and their relationship which are fundamental for further research on soft topology and will strengthen the foundations of the theory of soft topological spaces.

\*Correspondence: sabiriub@yahoo.com

<sup>3</sup>Department of Mathematics, Islamia University, Bahawalpur, 5000, Pakistan

<sup>4</sup>Present Address: Department of Mathematics, Yanbu University, P. O. Box 31387, Yanbu Alsinaiyah, Saudi Arabia

Full list of author information is available at the end of the article

**Preliminaries**

Now, we recall some definitions and results defined and discussed in [1,4,24]. Hereafter,  $U$  refers to an initial universe,  $E$  is a set of parameters,  $P(U)$  is the power set of  $U$ , and  $A$  is a nonempty subset of  $E$ .

**Definition 1.** A soft set  $F_A$  on the universe  $U$  is defined by the set of ordered pairs  $F_A = \{(x, f_A(x)) : x \in E, f_A(x) \in P(U)\}$ , where  $f_A : E \rightarrow P(U)$  such that  $f_A(x) = \phi$  if  $x \notin A$ . Here,  $f_A$  is called an approximate function of the soft set  $F_A$ . The value of  $f_A(x)$  may be arbitrary. Some of them may be empty, and some may have nonempty intersection. The class of all soft sets over  $U$  will be denoted by  $S(U)$ .

**Example 1.** Suppose that there are six houses in the universe  $U = \{h_1, h_2, h_3, h_4, h_5, h_6\}$  under consideration and that  $E = \{x_1, x_2, x_3, x_4, x_5\}$  is a set of decision parameters. The  $x_i$  ( $i = 1, 2, 3, 4, 5$ ) stand for the parameters ‘expensive,’ ‘beautiful,’ ‘wooden,’ ‘cheap,’ and ‘in green surroundings,’ respectively.

Consider the mapping  $f_A$  given by ‘houses (.),’ where (.) is to be filled in by one of the parameters  $x_i \in E$ . For instance,  $f_A(x_1)$  means ‘houses (expensive),’ and its functional value is the set  $\{h \in U : h \text{ is an expensive house}\}$ .

Suppose that  $A = \{x_1, x_3, x_4\} \subseteq E$  and  $f_A(x_1) = \{h_2, h_4\}$ ,  $f_A(x_3) = U$ , and  $f_A(x_4) = \{h_1, h_3, h_5\}$ . Then, we can view the soft set  $F_A$  as consisting of the following collection of approximations:

$$F_A = \{(x_1, \{h_2, h_4\}), (x_3, U), (x_4, \{h_1, h_3, h_5\})\}.$$

**Definition 2.** Let  $F_A \in S(U)$ . If  $f_A(x) = \phi$ , for all  $x \in E$ , then  $F_A$  is called an empty soft set, denoted by  $F_\phi$ .  $f_A(x) = \phi$  means that there is no element in  $U$  related to the parameter  $x \in E$ . Therefore, we do not display such elements in the soft sets as it is meaningless to consider such parameters.

**Definition 3.** Let  $F_A \in S(U)$ . If  $f_A(x) = U$ , for all  $x \in A$ , then  $F_A$  is called an  $A$ -universal soft set, denoted by  $F_{\bar{A}}$ . If  $A = E$ , then the  $A$ -universal soft set is called a universal soft set, denoted by  $F_{\bar{E}}$ .

**Definition 4.** Let  $F_A, F_B \in S(U)$ . Then,  $F_A$  is a soft subset of  $F_B$ , denoted by  $F_A \tilde{\subseteq} F_B$ , if  $f_A(x) \subseteq f_B(x)$ , for all  $x \in E$ .

**Definition 5.** Let  $F_A, F_B \in S(U)$ . Then,  $F_A$  and  $F_B$  are soft equal, denoted by  $F_A = F_B$ , if  $f_A(x) = f_B(x)$ , for all  $x \in E$ .

**Definition 6.** Let  $F_A, F_B \in S(U)$ . Then, the soft union  $F_A \tilde{\cup} F_B$ , the soft intersection  $F_A \tilde{\cap} F_B$ , and the soft difference  $F_A \tilde{\setminus} F_B$  of  $F_A$  and  $F_B$  are defined by the approximate

functions  $f_{A \tilde{\cup} B}(x) = f_A(x) \cup f_B(x)$ ,  $f_{A \tilde{\cap} B}(x) = f_A(x) \cap f_B(x)$ , and  $f_{A \tilde{\setminus} B}(x) = f_A(x) \setminus f_B(x)$ , respectively, and the soft complement  $F_A^{\tilde{c}}$  of  $F_A$  is defined by the approximate function  $f_A^{\tilde{c}}(x) = f_A^c(x)$ , where  $f_A^c(x)$  is the complement of the set  $f_A(x)$ , that is,  $f_A^c(x) = U \setminus f_A(x)$ , for all  $x \in E$ . It is easy to see that  $(F_A^{\tilde{c}})^{\tilde{c}} = F_A$  and  $F_{\tilde{\phi}} = F_{\bar{E}}$ .

**Proposition 1.** Let  $F_A \in S(U)$ . Then,

- (1)  $F_A \tilde{\cup} F_A = F_A, F_A \tilde{\cap} F_A = F_A.$
- (2)  $F_A \tilde{\cup} F_\phi = F_A, F_A \tilde{\cap} F_\phi = F_\phi.$
- (3)  $F_A \tilde{\cup} F_{\bar{E}} = F_{\bar{E}}, F_A \tilde{\cap} F_{\bar{E}} = F_A.$
- (4)  $F_A \tilde{\cup} F_A^{\tilde{c}} = F_{\bar{E}}, F_A \tilde{\cap} F_A^{\tilde{c}} = F_\phi.$

**Proposition 2.** Let  $F_A, F_B, F_C \in S(U)$ . Then,

- (1)  $F_A \tilde{\cup} F_B = F_B \tilde{\cup} F_A, F_A \tilde{\cap} F_B = F_B \tilde{\cap} F_A.$
- (2)  $(F_A \tilde{\cup} F_B)^{\tilde{c}} = F_B^{\tilde{c}} \tilde{\cap} F_A^{\tilde{c}}, (F_A \tilde{\cap} F_B)^{\tilde{c}} = F_B^{\tilde{c}} \tilde{\cup} F_A^{\tilde{c}}.$
- (3)  $F_A \tilde{\cup} (F_B \tilde{\cap} F_C) = (F_A \tilde{\cup} F_B) \tilde{\cap} (F_A \tilde{\cup} F_C).$
- (4)  $F_A \tilde{\cap} (F_B \tilde{\cup} F_C) = (F_A \tilde{\cap} F_B) \tilde{\cup} (F_A \tilde{\cap} F_C).$

**Definition 7.** Let  $F_A \in S(U)$ . The soft power set of  $F_A$  is defined by  $\tilde{P}(F_A) = \{F_{A_i} : F_{A_i} \tilde{\subseteq} F_A, i \in I \subseteq N\}$ , and its cardinality is defined by  $|\tilde{P}(F_A)| = 2^{\sum_{x \in E} |f_A(x)|}$ , where  $|f_A(x)|$  is the cardinality of  $f_A(x)$ .

**Example 2.** Let  $U = \{u_1, u_2, u_3\}$ ,  $E = \{x_1, x_2, x_3\}$ ,  $A = \{x_1, x_2\} \subseteq E$ , and

$F_A = \{(x_1, \{u_1, u_2\}), (x_2, \{u_2, u_3\})\}$ . Then,

- $F_{A_1} = \{(x_1, \{u_1\})\},$
- $F_{A_2} = \{(x_1, \{u_2\})\},$
- $F_{A_3} = \{(x_1, \{u_1, u_2\})\},$
- $F_{A_4} = \{(x_2, \{u_2\})\},$
- $F_{A_5} = \{(x_2, \{u_3\})\},$
- $F_{A_6} = \{(x_2, \{u_2, u_3\})\},$
- $F_{A_7} = \{(x_1, \{u_1\}), (x_2, \{u_2\})\},$
- $F_{A_8} = \{(x_1, \{u_1\}), (x_2, \{u_3\})\},$
- $F_{A_9} = \{(x_1, \{u_1\}), (x_2, \{u_2, u_3\})\},$
- $F_{A_{10}} = \{(x_1, \{u_2\}), (x_2, \{u_2\})\},$
- $F_{A_{11}} = \{(x_1, \{u_2\}), (x_2, \{u_3\})\},$
- $F_{A_{12}} = \{(x_1, \{u_2\}), (x_2, \{u_2, u_3\})\},$
- $F_{A_{13}} = \{(x_1, \{u_1, u_2\}), (x_2, \{u_2\})\},$
- $F_{A_{14}} = \{(x_1, \{u_1, u_2\}), (x_2, \{u_3\})\},$
- $F_{A_{15}} = F_A,$
- $F_{A_{16}} = F_\phi$

are all soft subsets of  $F_A$ . So,  $|\tilde{P}(F_A)| = 2^4 = 16$ .

**Definition 8.** Let  $F_A \in S(U)$ . A soft topology on  $F_A$ , denoted by  $\tilde{\tau}$ , is a collection of soft subsets of  $F_A$  having the following properties:

- (1)  $F_\Phi, F_A \in \tilde{\tau}$ .
- (2)  $\{F_{A_i} \subseteq F_A : i \in I \subseteq N\} \subseteq \tilde{\tau} \Rightarrow \bigcup_{i \in I} F_{A_i} \in \tilde{\tau}$ .
- (3)  $\{F_{A_i} \subseteq F_A : 1 \leq i \leq n, n \in N\} \subseteq \tilde{\tau} \Rightarrow \bigcap_{i=1}^n F_{A_i} \in \tilde{\tau}$ .

The pair  $(F_A, \tilde{\tau})$  is called a soft topological space.

**Example 3.** Let us consider the soft subsets of  $F_A$  that are given in Example 2. Then,  $\tilde{\tau}_1 = \{F_\Phi, F_A\}$ ,  $\tilde{\tau}_2 = P(\tilde{F}_A)$ , and  $\tilde{\tau}_3 = \{F_\Phi, F_A, F_{A_2}, F_{A_{11}}, F_{A_{13}}\}$  are soft topologies on  $F_A$ .

**Definition 9.** Let  $(F_A, \tilde{\tau})$  be a soft topological space. Then, every element of  $\tilde{\tau}$  is called a soft open set. Clearly,  $F_\Phi$  and  $F_A$  are soft open sets.

**Definition 10.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B \subseteq F_A$ . Then,  $F_B$  is said to be soft closed if the soft set  $F_B^c$  is soft open.

**Theorem 1.** Let  $(F_A, \tilde{\tau})$  be a soft topological space. Then, the following hold:

- (1) The universal soft set  $F_\Phi$  and  $F_A^c$  are soft closed sets.
- (2) Arbitrary soft intersections of the soft closed sets are soft closed sets.
- (3) Finite soft unions of the soft closed sets are soft closed sets.

**Definition 11.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B \subseteq F_A$ . Then, the soft interior of a soft set  $F_B$  is denoted by  $F_B^\circ$  and is defined as the soft union of all soft open subsets of  $F_B$ . Thus,  $F_B^\circ$  is the largest soft open set contained in  $F_B$ .

**Theorem 2.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B, F_C \subseteq F_A$ . Then,

- (1)  $F_\Phi^\circ = F_\Phi$ .
- (2)  $F_B^\circ \subseteq F_B$ .
- (3)  $(F_B^\circ)^\circ = F_B^\circ$ .
- (4)  $F_B$  is a soft open set if and only if  $F_B^\circ = F_B$ .
- (5)  $F_B \subseteq F_C$  implies  $F_B^\circ \subseteq F_C^\circ$ .
- (6)  $F_B^\circ \cap F_C^\circ = (F_B \cap F_C)^\circ$ .
- (7)  $F_B^\circ \cup F_C^\circ \subseteq (F_B \cup F_C)^\circ$ .

**Definition 12.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B \subseteq F_A$ . Then, the soft closure of  $F_B$ , denoted  $\overline{F_B}$ , is defined as the soft intersection of all soft closed supersets of  $F_B$ . Note that  $\overline{F_B}$  is the smallest soft closed set containing  $F_B$ .

**Theorem 3.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B \subseteq F_A$ .  $F_B$  is a closed soft set if and only if  $\overline{F_B} = F_B$ .

**Theorem 4.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B, F_C \subseteq F_A$ . Then,

- (1)  $\overline{\overline{F_B}} = \overline{F_B}$ .
- (2)  $F_C \subseteq F_B$  implies  $\overline{F_C} \subseteq \overline{F_B}$ .
- (3)  $\overline{F_B \cup F_C} = \overline{F_B} \cup \overline{F_C}$ .
- (4)  $\overline{F_B \cap F_C} \supseteq \overline{F_B} \cap \overline{F_C}$ .

**Definition 13.** Let  $(F_A, \tilde{\tau})$  be a soft topological and  $F_B \subseteq F_A$ . Then, the soft boundary of soft set  $F_B$  is denoted by  $F_B^b$  and is defined as  $F_B^b = \overline{F_B} \cap (F_B^c)$ .

**Definition 14.** Let  $(F_A, \tilde{\tau})$  be a soft topological space,  $F_B \subseteq F_A$ , and  $\alpha \in F_A$ . If every soft nbd of  $\alpha$  soft intersects  $F_B$  in some points other than  $\alpha$  itself, then  $\alpha$  is called a soft limit point of  $F_B$ . The set of all soft limit points of  $F_B$  is denoted by  $F_B'$ . In other words, if  $(F_A, \tilde{\tau})$  is a soft topological space,  $F_B, F_C \subseteq F_A$ , and  $\alpha \in F_A$ , then  $\alpha \in F_B'$  if and only if  $F_C \cap (F_B \setminus \{\alpha\}) \neq F_\Phi$ , for all  $F_C \in \tilde{\nu}(\alpha)$ .

### Soft topology

**Definition 15.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $\alpha \in F_A$ . If there is a soft open set  $F_B$  such that  $\alpha \in F_B$ , then  $F_B$  is called a soft open neighborhood (soft nbd) of  $\alpha$ . The set of all soft nbds of  $\alpha$ , denoted  $\tilde{\nu}(\alpha)$ , is called the family of soft nbds of  $\alpha$ , that is,  $\tilde{\nu}(\alpha) = \{F_B : F_B \in \tilde{\tau} : \alpha \in F_B\}$ .

**Example 4.** Let us consider the topological space  $(F_A, \tilde{\tau}_3)$  in Example 3 and  $\alpha = (x_1, \{u_1, u_2\}) \in F_A$ . Then,  $\tilde{\nu}(\alpha) = \{F_A, F_{A_{13}}\}$ .

The following theorem gives important properties of the soft nbd system:

**Proposition 3.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B, F_C \subseteq F_A$ . Then, the collection of soft nbd  $\tilde{\nu}(\alpha)$  at  $\alpha$  in  $(F_A, \tilde{\tau})$  has the following properties:

- (1) If  $F_B \in \tilde{\nu}(\alpha)$ , then  $\alpha \in F_B$ .
- (2) If  $F_B, F_C \in \tilde{\nu}(\alpha)$ , then  $F_B \cap F_C \in \tilde{\nu}(\alpha)$ .
- (3) If  $F_B \in \tilde{\nu}(\alpha)$  and  $F_B \subseteq F_C$ , then  $F_C \in \tilde{\nu}(\alpha)$ .
- (4) If  $F_B \in \tilde{\nu}(\alpha)$ , then there is an  $F_C \in \tilde{\nu}(\alpha)$  such that  $F_B \in \tilde{\nu}(\beta)$ , for each  $\beta \in F_C$ .
- (5)  $F_B \subseteq F_A$  is soft open if and only if  $F_B$  contains a soft nbd of each of its points.

*Proof.* (1) is obvious since  $F_B$  is a soft open nbd of  $\alpha \in F_A$ . So,  $F_B$  is a soft open set such that  $\alpha \in F_B$ .  
 (2) If  $F_B, F_C \in \tilde{\nu}(\alpha)$ , then there exist soft open sets  $F_D$  and  $F_G$  such that  $\alpha \in F_D \subseteq F_B$  and  $\alpha \in F_G \subseteq F_C$ . Therefore,  $\alpha \in F_D \cap F_G \subseteq F_B \cap F_C$ , and hence,  $F_B \cap F_C \in \tilde{\nu}(\alpha)$ .

- (3) Since  $F_B \in \tilde{v}(\alpha)$ , therefore there exists a soft open set  $F_D$  such that  $\alpha \in F_D \subseteq F_B$ . Then,  $\alpha \in F_D \subseteq F_B \subseteq F_C$  or  $\alpha \in F_D \subseteq F_C$ . Hence,  $F_C \in \tilde{v}(\alpha)$ .
- (4) Since  $F_B \in \tilde{v}(\alpha)$ , then  $\alpha \in F_C \subseteq F_B$ , for  $F_C$  soft open in  $F_A$ . Since  $\alpha \in F_C \subseteq F_C$ , then  $F_C \in \tilde{v}(\alpha)$ . If  $\beta \in F_C$ , then by (3)  $F_C \subseteq F_B$  implies that  $F_B \in \tilde{v}(\beta)$ , for each  $\beta \in F_C$ .
- (5) (i) Suppose  $F_B$  is a soft open in  $F_A$ , then  $\alpha \in F_B \subseteq F_B$  implies that  $F_B$  is a soft nbd of each  $\alpha \in F_B$ .
- (ii) If each  $\alpha \in F_B$  has a soft nbd  $F_{C_\alpha} \subseteq F_B$ , then  $F_B = \{\alpha : \alpha \in F_B\} \subseteq \bigcup_{\alpha \in F_B} F_{C_\alpha} \subseteq F_B$  or  $F_B = \bigcup_{\alpha \in F_B} F_{C_\alpha}$ . This implies that  $F_B$  is soft open in  $F_A$ . □

**Definition 16.** Let  $(F_A, \tilde{\tau})$  be a soft topological space. A soft nbd base at  $\alpha \in F_A$  is a subcollection  $\tilde{\delta}(\alpha)$  of soft nbd  $\tilde{v}(\alpha)$  having the property that each  $F_B \in \tilde{v}(\alpha)$  contains some  $F_C \in \tilde{\delta}(\alpha)$ , that is,  $\tilde{v}(\alpha)$  must be determined by  $\tilde{\delta}(\alpha)$  as follows:

$$\tilde{v}(\alpha) = \{F_B \subseteq F_A : F_C \subseteq F_B, \text{ for some } F_C \in \tilde{\delta}(\alpha)\}.$$

Each  $F_C \in \tilde{\delta}(\alpha)$  is called a basic soft open neighborhood of  $\alpha$ .

For the soft basic nbd system, we have the following properties:

**Proposition 4.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and for each  $\alpha \in F_A$ , let  $\tilde{\delta}(\alpha)$  be a soft nbd base at  $\alpha$ . Then,

- (1) If  $F_C \in \tilde{\delta}(\alpha)$ , then  $\alpha \in F_C$ .
- (2) If  $F_{C_1}, F_{C_2} \in \tilde{\delta}(\alpha)$ , then there is some  $F_{C_3} \in \tilde{\delta}(\alpha)$  such that  $F_{C_3} \subseteq F_{C_1} \tilde{\cap} F_{C_2}$ .
- (3) If  $F_C \in \tilde{\delta}(\alpha)$ , then there is some  $F_{C_0} \in \tilde{\delta}(\alpha)$  such that if  $\gamma \in F_{C_0}$ , then there is some  $F_D \in \tilde{\delta}(\gamma)$  with  $F_D \subseteq F_C$ .
- (4)  $F_B \subseteq F_A$  is soft open if and only if  $F_B$  contains a soft basic nbd of each of its points.

*Proof.* These properties are easily verified for soft basic nbds by referring to the corresponding properties of soft nbds in Proposition 3. □

Next, we prove the following theorem which relates the concepts of soft interior and soft closure:

**Theorem 5.** Let  $F_B$  be a soft set of soft topological space  $(F_A, \tilde{\tau})$ . Then,

- (1)  $\overline{(F_B^{\tilde{c}})} = (F_B^{\circ})^{\tilde{c}}$ .
- (2)  $(F_B^{\circ})^{\circ} = \overline{(F_B)^{\tilde{c}}}$ .
- (3)  $F_B^{\circ} = \overline{((F_B^{\tilde{c}}))^{\circ}}$ .

- (4)  $\overline{F_B} = (((F_B^{\tilde{c}})^{\circ})^{\tilde{c}})$ .
- (5)  $(F_B \tilde{\setminus} F_C)^{\circ} \subseteq F_B^{\circ} \tilde{\setminus} F_C^{\circ}$ .

*Proof.* (1) Let  $\alpha \in F_B$  such that  $\alpha \notin F_B^{\circ}$ . Then, for each soft open nbd  $F_C$  of  $\alpha$ ,  $F_C$  soft intersects  $F_B^{\tilde{c}}$ . Otherwise, for some soft open nbd  $F_C$  of  $\alpha$ ,  $F_C \tilde{\cap} F_B^{\tilde{c}} = F_{\Phi}$  or  $F_C \subseteq F_B$ . Since  $F_B^{\circ}$  is the largest soft open set in  $F_B$ , therefore  $\alpha \in F_C \subseteq F_B^{\circ}$ , which is a contradiction. Therefore, by Theorem 12(a) [24],  $\alpha \in \overline{(F_B^{\tilde{c}})}$ . Hence,  $(F_B^{\tilde{c}})^{\circ} \subseteq \overline{(F_B^{\tilde{c}})}$ .

Conversely, suppose  $\alpha \in \overline{(F_B^{\tilde{c}})}$ , then by Definition 17 [24],  $\alpha \in F_B^{\tilde{c}}$  or  $\alpha$  is a soft limit point of  $F_B^{\tilde{c}}$ . If  $\alpha \in F_B^{\tilde{c}}$ , then  $\alpha \in (F_B^{\tilde{c}})^{\circ}$ . In the second case,  $\alpha \notin F_B^{\circ}$ . Otherwise, by the definition of soft limit point [24],  $F_B^{\circ} \tilde{\cap} F_B^{\tilde{c}} \neq F_{\Phi}$ , which is false. Therefore,  $\alpha \in (F_B^{\tilde{c}})^{\circ}$ . This shows that  $F_B^{\tilde{c}} \subseteq (F_B^{\tilde{c}})^{\circ}$ . Combining, we get (1).

(2) is proved in [24].

(3) and (4) are directly obtained by taking the complements of (1) and (2), respectively.

$$\begin{aligned} (5) \quad (F_B \tilde{\setminus} F_C)^{\circ} &= (F_B \tilde{\cap} F_C^{\tilde{c}})^{\circ} \\ &= F_B^{\circ} \tilde{\cap} (F_C^{\tilde{c}})^{\circ} \quad (\text{by Theorem 8(3)[24]}) \\ &= F_B^{\circ} \tilde{\cap} (F_C)^{\tilde{c}} \quad (\text{by Theorem 5(2)}) \\ &\subseteq F_B^{\circ} \tilde{\cap} (F_C)^{\tilde{c}} \\ &= F_B^{\circ} \tilde{\setminus} F_C^{\circ}. \end{aligned}$$

□

Now, we define the following:

**Definition 17.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B \subseteq F_A$ . Then, the soft exterior of a soft set  $F_B$  is denoted by  $(F_B)^{\tilde{e}}$  and is defined as  $(F_B)^{\tilde{e}} = (F_B^{\tilde{c}})^{\circ}$ .

Thus,  $\alpha$  is called a soft exterior point of  $F_B$  if there exists a soft open set  $F_C$  such that  $\alpha \in F_C \subseteq F_B^{\tilde{c}}$ . We observe that  $(F_B)^{\tilde{e}}$  is the largest soft open set contained in  $F_B^{\tilde{c}}$ .

**Example 5.** In Example 2, we take  $F_B = \{(x_1, U), (x_2, \{u_1, u_2\})\}$ . Then, the soft exterior of  $F_B$  is  $F_{\Phi}$  since  $(F_B)^{\tilde{c}} = \{(x_2, \{u_3\})\}$ , and thus,  $(F_B)^{\tilde{e}} = ((F_B)^{\tilde{c}})^{\circ} = F_{\Phi}$ .

**Theorem 6.** Let  $F_B$  and  $F_C$  be soft sets of a soft topological space  $(F_A, \tilde{\tau})$ . Then,

- (1)  $(F_B)^{\tilde{e}} = (F_B^{\tilde{c}})^{\tilde{e}}$ .
- (2)  $(F_B \tilde{\cup} F_C)^{\tilde{e}} = (F_B)^{\tilde{e}} \tilde{\cap} (F_C)^{\tilde{e}}$ .
- (3)  $(F_B)^{\tilde{e}} \tilde{\cup} (F_C)^{\tilde{e}} \subseteq (F_B \tilde{\cap} F_C)^{\tilde{e}}$ .

*Proof.* (1) The proof follows from the definition.

$$\begin{aligned} (2) \quad (F_B \tilde{\cup} F_C)^{\tilde{e}} &= ((F_B \tilde{\cup} F_C)^{\tilde{c}})^{\circ} \\ &= (F_B^{\tilde{c}} \tilde{\cap} F_C^{\tilde{c}})^{\circ} \end{aligned}$$

$$= (F_B^c)^\circ \tilde{\cap} (F_C^c)^\circ \quad (\text{by Theorem 8(3)[24]})$$

$$= (F_B)^\tilde{e} \tilde{\cap} (F_C)^\tilde{e}.$$

$$(3) (F_B)^\tilde{e} \tilde{\cup} (F_C)^\tilde{e} = (F_B^c)^\circ \tilde{\cup} (F_C^c)^\circ$$

$$\subseteq (F_B^c \tilde{\cup} F_C^c)^\circ \quad (\text{by Theorem 8(4)[24]})$$

$$= ((F_B \tilde{\cap} F_C)^\tilde{e})^\circ$$

$$= (F_B \tilde{\cap} F_C)^\tilde{e}.$$

□

**Theorem 7.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B \tilde{\subseteq} F_A$ . Then, the following hold:

- (1)  $(F_B^b)^\tilde{c} = F_B^\circ \tilde{\cup} (F_B^c)^\circ = F_B^\circ \tilde{\cup} (F_B)^\tilde{e}.$
- (2)  $\overline{F_B} = F_B \tilde{\cup} F_B^b.$
- (3)  $F_B^\circ = (F_B) \tilde{\setminus} F_B^b.$

*Proof.* We use Theorem 5(1) and (2) to prove (1):

$$(1) F_B^\circ \tilde{\cup} (F_B^c)^\circ = ((F_B^\circ)^\tilde{c}) \tilde{\cup} (((F_B^c)^\circ)^\tilde{c})^\tilde{c}$$

$$= [(F_B^\circ)^\tilde{c} \tilde{\cap} (((F_B^c)^\circ)^\tilde{c})]^\tilde{c}$$

$$= [\overline{(F_B^c) \tilde{\cap} (F_B)^\tilde{e}}]^\tilde{c}$$

$$= (F_B^b)^\tilde{c}.$$

$$(2) F_B \tilde{\cup} F_B^b = F_B \tilde{\cup} (\overline{F_B \tilde{\cap} F_B^c})$$

$$= [F_B \tilde{\cup} \overline{F_B}] \tilde{\cap} [F_B \tilde{\cup} F_B^c]$$

$$= \overline{F_B} \tilde{\cap} [F_B \tilde{\cup} (F_B^c)]$$

$$= \overline{F_B} \tilde{\cap} F_A$$

$$= \overline{F_B}.$$

$$(3) F_B \tilde{\setminus} F_B^b = F_B \tilde{\cap} (F_B^b)^\tilde{c}$$

$$= F_B \tilde{\cap} (F_B^\circ \tilde{\cup} (F_B^c)^\circ) \quad (\text{by (1)})$$

$$= [F_B \tilde{\cap} F_B^\circ] \tilde{\cup} [F_B \tilde{\cap} (F_B^c)^\circ]$$

$$= F_B^\circ \tilde{\cup} F_\Phi$$

$$= F_B^\circ$$

□

**Remark 1.** (a) In [24], it is known that  $F_B^b = (F_B^c)^\tilde{b}.$   
 (b) From Theorem 7(1), it follows that  $F_A = F_B^\circ \tilde{\cup} (F_B)^\tilde{e} \tilde{\cup} F_B^b.$

**Theorem 8.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B \tilde{\subseteq} F_A$ . Then, the following hold:

- (1)  $F_B^b \tilde{\cap} F_B^\circ = F_\Phi.$
- (2)  $F_B^b \tilde{\cap} (F_B)^\tilde{e} = F_\Phi.$

*Proof.*

$$(1) F_B^b \tilde{\cap} F_B^\circ = F_B^\circ \tilde{\cap} (\overline{F_B \tilde{\cap} F_B^c})$$

$$= F_B^\circ \tilde{\cap} \overline{F_B \tilde{\cap} (F_B^c)^\circ} \quad (\text{by Theorem 5(1)})$$

$$= F_\Phi.$$

$$(2) F_B^b \tilde{\cap} (F_B)^\tilde{e} = (F_B^c)^\circ \tilde{\cap} (\overline{F_B \tilde{\cap} F_B^c})$$

$$= (F_B^c)^\circ \tilde{\cap} \overline{F_B \tilde{\cap} F_B^c}$$

$$= (\overline{F_B})^\tilde{c} \tilde{\cap} \overline{F_B \tilde{\cap} F_B^c} \quad (\text{by Theorem 5(1)})$$

$$= F_\Phi.$$

□

**Theorem 9.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B \tilde{\subseteq} F_A$ . Then,

- (1)  $F_B$  is soft open if and only if  $F_B \tilde{\cap} F_B^b = F_\Phi.$
- (2)  $F_B$  is soft closed if and only if  $F_B^b \tilde{\subseteq} F_B.$

*Proof.* (1) Let  $F_B$  be a soft open set. Then,  $F_B^\circ = F_B.$

Thus,  $F_B \tilde{\cap} F_B^b = F_B^\circ \tilde{\cap} F_B^b = F_\Phi$  (by Theorem 8(1)).

Conversely, let  $F_B \tilde{\cap} F_B^b = F_\Phi$ . Then,

$F_B \tilde{\cap} [\overline{F_B \tilde{\cap} F_B^c}] = F_\Phi$ ,  $F_B \tilde{\cap} F_B^c = F_\Phi$ , or  $\overline{F_B^c} \tilde{\subseteq} F_B^c$ , which implies that  $F_B^c$  is soft closed, and hence,  $F_B$  is soft open.

- (2) Let  $F_B$  be a soft closed set. Then,  $\overline{F_B} = F_B$ . Now,  $F_B^b = \overline{F_B \tilde{\cap} F_B^c} \tilde{\subseteq} \overline{F_B} = F_B$ , or  $F_B^b \tilde{\subseteq} F_B$  and conversely.

□

**Theorem 10.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B, F_C \tilde{\subseteq} F_A$ . Then, the following hold:

- (1)  $(F_B \tilde{\cup} F_C)^\tilde{b} \tilde{\subseteq} [F_B \tilde{\cap} F_C^c]^\tilde{b} \tilde{\cup} [F_C^b \tilde{\cap} \overline{F_B^c}].$
- (2)  $[F_B \tilde{\cap} F_C]^\tilde{b} \tilde{\subseteq} [F_B^b \tilde{\cap} \overline{F_C}] \tilde{\cup} [F_C^b \tilde{\cap} \overline{F_B}].$

*Proof.*

$$(1) (F_B \tilde{\cup} F_C)^\tilde{b} = \overline{(F_B \tilde{\cup} F_C)} \tilde{\cap} \overline{(F_B \tilde{\cup} F_C)^\tilde{c}}$$

$$= (\overline{F_B \tilde{\cup} F_C}) \tilde{\cap} (\overline{F_B^c \tilde{\cap} F_C^c}) \quad (\text{by Theorem 11(5)[24]})$$

$$\tilde{\subseteq} (\overline{F_B \tilde{\cup} F_C}) \tilde{\cap} (\overline{F_B^c} \tilde{\cap} \overline{F_C^c}) \quad (\text{by Theorem 11(4)[24]})$$

$$= [(F_B \tilde{\cap} \overline{F_B^c}) \tilde{\cap} \overline{F_C^c}] \tilde{\cup} [(F_C \tilde{\cap} \overline{F_B^c}) \tilde{\cap} \overline{F_C^c}]$$

$$= [(F_B \tilde{\cap} \overline{F_B^c}) \tilde{\cap} \overline{F_C^c}] \tilde{\cup} [(\overline{F_C}) \tilde{\cap} \overline{F_C^c}] \tilde{\cap} \overline{F_B^c}$$

$$= [F_B^b \tilde{\cap} \overline{F_B^c}] \tilde{\cup} [F_C^b \tilde{\cap} \overline{F_B^c}]$$

$$\tilde{\subseteq} F_B^b \tilde{\cup} F_C^b.$$

$$\begin{aligned}
 (2) [F_B \tilde{\cap} F_C]^{\tilde{b}} &= \overline{(F_B \tilde{\cap} F_C)} \tilde{\cap} \overline{(F_B \tilde{\cap} F_C)}^{\tilde{c}} \\
 &\subseteq [\overline{F_B \tilde{\cap} F_C}] \tilde{\cap} [\overline{F_B \tilde{\cap} F_C}]^{\tilde{c}} \quad (\text{by Theorem 11(4)[24]}) \\
 &= [\overline{F_B \tilde{\cap} F_C}] \tilde{\cap} [\overline{F_B} \tilde{\cap} \overline{F_C}]^{\tilde{c}} \quad (\text{by Theorem 11(5)[24]}) \\
 &= [(\overline{F_B} \tilde{\cap} \overline{F_C}) \tilde{\cap} \overline{F_B}] \tilde{\cap} [(\overline{F_B} \tilde{\cap} \overline{F_C}) \tilde{\cap} \overline{F_C}] \\
 &= (F_B^{\tilde{b}} \tilde{\cap} \overline{F_C}) \tilde{\cap} (\overline{F_B} \tilde{\cap} F_C^{\tilde{b}}).
 \end{aligned}$$

□

**Theorem 11.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B \subseteq F_A$ . Then, the following holds:  $((F_B^{\tilde{b}})^{\tilde{b}})^{\tilde{b}} = (F_B^{\tilde{b}})^{\tilde{b}}$ .

*Proof.*

$$\begin{aligned}
 ((F_B^{\tilde{b}})^{\tilde{b}})^{\tilde{b}} &= \overline{(F_B^{\tilde{b}})^{\tilde{b}}} \tilde{\cap} \overline{((F_B^{\tilde{b}})^{\tilde{b}})^{\tilde{b}}} \\
 &= (F_B^{\tilde{b}})^{\tilde{b}} \tilde{\cap} \overline{((F_B^{\tilde{b}})^{\tilde{b}})^{\tilde{c}}} \quad \dots \quad (1)
 \end{aligned}$$

Now, consider

$$\begin{aligned}
 ((F_B^{\tilde{b}})^{\tilde{b}})^{\tilde{c}} &= \overline{(F_B^{\tilde{b}})^{\tilde{b}}} \tilde{\cap} \overline{(F_B^{\tilde{b}})^{\tilde{b}}}^{\tilde{c}} \\
 &= [F_B^{\tilde{b}} \tilde{\cap} \overline{(F_B^{\tilde{b}})^{\tilde{b}}}^{\tilde{c}}] \\
 &= (F_B^{\tilde{b}})^{\tilde{c}} \tilde{\cap} \overline{(F_B^{\tilde{b}})^{\tilde{b}}}^{\tilde{c}}.
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 \overline{((F_B^{\tilde{b}})^{\tilde{b}})^{\tilde{c}}} &= \overline{[(F_B^{\tilde{b}})^{\tilde{c}} \tilde{\cap} \overline{(F_B^{\tilde{b}})^{\tilde{b}}}^{\tilde{c}}]} \\
 &= \overline{((F_B^{\tilde{b}})^{\tilde{c}}) \tilde{\cap} \overline{((F_B^{\tilde{b}})^{\tilde{b}})^{\tilde{c}}}} \quad (\text{by Theorem 11(5)[24]}) \\
 &= (F_C \tilde{\cap} \overline{(F_C)^{\tilde{c}}}) = F_A \quad \dots \quad (2)
 \end{aligned}$$

where  $F_C = \overline{((F_B^{\tilde{b}})^{\tilde{b}})^{\tilde{c}}}$ . From (1) and (2), we have  $((F_B^{\tilde{b}})^{\tilde{b}})^{\tilde{b}} = (F_B^{\tilde{b}})^{\tilde{b}} \tilde{\cap} F_A = (F_B^{\tilde{b}})^{\tilde{b}}$ . □

**Theorem 12.** Let  $(F_A, \tilde{\tau})$  be a soft topological space and  $F_B \subseteq F_A$ .  $F_B^{\tilde{b}} = F_\Phi$  if and only if  $F_B$  is a soft clopen set.

*Proof.* Suppose that  $F_B^{\tilde{b}} = F_\Phi$ .

(1) First, we prove that  $F_B$  is a soft closed set. Consider

$$\begin{aligned}
 F_B^{\tilde{b}} = F_\Phi &\Rightarrow \overline{F_B} \tilde{\cap} \overline{(F_B^{\tilde{c}})} = F_\Phi \\
 &\Rightarrow \overline{F_B} \subseteq \overline{(F_B^{\tilde{c}})}^{\tilde{c}} = F_B^{\tilde{c}} \tilde{\cap} F_B \quad (\text{by Theorem 5(3)}) \\
 &\Rightarrow \overline{F_B} \subseteq F_B \Rightarrow \overline{F_B} = F_B.
 \end{aligned}$$

This implies that  $F_B$  is a soft closed set.

(2) Using Theorem 5(3), we now prove that  $F_B$  is a soft open set:

$$\begin{aligned}
 F_B^{\tilde{b}} = F_\Phi &\Rightarrow \overline{F_B} \tilde{\cap} \overline{F_B}^{\tilde{c}} \text{ or } F_B \tilde{\cap} (F_B^{\tilde{c}})^{\tilde{c}} \\
 &= F_\Phi \Rightarrow F_B \subseteq F_B^{\tilde{c}} \tilde{\cap} F_B \Rightarrow F_B = F_B.
 \end{aligned}$$

This implies that  $F_B$  is a soft open set.

Conversely, suppose that  $F_B$  is a soft clopen set. Then,

$$\begin{aligned}
 F_B^{\tilde{b}} &= \overline{F_B} \tilde{\cap} \overline{(F_B^{\tilde{c}})} \\
 &= \overline{F_B} \tilde{\cap} (F_B^{\tilde{c}})^{\tilde{c}} \quad (\text{by Theorem 5(1)}) \\
 &= F_B \tilde{\cap} F_B^{\tilde{c}} = F_\Phi.
 \end{aligned}$$

□

## Conclusions

In the present work, we have defined and studied the important properties of soft exterior. We have established several results relating soft interior, soft exterior, soft closure, and soft boundary. Moreover, we have characterized soft open sets, soft closed sets, and soft clopen sets in terms of soft boundary. We have presented the base of the theory of soft topological spaces. These findings will strengthen the foundation of soft topological spaces and will help to establish a general framework for practical applications. We hope that researchers working on soft topological structures will be benefited.

## Competing interests

The authors declare that they have no competing interests.

## Author's contributions

Both authors contributed equally and significantly in writing this article. Both authors read and approved the final manuscript.

## Author details

<sup>1</sup>Center For Advanced Studies in Pure and Applied Mathematics, Bahauddin Zakariya University, Multan, 6000, Pakistan. <sup>2</sup>Present Address: Department of Mathematics, King Abdul Aziz University, P. O. Box 80203, Jeddah, 21589, Saudi Arabia. <sup>3</sup>Department of Mathematics, Islamia University, Bahawalpur, 5000, Pakistan. <sup>4</sup>Present Address: Department of Mathematics, Yanbu University, P. O. Box 31387, Yanbu Alsinaiyah, Saudi Arabia.

Received: 3 September 2012 Accepted: 7 October 2012

Published: 2 November 2012

## References

- Molodtsov, D: Soft set theory - first results. *Comput. Math. Appl.* **37**, 19–31 (1999)
- Molodtsov, D, Leonov, VY, Kovkov, DV: Soft sets technique and its application. *Nechetkie Sistemy i Myagkie Vychisleniya.* **1**(1), 8–39 (2006)
- Maji, PK, Biswas, R, Roy, R: An application of soft sets in a decision making problem. *Comput. Math. Appl.* **44**, 1077–1083 (2002)
- Maji, PK, Biswas, R, Roy, R: Soft set theory. *Comput. Math. Appl.* **45**, 555–562 (2003)
- Pie, D, Miao, D: From soft sets to information systems. *Granular Comput. 2005 IEEE Inter. Conf.* **2**, 617–621 ((2005))
- Chen, D: The parametrization reduction of soft sets and its applications. *Comput. Math. Appl.* **49**, 757–763 (2005)
- Acar, U, Koyuncu, F, Tanay, B: Soft sets and soft rings. *Comput. Math. Appl.* **59**, 3458–3463 (2010)
- Aktas, H, Cagman, N: Soft sets and soft groups. *Inf. Sci.* **1**(77), 2726–2735 (2007)
- Jun, YB: Soft BCK/BCI-algebras. *Comput. Math. Appl.* **56**, 1408–1413 (2008)
- Jun, YB, Park, CH: Applications of soft sets in ideal theory of BCK/BCI-algebras. *Inf. Sci.* **178**, 2466–2475 (2008)
- Jun, YB, Park, CH: Applications of soft sets in Hilbert algebras. *Iran. J. Fuzzy Syst.* **6**(2), 55–86 (2009)
- Jun, YB, Kim, HS, Negggersm, J: Pseudo d-algebras. *Inf. Sci.* **179**, 1751–1759 (2009)
- Jun, YB, Lee, KJ, Khan, A: Soft ordered semigroups. *Math. Logic Q.* **56**(1), 42–50 (2010)

14. Jun, YB, Lee, KJ, Park, CH: Soft set theory applied to commutative ideals in BCK-algebras. *J. Appl. Math. Inform.* **26**(3-4), 707–720 (2008)
15. Jun, YB, Lee, KJ, Park, CH: Soft set theory applied to ideals in d-algebras. *Comput. Math. Appl.* **57**, 367–378 (2009)
16. Jun, YB, Lee, KJ, Park, CH: Fuzzy soft set theory applied to BCK/BCI-algebras. *Comput. Math. Appl.* **59**, 3180–3192 (2010)
17. Jun, YB, Park, CH: Applications of soft sets in ideal theory of BCK/BCI-algebras. *Inf. Sci.* **178**, 2466–2475 (2008)
18. Park, CH, Jun, YB, Ozturk, MA: Soft WS-algebras. *Commun. Korean Math. Soc.* **23**(3), 313–324 (2008)
19. Sun, QM, Zhang, ZL, Liu, J: Soft sets and soft modules. In *Proceedings of Rough Sets and Knowledge Technology, Third International Conference, RSKT 2008 Chengdu, China, May 2008*. Edited by: Wang, G, Li, T, Grzymala-Busse, JW, Miao, D, Skowron, A, Yao, Y. Springer, Berlin, 403–409 (2008)
20. Feng, F, Jun, YB, Zhao, XZ: Soft semirings. *Comput. Math. Appl.* **56**, 2621–2628 (2008)
21. Ali, MI, Feng, F, Liu, XY, Min, WK, Shabir, M: On some new operations in soft set theory. *Comput. Math. Appl.* **57**, 1547–1553 (2009)
22. Shabir, M, Irfan, Ali, M: Soft ideals and generalized fuzzy ideals in semigroups. *New Math. Nat. Comput.* **5**, 599–615 (2009)
23. Shabir, M, Naz, M: On soft topological spaces. *Comput. Math. Appl.* **61**, 1786–1799 (2011)
24. Cagman, N, Karatas, S, Enginoglu, S: Soft topology. *Comput. Math. Appl.* **62**, 351–358 (2011)
25. Hussain, S, Ahmad, B: Some properties of soft topological spaces. *Comput. Math. Appl.* **62**, 4058–4067 (2011)

doi:10.1186/2251-7456-6-64

Cite this article as: Ahmad and Hussain: On some structures of soft topology. *Mathematical Sciences* 2012 **6**:64.

Archive of SID

Submit your manuscript to a SpringerOpen<sup>®</sup> journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► [springeropen.com](http://springeropen.com)