

ORIGINAL RESEARCH

Open Access

Unification of sixth-order iterative methods

Sanjay K Khattri^{1*} and Ioannis K Argyros²

Abstract

There exist many sixth-order iterative methods for solving nonlinear scalar equations. The purpose of this work is to bring them together. We develop a scheme for unifying sixth-order iterative methods. Convergence analysis shows that the methods, and family of methods, formulated through the scheme are sixth-order convergent. Finally, some computational results are reported to verify the developed theory.

Keywords: Iterative methods, Sixth order, Newton, Unification, Convergence, Numerical analysis, Infinite series, Nonlinear

MSC: 65H05; 41A25; 65B10

Introduction

Many problems in science and engineering require solving a nonlinear scalar equation $f(x) = 0$ [1-16]. As a result, solving nonlinear equations is an important part of scientific computing. There exist various iterative methods for solving nonlinear scalar equations. We are interested in sixth-order iterative methods, and their dynamics, to find a simple zero, that is $f(\gamma) = 0$ and $f'(\gamma) \neq 0$, of a nonlinear equation $f(x) = 0$. There exist many sixth-order iterative methods (see, e.g., [5-7,12,13,15]) for solving nonlinear scalar equations. The paper develops a scheme for constructing sixth-order iterative methods or family of methods. The scheme unifies existing sixth-order iterative methods. It is shown that various existing sixth-order iterative methods can be generated by the scheme through a proper choice of the independent parameters.

Let us first explore the existing sixth-order convergent iterative methods and family of methods. Lately, Sharma and Guha [13], through a modification of the well-known Ostrowski's method [11], developed the following sixth-order convergent method (SG):

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(x_n) - f(y_n)}{f(x_n) - 2f(y_n)} \frac{f'(y_n)}{f'(x_n)}, \\ x_{n+1} = z_n - \frac{f(x_n) + af(y_n) - f(z_n)}{f(x_n) + (a-2)f(y_n)} \frac{f'(z_n)}{f'(x_n)}, \end{cases} \quad (1)$$

where $a \in \mathbb{R}$. From here onwards, the preceding method is referred through the initials of the authors, i.e., SG. Neta [7] proposed a family, consisting of three steps and one parameter, of sixth-order convergent iterative methods (NETA):

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(x_n) + \alpha f(y_n) - f(z_n)}{f(x_n) + (\alpha - 2)f(y_n)} \frac{f'(y_n)}{f'(x_n)}, \\ x_{n+1} = z_n - \frac{f(x_n) - f(y_n) - f(z_n)}{f(x_n) - 3f(y_n)} \frac{f'(z_n)}{f'(x_n)}. \end{cases} \quad (2)$$

From here onwards, the above method is referred to as NETA. We observe that for $\alpha = 0$, the second step of the methods NETA and SG is the same, while for the choice $a = -1$, the third step of the methods NETA and SG is the same. In [6], Grau and Díaz-Barrero developed yet another sixth-order variant of Ostrowski's method:

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(x_n) - 2f(y_n)}{f(x_n) - 2f(y_n)} \frac{f'(y_n)}{f'(x_n)}, \\ x_{n+1} = z_n - \frac{f(x_n) - f(y_n)}{f(x_n) - 2f(y_n)} \frac{f'(z_n)}{f'(x_n)}. \end{cases} \quad (3)$$

The term $f(x_n)/(f(x_n) - 2f(y_n))$ is referred to as Ostrowski's correction factor [6]. From here onwards, the above method is referred to as GD. Through a suitable modification of Ostrowski's method, Chun and Ham

*Correspondence: sanjay.khattri@hsh.no

¹Department of Engineering, Stord/Haugesund University College, Stord, 5414, Norway

Full list of author information is available at the end of the article



[5] derived the following family of sixth-order iterative methods:

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(x_n)}{f(x_n) - 2f(y_n)} \frac{f(y_n)}{f'(x_n)}, \\ x_{n+1} = z_n - \mathcal{H}(u_n) \frac{f(z_n)}{f'(x_n)}, \end{cases} \quad (4)$$

where $u_n = f(y_n)/f(x_n)$ and $\mathcal{H}(t)$ represents a real valued function satisfying $\mathcal{H}(0) = 1$ and $\mathcal{H}'(0) = 2$. If one chooses $\mathcal{H}(t) = \frac{1+\beta t}{1+(\beta-2)t}$, one obtains the method SG, while if one chooses $\mathcal{H}(t) = \frac{1}{1-2t}$, one obtains the method GD. From here onwards, the above method is referred to as CH. It may be noticed that the methods SG, CH, and GD are formulated by modifying Ostrowski's method; as a result, the first two steps of these methods are the same. Through a graceful modification of the well-known Kung and Traub method [17], recently Chun and Neta [16] proposed the following sixth-order iterative method:

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(y_n)}{f'(x_n)} \frac{1}{\left[1 - \frac{f(y_n)}{f(x_n)}\right]^2}, \\ x_{n+1} = z_n - \frac{f(z_n)}{f'(x_n)} \frac{1}{\left[1 - \frac{f(y_n)}{f(x_n)} - \frac{f(z_n)}{f(x_n)}\right]^2}. \end{cases} \quad (5)$$

From here onwards, the above method is referred to as CN.

In this work, we propose a scheme for forming sixth-order convergent iterative methods. The methods formulated through the scheme consist of three steps. During each iteration, the developed methods require three functional evaluations and one evaluation of the derivative of the function. It is also shown that methods, such as SG, NETA, GD, and CH, that require three functional and one derivative evaluations can be formed through the proposed scheme. In consequence, the proposed scheme offers a unification of the sixth-order iterative methods, which is the main purpose of this work. The rest of the paper is organized as follows: The 'Scheme for constructing sixth-order iterative methods' section presents the scheme. In the 'Unification of sixth-order iterative methods' section, through the scheme, we generate various well-known sixth-order iterative methods. In the 'Numerical work' section, numerical and dynamical comparisons of various methods are shown. Finally, the 'Conclusions' section concludes the article.

Scheme for constructing sixth-order iterative methods

Our aspiration is to develop a sixth-order iterative scheme which unifies sixth-order methods published in the literature. For this purpose, we consider the following three-step iterative scheme:

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(y_n)}{f'(x_n)} \left(1 + \sum_{j=1}^m a_j \left(\frac{f(y_n)}{f(x_n)}\right)^j\right), \\ x_{n+1} = z_n - \frac{f(z_n)}{f'(x_n)} \left(1 + \sum_{k=1}^l b_k \left(\frac{\mu_1 f(y_n) + \mu_2 f(z_n)}{f(x_n)}\right)^k\right). \end{cases} \quad (6)$$

Here, a_j , b_k , μ_1 , μ_2 , m , and l are independent parameters. The parameters a_j , b_k , μ_1 , $\mu_2 \in \mathbb{R}$, while the parameters m and l are positive integers. The parameters are established through the following convergence theorem:

Theorem 1. *Let γ be a simple zero of a sufficiently differentiable function $f : \mathbf{D} \subset \mathbb{R} \mapsto \mathbb{R}$ in an open interval \mathbf{D} . If the initialization x_0 is sufficiently close to γ , then the scheme (6) defines sixth-order iterative methods iff $a_1 = 2$ and $b_1 = 2/\mu_1$, and the error equation for the family of methods is given as*

$$e_{n+1} = \frac{c_2 (c_3 c_1 - 5 c_2^2 + a_2 c_2^2) (-6 c_2^2 + c_2^2 b_2 \mu_1^2 + c_3 c_1)}{c_1^5} \times e_n^6 + O(e_n^7), \quad (7)$$

where $e_n = x_n - \gamma$ and $c_m = f^{(m)}(\gamma)/m!$ with $m \geq 1$.

Proof. The Taylor expansion of the function $f(x)$ around the solution γ is given as

$$f(x_n) = c_1 e_n + c_2 e_n^2 + c_3 e_n^3 + c_4 e_n^4 + O(e_n^5). \quad (8)$$

Furthermore, from the preceding equation, we have

$$f'(x_n) = c_1 + 2 c_2 e_n + 3 c_3 e_n^2 + 4 c_4 e_n^3 + O(e_n^4). \quad (9)$$

Dividing Equations 8 and 9,

$$\frac{f(x)}{f'(x)} = e_n - \frac{c_2}{c_1} e_n^2 - 2 \frac{c_3 c_1 - c_2^2}{c_1^2} e_n^3 - \frac{3 c_4 c_1^2 - 7 c_2 c_3 c_1 + 4 c_2^3}{c_1^3} e_n^4 + O(e_n^5). \quad (10)$$

From the first step of our scheme, we write

$$y_n - \gamma = e_n - \frac{f(x_n)}{f'(x_n)}; \quad (11)$$

substituting Equation 10 into the preceding equation yields

$$y_n - \gamma = \frac{c_2}{c_1} e_n^2 + 2 \frac{c_3 c_1 - c_2^2}{c_1^2} e_n^3 + \frac{3 c_4 c_1^2 - 7 c_2 c_3 c_1 + 4 c_2^3}{c_1^3} e_n^4 + O(e_n^5). \quad (12)$$

Expanding $f(y_n)$, around the solution γ , through the Taylor series and using $f(\gamma) = 0$,

$$f(y_n) = \sum_{j=1}^{\infty} c_j (y_n - \gamma)^j;$$

substituting Equation 12 into the above equation, we obtain

$$f(y_n) = c_2 e_n^2 + 2 \frac{c_3 c_1 - c_2^2}{c_1} e_n^3 + \frac{3 c_4 c_1^2 - 7 c_2 c_3 c_1 + 5 c_2^3}{c_1^2} e_n^4 + O(e_n^5). \quad (13)$$

Dividing Equations 8 and 13 gives

$$\frac{f(y_n)}{f(x_n)} = \frac{c_2}{c_1} e_n + \frac{2 c_3 c_1 - 3 c_2^2}{c_1^2} e_n^2 + \frac{3 c_4 c_1^2 - 10 c_2 c_3 c_1 + 8 c_2^3}{c_1^3} e_n^3 + O(e_n^4). \quad (14)$$

From the second step of our scheme (6), we may write

$$z_n = x_n - \frac{f(x_n)}{f'(x_n)} \left[1 + \frac{f(y_n)}{f(x_n)} \left(1 + a_1 \left(\frac{f(y_n)}{f(x_n)} \right) + a_2 \left(\frac{f(y_n)}{f(x_n)} \right)^2 + \dots \right) \right]; \quad (15)$$

substituting $f(x_n)/f'(x_n)$, from Equation 10, and $f(y_n)/f(x_n)$, from Equation 14, into the above equation yields

$$z_n - \gamma = \gamma - \frac{c_2^2 (a_1 - 2)}{c_1^2} e_n^3 - \frac{c_2 (4 a_1 c_3 c_1 - 7 a_1 c_2^2 + a_2 c_2^2 - 7 c_3 c_1 + 9 c_2^2)}{c_1^3} e_n^4 + O(e_n^5). \quad (16)$$

The Taylor expansion of $f(z_n)$ around the solution γ is given as

$$f(z_n) = \sum_{k=1}^{\infty} c_k (z_n - \gamma)^k; \quad (17)$$

substituting $z_n - \gamma$, from Equation 16, into the above equation returns

$$f(z_n) = - \frac{c_2^2 (a_1 - 2)}{c_1} e_n^3 - \frac{c_2 (4 a_1 c_3 c_1 - 7 a_1 c_2^2 + a_2 c_2^2 - 7 c_3 c_1 + 9 c_2^2)}{c_1^2} e_n^4 + O(e_n^5). \quad (18)$$

From the third step of our scheme (6), we have

$$x_{n+1} = z_n - \frac{f(z_n)}{f'(x_n)} \left[1 + b_1 \left(\mu_1 \frac{f(y_n)}{f(x_n)} + \mu_2 \frac{f(z_n)}{f(x_n)} \right) + b_2 \left(\mu_1 \frac{f(y_n)}{f(x_n)} + \mu_2 \frac{f(z_n)}{f(x_n)} \right)^2 + \dots \right];$$

substituting Equations 9, 14, and 18 into the above equation furnishes the following error relation:

$$x_{n+1} = \gamma + \frac{c_2^3 (-2 + b_1 \mu_1) a_1 - 2 e_n^4}{c_1^3} + \frac{c_2^2}{c_1^4} ((-11 a_1 - 11 b_1 \mu_1 + 6 b_1 a_1 \mu_1 + 20) c_1 c_3 + ((-2 + a_1) \mu_1^2 b_2 + ((-12 a_1 + a_2 + 19) \mu_1 + (-4 + 4 a_1 - a_1^2) \mu_2) b_1 - 2 a_2 + 18 a_1 - 26) c_2^2) e_n^5 + \frac{c_2}{c_1^5} ((-16 a_1 - 16 b_1 \mu_1 + 28 + 9 a_1 b_1 \mu_1) c_1^2 c_2 c_4 + (12 a_1 b_1 \mu_1 - 20 b_1 \mu_1 + 33 - 20 a_1) c_1^2 c_3^2 + (131 b_1 \mu_1 - 89 a_1 b_1 \mu_1 - 28 b_1 \mu_2 - 15 b_2 \mu_1^2 + 8 a_1 b_2 \mu_1^2 + 8 b_1 \mu_1 a_2 + 30 b_1 \mu_2 a_1 - 8 b_1 \mu_2 a_1^2 - 15 a_2 + 125 a_1 - 167) c_1 c_2^2 c_3 + (86 a_1 b_1 \mu_1 - 15 b_1 \mu_1 a_2 - 58 b_1 \mu_2 a_1 + 17 b_1 \mu_2 a_1^2 - 15 a_1 b_2 \mu_1^2 + 4 a_2 b_1 \mu_2 + a_2 b_2 \mu_1^2 - 8 b_2 \mu_1 \mu_2 + a_1 b_3 \mu_1^3 + b_1 \mu_1 a_3 - 2 a_2 b_1 \mu_2 a_1 + 8 b_2 \mu_1 \mu_2 a_1 - 2 b_2 \mu_1 \mu_2 a_1^2 - a_1^2 - 98 a_1 + 24 a_2 - 2 a_3 + 108 + 48 b_1 \mu_2 - 111 b_1 \mu_1 + 25 b_2 \mu_1^2 - 2 b_3 \mu_1^3) c_2^4) e_n^6 + O(e_n^7). \quad (19)$$

From the above error relation, we may deduce that the three-step scheme (6) will define sixth-order methods, or family of methods, if the following three equations are satisfied simultaneously:

$$\begin{cases} (a_1 - 2)(b_1 \mu_1 - 2) & = 0, \\ -11 a_1 - 11 b_1 \mu_1 + 6 b_1 a_1 \mu_1 + 20 & = 0, \\ (-2 + a_1) \mu_1^2 b_2 + ((-12 a_1 + a_2 + 19) \mu_1 + (-4 + 4 a_1 - a_1^2) \mu_2) b_1 - 2 a_2 + 18 a_1 - 26 & = 0. \end{cases}$$

From a simple calculation, we see that a solution is

$$a_1 = 2 \quad \text{and} \quad b_1 = \frac{2}{\mu_1}.$$

Substituting $a_1 = 2$ and $b_1 = 2/\mu_1$ in Equation 19 produces the required error equation (7). This proves our theorem. \square

Consequently, this work contributes the following three-step sixth-order convergent iterative scheme for solving nonlinear equations:

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(y_n)}{f'(x_n)} \left(1 + 2 \left(\frac{f(y_n)}{f(x_n)} \right) + \sum_{j=2}^m a_j \left(\frac{f(y_n)}{f(x_n)} \right)^j \right), \\ x_{n+1} = z_n - \frac{f(z_n)}{f'(x_n)} \left(1 + \frac{2}{\mu_1} \left(\frac{\mu_1 f(y_n) + \mu_2 f(z_n)}{f(x_n)} \right) + \sum_{k=2}^l b_k \left(\frac{\mu_1 f(y_n) + \mu_2 f(z_n)}{f(x_n)} \right)^k \right). \end{cases} \quad (20)$$

In the preceding scheme, the parameters a_j, b_k (with $j \geq 2$ and $k \geq 2$), and μ_m (with $m = 1, 2$) are free to choose. From here onwards, the above scheme is referred to as *USS* for short. Accordingly, *USS* presents opportunities to form various sixth-order methods. The next section explores few interesting choices of these parameters to formulate methods and family of methods from the published literature.

Unification of sixth-order iterative methods

Let us now derive methods from the published literature. Let us first construct the family of methods developed by Sharma and Guha [13]. For this purpose, we consider

$$\begin{aligned} a_j &= 2^j & \text{for } j \geq 2, \quad m = \infty, \quad \mu_1 = 1, \\ b_k &= 2(2 - \beta)^{k-1} & \text{for } k \geq 2, \quad l = \infty, \quad \mu_2 = 0. \end{aligned}$$

Here, $\beta \in \mathbb{R}$. Substituting these values in the second and third steps of the proposed scheme *USS*, we get

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(y_n)}{f'(x_n)} \left(1 + 2 \left(\frac{f(y_n)}{f(x_n)} \right) + \sum_{j=2}^{\infty} 2^j \left(\frac{f(y_n)}{f(x_n)} \right)^j \right), \\ x_{n+1} = z_n - \frac{f(z_n)}{f'(x_n)} \left(1 + 2 \left(\frac{f(y_n)}{f(x_n)} \right) + \sum_{k=2}^{\infty} 2(2 - \beta)^{k-1} \left(\frac{f(y_n)}{f(x_n)} \right)^k \right). \end{cases}$$

Using $1 + r + r^2 + r^3 + \dots = 1/(1 - r)$ for $|r| < 1$ in the second and third steps of the preceding equation, we obtain the method, *SG*, developed by Sharma and Guha

[13]. Now, to formulate the family of methods developed by Chun and Ham [5], we consider

$$\begin{aligned} a_j &= 2^j & \text{for } j \geq 2, \quad m = \infty, \quad \mu_1 = 1, \\ b_k &= \omega_k & \text{for } k \geq 2, \quad l = \infty, \quad \mu_2 = 0. \end{aligned}$$

Here, $a \in \mathbb{R}$. Substituting the preceding choices in the second and third steps of the scheme *USS*, we get

$$\begin{cases} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(y_n)}{f'(x_n)} \left(1 + 2 \left(\frac{f(y_n)}{f(x_n)} \right) + \sum_{j=2}^{\infty} 2^j \left(\frac{f(y_n)}{f(x_n)} \right)^j \right), \\ x_{n+1} = z_n - \frac{f(z_n)}{f'(x_n)} \left(1 + 2 \left(\frac{f(y_n)}{f(x_n)} \right) + \sum_{k=2}^{\infty} \omega_k \left(\frac{f(y_n)}{f(x_n)} \right)^k \right). \end{cases} \quad (21)$$

Using $1 + r + r^2 + r^3 + \dots = 1/(1 - r)$ for $|r| < 1$ in the second step of the preceding equation, we obtain the second step of the method, *CH*, developed by Chun and Ham [5]. The third step of the method *CH* is given as $x_{n+1} = z_n - f(z_n)/f'(x_n) \mathcal{H}(u_n)$ (see Equation 4). Here, $u_n = f(y_n)/f(x_n)$ and $\mathcal{H}(t)$ is a real valued function satisfying $\mathcal{H}(0) = 1$ and $\mathcal{H}'(0) = 2$. Through a simple calculation, the function $\mathcal{H}(u_n)$ may be expressed as

$$\mathcal{H} \left(\frac{f(y_n)}{f(x_n)} \right) = 1 + 2 \left(\frac{f(y_n)}{f(x_n)} \right) + \sum_{k=2}^{\infty} \omega_k \left(\frac{f(y_n)}{f(x_n)} \right)^k.$$

As a consequence, the formulated method (21), through the scheme *USS*, is the method developed by Chun and Ham [5]. The method of Grau and Diaz-Barrero [6] can be derived by considering the following:

$$\begin{aligned} a_j &= 2^j & \text{for } j \geq 2, \quad m = \infty, \quad \mu_1 = 1, \\ b_k &= 2^k & \text{for } k \geq 2, \quad l = \infty, \quad \mu_2 = 0, \end{aligned}$$

in the second and third steps of the developed scheme *USS*. To derive the family of methods developed by Neta et al. (2), the choices are

$$\begin{aligned} a_j &= 2(-a - 2)^{j-1} & \text{for } j \geq 2, \quad m = \infty, \quad \mu_1 = 1, \\ b_k &= 2(-(-3))^{k-1} & \text{for } k \geq 2, \quad l = \infty, \quad \mu_2 = 0. \end{aligned}$$

Here, $a \in \mathbb{R}$. To formulate the recently developed sixth-order method *CN* by Chun and Neta [16], the parameters, in the scheme *USS*, are chosen as follows:

$$\begin{aligned} a_j &= (j + 1) & \text{for } j \geq 2, \quad m = \infty, \quad \mu_1 = 1, \\ b_k &= (k + 1) & \text{for } k \geq 2, \quad l = \infty, \quad \mu_2 = 1. \end{aligned}$$

Table 1 The (number of function evaluations, COC) for various sixth-order iterative methods

$f(x)$	x_0	SG	NETA	GD	CH	CN
$f_1(x)$	100	(36,6)	(36,6)	(28,6)	(36,6)	(28,6)
$f_2(x)$	100	(172,6)	(884,6)	(160,6)	(172,6)	(108,6)
$f_3(x)$	-1	(20,6)	(20,6)	(20,6)	(20,6)	(32,6)
$f_4(x)$	1	(20,6)	(20,6)	(20,6)	(20,6)	(20,6)
$f_5(x)$	1	(24,6)	(24,6)	(24,6)	(24,6)	(24,6)
$f_5(x)$	3	(24,6)	(28,6)	(20,6)	(24,6)	(20,6)
$f_6(x)$	-1.5	(24,6)	(28,6)	(20,6)	(24,6)	(20,6)

Substituting the above values in the scheme USS, we get

$$\left\{ \begin{array}{l} y_n = x_n - \frac{f(x_n)}{f'(x_n)}, \\ z_n = y_n - \frac{f(y_n)}{f'(x_n)} \left(1 + 2 \left(\frac{f(y_n)}{f(x_n)} \right) + \sum_{j=2}^{\infty} (j+1) \left(\frac{f(y_n)}{f(x_n)} \right)^j \right), \\ x_{n+1} = z_n - \frac{f(z_n)}{f'(x_n)} \left(1 + 2 \left(\frac{f(y_n) + f(z_n)}{f(x_n)} \right) + \sum_{k=2}^{\infty} (k+1) \left(\frac{f(y_n) + f(z_n)}{f(x_n)} \right)^k \right). \end{array} \right.$$





Figure 1 The attraction basins for the polynomial $f(z) = z^3 - 1$. (a) SG, $\bar{\mu} = 4.46$ and $\omega = 10.37$, $a = 2$; (b) CN, $\bar{\mu} = 3.98$ and $\omega = 1.98$; (c) GD, $\bar{\mu} = 3.94$ and $\omega = 0.56$; (d) NETA, $\bar{\mu} = 4.25$ and $\omega = 10.95$, $\alpha = 5$.



Using $1 + 2r + 3r^2 + 4r^3 + \dots = 1/(1-r)^2$ for $|r| < 1$ in the second and third steps of the preceding method yields the method of Chun and Neta [16].

Numerical work

Let $\{x_n\}_{n=0}^{\infty}$ be a sequence, generated by an iterative method, converging to γ and $e_n = x_n - \gamma$. If there exist a real number $\xi \in [1, \infty)$ and a nonzero constant C such that

$$\lim_{n \rightarrow \infty} |e_{n+1}|/|e_n|^\xi = C,$$

then ξ is called the convergence order of the sequence and the constant C is called the asymptotic error constant.

From the preceding relation, the computational order of convergence (COC) is approximated as follows:

$$\rho \approx \frac{\log |(x_{n+1} - \gamma)/(x_n - \gamma)|}{\log |(x_n - \gamma)/(x_{n-1} - \gamma)|}.$$

All the computations are performed in the programming language C++. For numerical precision, the C++ library ARPREC [1] is being used. For the convergence of the method, it is required that the distance of two consecutive iterates ($|x_{n+1} - x_n|$) and the absolute value of the function ($|f(x_n)|$), also referred to as residual, be less than 10^{-300} . The maximum allowed iteration is 200.

Solving nonlinear equations

The methods are tested for the following functions:

$$\begin{aligned}
 f_1(x) &= x^3 + 4x^2 - 10, & \gamma &\approx 1.365. \\
 f_2(x) &= x^2 - \exp(x) - 3x + 2, & \gamma &\approx 0.257. \\
 f_3(x) &= x \exp(x^2) - \sin^2(x) + 3 \cos(x) + 5, & \gamma &\approx -1.207. \\
 f_4(x) &= x^4 + \sin \frac{\pi}{x^2} - 5, & \gamma &= \sqrt{2}. \\
 f_5(x) &= e^x \sin x + \log(1 + x^2), & \gamma &= 0. \\
 f_6(x) &= \sqrt{2 + x^2} \sin \frac{\pi}{x^2} + \frac{1}{x^4 + 1} - \frac{17\sqrt{3} + 1}{17}, & \gamma &= -2.
 \end{aligned}$$

Various free parameters are chosen as in the method SG: $a = 2$, in the method NETA: $\alpha = 5$, and in the method CH: $\beta = 3$. The outcome of the numerical experimentation is presented in Table 1. Table 1 reports the number of function evaluations and COC during the second-to-the-last iterative step. COC is rounded to the nearest significant digits. In Table 1, we can see that the methods GD and CN show better results.

Dynamic behavior of various sixth-order methods

Let $f(x)$ be a complex polynomial and γ be one of its zeros. Furthermore, let $x_0 \in \mathbb{C}$ be a starting point for an iterative method. Then, the sequence $\{x_n\}_{n=0}^{\infty}$, generated by the iterative methods, may converge or may not converge to the zero γ . If the sequence converges to the zero γ , then the starting point x_0 is attracted to γ . The basin of attraction, corresponding to a zero γ of the complex polynomial $f(x)$, is the set of all the starting points x_0 which are attracted to γ .

To make the figures, first we take a rectangle D , of size $[-1.5, 1.5] \times [-1.5, 1.5]$, and then divide the rectangle into $1,000 \times 1,000$ grids [18–20]. Further, we apply the iterative method starting at each grid point. The iterative methods converge if the residual, in a maximum of 10 iterations, is less than 10^{-15} . Given that the iterative method does not generate a residual less than 10^{-10} in the maximum allowed iterations, we say that the initial point does not converge to any root. Let us denote the number of mean iterations by $\bar{\mu}$ and the percentage of diverging points by ω [14].

We consider the polynomial $f(x) = x^n - 1$, $x \in \mathbb{C}$ for finding the n th roots of unity. The n th roots of unity are given by [18–20]

$$\alpha_j = \cos\left(\frac{2\pi(j-1)}{n}\right) + i \sin\left(\frac{2\pi(j-1)}{n}\right), \quad j = 1, 2, \dots, n.$$

The outcome of our numerical experimentation is reported in Figures 1 and 2. From these figures, we notice that the methods GD and CN which require the least number of iterations $\bar{\mu}$ while diverging at the least number of points show better results.

Conclusions

This work has developed a scheme to unify various sixth-order iterative methods. Comparison among iterative methods, by using the basins of attraction and also through numerical computations, is also presented. Ideas presented in this work can be further developed and extended to include iterative methods of higher orders such as seven or eight.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

This work was done through the cooperation of both authors. Both authors have contributed in it equally. Both authors read and approved the final manuscript.

Acknowledgements

We gratefully acknowledge the help from Professor Hadi Susanto during the numerical experimentation. We appreciate the reviewers for their insightful comments which improved the quality of our paper.

Author details

¹Department of Engineering, Stord/Haugesund University College, Stord, 5414, Norway. ²Department of Mathematical Sciences, Cameron University, Lawton, OK, 73505, USA.

Received: 21 June 2012 Accepted: 22 November 2012

Published: 22 January 2013

References

- Bailey, DH, Hida, Y, Li, XS, Thompson, B: ARPREC (C++/Fortran-90 arbitrary precision package). [http://crd.lbl.gov/~dhbailey/mpdist/] (2012). Accessed Jan 2012
- Argyros, IK: Computational Theory of Iterative Methods. In *Studies in Computational Mathematics*. Edited by: Chui, CK, Wuytack, L. Elsevier, New York (2007)
- Argyros, IK, Hilout, S: A unifying theorem for Newton's method on spaces with a convergence structure. *J Complexity*. **27**(1), 39–54 (2011)
- Argyros, IK, Chen, D, Qian, Q: The Jarrat method in Banach space setting. *J. Comput. Appl. Math.* **51**, 103–106 (1994)
- Chun, C, Ham, Y: Some sixth-order variants of Ostrowski root-finding methods. *Appl. Math. Comput.* **193**, 389–394 (2007)
- Grau, M, Díaz-Barrero, JL: An improvement to Ostrowski root-finding method. *Appl. Math. Comput.* **173**(1), 450–456 (2006)
- Neta, B: A sixth-order family of methods for nonlinear equations. *Int. J. Comput. Math.* **7**, 157–161 (1979)
- Khattri, SK, Argyros, IK: Sixth order derivative free family of iterative methods. *Appl. Math. Comput.* **217**(12), 5500–5507 (2011)
- Khattri, SK, Argyros, IK: How to develop fourth and seventh order iterative methods? *Novi. Sad. J. Math.* **40**, 61–67 (2010)
- Khattri, SK, Log, T: Constructing third-order derivative-free iterative methods. *Int. J. Comput. Math.* **88**(7), 1509–1518 (2011)
- Ostrowski, AM: *Solutions of Equations and System Equations*. Academic, New York (1960)
- Ren, H, Wu, Q, Bi, W: New variants of Jarratt's method with sixth-order convergence. *Numer. Algorithm.* **52**, 585–603 (2009)
- Sharma, JR, Guha, RK: A family of modified Ostrowski methods with accelerated sixth order convergence. *Appl. Math. Comput.* **190**, 111–115 (2007)
- Traub, JF: *Iterative Methods for the Solution of Equations*. Prentice Hall, Englewood Cliffs (1964)
- Wang, X, Kou, J, Li, Y: A variant of Jarratt method with sixth-order convergence. *Appl. Math. Comput.* **204**, 14–19 (2008)
- Changbum, C, Neta, B: A new sixth-order scheme for nonlinear equations. *Appl. Math. Lett.* **25**(2), 185–189 (2012)
- Kung, HT, Traub, JF: Optimal order of one-point and multipoint iterations. *J. Assoc. Comput. Mach.* **21**, 643–651 (1974)

18. Ardelean, G: A comparison between iterative methods by using the basins of attraction. *Appl. Math. Comput.* **218**(1), 88–95 (2011)
19. Amat, S, Busquier, S, Plaza, S: Dynamics of the King and Jarratt iterations. *Aequationes Math.* **69**, 212–223 (2005)
20. Susanto, H, Karjanto, N: Newtons method basin of attraction revisited. *Appl. Math. Comput.* **215**, 1084–1090 (2009)

doi:10.1186/2251-7456-7-5

Cite this article as: Khattri and Argyros: Unification of sixth-order iterative methods. *Mathematical Sciences* 2013 **7**:5.

Archive of SID

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Immediate publication on acceptance
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ springeropen.com
