

ORIGINAL RESEARCH

Open Access

A new generalization of some integral inequalities for (α, m) -convex functions

İmdat İşcan

Abstract

In this paper, we derive new estimates for the remainder term of the midpoint, trapezoid, and Simpson formulae for functions whose derivatives in absolute value at certain power are (α, m) -convex.

Keywords: (α, m) -Convex function, Hermite-Hadamard's inequality, Simpson type inequality, Trapezoid inequality, Midpoint inequality

Introduction

Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a convex function defined on the interval I of real numbers and $a, b \in I$ with $a < b$. The following double inequality is well known in the literature as Hermite-Hadamard integral inequality

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{2}. \quad (1)$$

The class of (α, m) -convex functions was first introduced in [1], and it is defined as follows:

The function $f : [0, b] \rightarrow \mathbb{R}$, $b > 0$, is said to be (α, m) -convex where $(\alpha, m) \in [0, 1]^2$, if we have

$$f(tx + m(1-t)y) \leq t^\alpha f(x) + m(1-t^\alpha)f(y)$$

for all $x, y \in [0, b]$ and $t \in [0, 1]$. It can be easily deduced that for $(\alpha, m) \in \{(0, 0), (\alpha, 0), (1, 0), (1, m), (1, 1), (\alpha, 1)\}$ one obtains the following classes of functions: increasing, α star-shaped, star-shaped, m -convex, convex, and α -convex functions, respectively.

As denoted by $K_m^\alpha(b)$, the set of all (α, m) -convex functions is on $[0, b]$ for which $f(0) \leq 0$. For recent results and generalizations concerning (α, m) -convex functions, see [1-8].

In [7], Set et al. proved the following Hadamard type inequality for (α, m) -convex functions.

Theorem 1. Let $f : [0, \infty) \rightarrow \mathbb{R}$ be an (α, m) -convex function with $(\alpha, m) \in (0, 1]^2$. If $0 \leq a < b < \infty$ and $f \in L[a, b]$, then one has the inequality

$$\frac{1}{b-a} \int_a^b f(x) dx \leq \min \left\{ \frac{f(a) + \alpha m f\left(\frac{b}{m}\right)}{\alpha + 1}, \frac{f(b) + \alpha m f\left(\frac{a}{m}\right)}{\alpha + 1} \right\}. \quad (2)$$

The following inequality is well known in the literature as Simpson's inequality.

Let $f : [a, b] \rightarrow \mathbb{R}$ be a four-time continuously differentiable mapping on (a, b) and $\|f^{(4)}\|_\infty = \sup_{x \in (a, b)} |f^{(4)}(x)| < \infty$. Then the following inequality holds:

$$\left| \frac{1}{3} \left[\frac{f(a)+f(b)}{2} + 2f\left(\frac{a+b}{2}\right) \right] - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \frac{1}{2880} \|f^{(4)}\|_\infty (b-a)^4.$$

In recent years, many authors have studied error estimations for Simpson's inequality; for refinements, counterparts, generalizations, and new Simpson type inequalities, see [9-12].

In this paper, in order to provide a unified approach to establish midpoint, trapezoid, and Simpson type inequality for functions whose derivatives in absolute value at certain power are (α, m) -convex, we derive a general integral identity for convex functions.

Correspondence: imdat.iscan@giresun.edu.tr
Department of Mathematics, Faculty of Arts and Sciences, Giresun University
28100, Giresun, Turkey

Main results

In order to generalize the classical Trapezoid, midpoint and Simpson type inequalities and prove them, we need the following Lemma:

Lemma 1. *Let $f : I \subseteq \mathbb{R} \rightarrow \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a, b]$, where $a, b \in I$ with $a < b$, $\lambda, \mu \in [0, 1]$ and $m \in (0, 1]$. Then the following equality holds:*

$$\begin{aligned} & \lambda (\mu f(a) + (1 - \mu)f(mb)) + (1 - \lambda)f(\mu a + m(1 - \mu)b) \\ & - \frac{1}{mb - a} \int_a^{mb} f(x)dx = (mb - a) \\ & \times \left[\int_0^\mu [-t + \lambda(1 - \mu)]f'(ta + m(1 - t)b) dt \right. \\ & \left. + \int_\mu^1 [-t + (1 - \mu)\lambda]f'(ta + m(1 - t)b) dt \right]. \end{aligned} \tag{3}$$

A simple proof of the equality can be done by performing an integration by parts in the integrals from the right side and changing the variable. The details are left to the interested reader.

Theorem 2. *Let $f : I \subseteq [0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a, b]$, where $a, b \in I^\circ$ with $a < b$ and $\lambda, \mu \in [0, 1]$. If $|f'|^q$ is (α, m) -convex on $[a, b]$, for $(\alpha, m) \in (0, 1]^2$, $mb > a$, $q \geq 1$, then the following inequality holds:*

where

$$\varepsilon_1 = -\frac{\mu^2}{2} + \lambda(1 - \mu)\mu, \quad \varepsilon_2 = [\lambda(1 - \mu)]^2 - \varepsilon_1,$$

$$\varepsilon_3 = (1 - \lambda\mu)^2 - (1 - \lambda\mu)(1 + \mu) + \frac{1 + \mu^2}{2},$$

$$\varepsilon_4 = \frac{1 - \mu^2}{2} - (1 - \lambda\mu)(1 - \mu),$$

$$\delta_1 = \frac{\lambda(1 - \mu)\mu^{\alpha+1}}{\alpha + 1} - \frac{\mu^{\alpha+2}}{\alpha + 2},$$

$$\delta_2 = \lambda(1 - \mu)\mu - \frac{\mu^2}{2} - \delta_1,$$

$$\delta_3 = \frac{2[\lambda(1 - \mu)]^{\alpha+2}}{(\alpha + 1)(\alpha + 2)} - \frac{\lambda(1 - \mu)\mu^{\alpha+1}}{\alpha + 1} + \frac{\mu^{\alpha+2}}{\alpha + 2},$$

$$\delta_4 = [\lambda(1 - \mu)]^2 - \lambda\mu(1 - \mu) + \frac{\mu^2}{2} - \delta_3,$$

$$\beta_1 = \frac{2(1 - \lambda\mu)^{\alpha+2}}{(\alpha + 1)(\alpha + 2)} - \frac{(1 - \lambda\mu)(1 + \mu^{\alpha+1})}{\alpha + 1} + \frac{1 + \mu^{\alpha+2}}{\alpha + 2},$$

$$\beta_2 = (1 - \lambda\mu)^2 - (1 - \lambda\mu)(1 + \mu) + \frac{1 + \mu^2}{2} - \beta_1,$$

$$\beta_3 = \frac{1 - \mu^{\alpha+2}}{\alpha + 2} - (1 - \lambda\mu)\frac{1 - \mu^{\alpha+1}}{\alpha + 1},$$

$$\beta_4 = (1 - \lambda\mu)(\mu - 1) + \frac{1 - \mu^2}{2} - \beta_3.$$

Proof. From Lemma 1 and using the properties of modulus and the well known power mean inequality, we have

$$\begin{aligned} & \left| \lambda (\mu f(a) + (1 - \mu)f(mb)) + (1 - \lambda)f(\mu a + m(1 - \mu)b) - \frac{1}{mb - a} \int_a^{mb} f(x)dx \right| \tag{4} \\ & \leq \begin{cases} (mb - a) \left\{ \varepsilon_2^{1-\frac{1}{q}} (\delta_3 |f'(a)|^q + m\delta_4 |f'(b)|^q)^{\frac{1}{q}} \right. \\ \quad \left. + \varepsilon_3^{1-\frac{1}{q}} (\beta_1 |f'(a)|^q + m\beta_2 |f'(b)|^q)^{\frac{1}{q}} \right\}, & \lambda(1 - \mu) \leq \mu \leq 1 - \lambda\mu \\ (mb - a) \left\{ \varepsilon_1^{1-\frac{1}{q}} (\delta_1 |f'(a)|^q + m\delta_2 |f'(b)|^q)^{\frac{1}{q}} \right. \\ \quad \left. + \varepsilon_3^{1-\frac{1}{q}} (\beta_1 |f'(a)|^q + m\beta_2 |f'(b)|^q)^{\frac{1}{q}} \right\}, & \mu \leq \lambda(1 - \mu) \leq 1 - \lambda\mu, \\ (mb - a) \left\{ \varepsilon_2^{1-\frac{1}{q}} (\delta_3 |f'(a)|^q + m\delta_4 |f'(b)|^q)^{\frac{1}{q}} \right. \\ \quad \left. + \varepsilon_4^{1-\frac{1}{q}} (\beta_3 |f'(a)|^q + m\beta_4 |f'(b)|^q)^{\frac{1}{q}} \right\}, & \lambda(1 - \mu) \leq 1 - \lambda\mu \leq \mu \end{cases} \end{aligned}$$

$$\begin{aligned}
 & \left| \lambda (\mu f(a) + (1 - \mu)f(mb)) + (1 - \lambda)f(\mu a + m(1 - \mu)b) \right. \\
 & \left. - \frac{1}{mb - a} \int_a^{mb} f(x) dx \right| \\
 & \leq (mb - a) \left[\int_0^\mu |-t + \lambda(1 - \mu)| |f'(ta + m(1 - t)b)| dt \right. \\
 & \quad \left. + \int_\mu^1 |-t + (1 - \lambda\mu)| |f'(ta + m(1 - t)b)| dt \right] \\
 & \leq (mb - a) \left\{ \left(\int_0^\mu |-t + \lambda(1 - \mu)| dt \right)^{1 - \frac{1}{q}} \right. \\
 & \quad \times \left(\int_0^\mu |-t + \lambda(1 - \mu)| |f'(ta + m(1 - t)b)|^q dt \right)^{\frac{1}{q}} \\
 & \quad \left. + \left(\int_\mu^1 |-t + (1 - \lambda\mu)| dt \right)^{1 - \frac{1}{q}} \right. \\
 & \quad \left. \times \left(\int_\mu^1 |-t + (1 - \lambda\mu)| |f'(ta + m(1 - t)b)|^q dt \right)^{\frac{1}{q}} \right\} \tag{5}
 \end{aligned}$$

Since $|f'|^q$ is (α, m) -convex on $[a, b]$, we know that for $t \in [0, 1]$

$$|f'(ta + m(1 - t)b)|^q \leq t^\alpha |f'(a)|^q + m(1 - t^\alpha) |f'(b)|^q,$$

hence, by simple computation

$$\int_0^\mu |-t + \lambda(1 - \mu)| dt = \begin{cases} \varepsilon_1, & \mu \leq \lambda(1 - \mu) \\ \varepsilon_2, & \mu \geq \lambda(1 - \mu) \end{cases}, \tag{6}$$

$$\int_\mu^1 |-t + (1 - \lambda\mu)| dt = \begin{cases} \varepsilon_3, & \mu \leq 1 - \lambda\mu \\ \varepsilon_4, & \mu \geq 1 - \lambda\mu \end{cases}, \tag{7}$$

$$\begin{aligned}
 & \int_0^\mu |-t + \lambda(1 - \mu)| |f'(ta + m(1 - t)b)|^q dt \\
 & \leq \int_0^\mu |-t + \lambda(1 - \mu)| [t^\alpha |f'(a)|^q + m(1 - t^\alpha) |f'(b)|^q] dt \\
 & = \begin{cases} \delta_1 |f'(a)|^q + m\delta_2 |f'(b)|^q, & \mu \leq \lambda(1 - \mu) \\ \delta_3 |f'(a)|^q + m\delta_4 |f'(b)|^q, & \mu \geq \lambda(1 - \mu) \end{cases}, \tag{8} \\
 & \int_\mu^1 |-t + (1 - \lambda\mu)| |f'(ta + m(1 - t)b)|^q dt
 \end{aligned}$$

$$\begin{aligned}
 & \leq \int_\mu^1 |-t + (1 - \lambda\mu)| [t^\alpha |f'(a)|^q + m(1 - t^\alpha) |f'(b)|^q] dt \\
 & = \begin{cases} \beta_1 |f'(a)|^q + m\beta_2 |f'(b)|^q, & \mu \leq 1 - \lambda\mu \\ \beta_3 |f'(a)|^q + m\beta_4 |f'(b)|^q, & \mu \geq 1 - \lambda\mu \end{cases}. \tag{9}
 \end{aligned}$$

Thus, using (6)-(9) in (5), we obtain the inequality (4). This completes the proof. \square

Corollary 2.1. Under the assumptions of Theorem 2 with $q = 1$, we have

$$\begin{aligned}
 & \left| \lambda (\mu f(a) + (1 - \mu)f(mb)) + (1 - \lambda)f(\mu a + m(1 - \mu)b) \right. \\
 & \left. - \frac{1}{mb - a} \int_a^{mb} f(x) dx \right| \\
 & \leq \begin{cases} (mb - a) \{ (\delta_3 + \beta_1) |f'(a)| + m(\delta_4 + \beta_2) |f'(b)| \}, & \lambda(1 - \mu) \leq \mu \leq 1 - \lambda\mu \\ (mb - a) \{ (\delta_1 + \beta_1) |f'(a)| + m(\delta_2 + \beta_2) |f'(b)| \}, & \mu \leq \lambda(1 - \mu) \leq 1 - \lambda\mu \\ (mb - a) \{ (\delta_3 + \beta_3) |f'(a)| + m(\delta_4 + \beta_4) |f'(b)| \}, & \lambda(1 - \mu) \leq 1 - \lambda\mu \leq \mu \end{cases}.
 \end{aligned}$$

Remark 1. In Corollary 2.1,

(i) If we choose $\mu = \frac{1}{2}$, $\lambda = \frac{1}{3}$ and $\alpha = 1$, we have the following inequality

$$\begin{aligned}
 & \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a + mb}{2}\right) + f(mb) \right] - \frac{1}{mb - a} \int_a^{mb} f(x) dx \right| \\
 & \leq \frac{5}{72} (mb - a) \{ |f'(a)| + m |f'(b)| \},
 \end{aligned}$$

which is the same of the Simpson type inequality in Corollary 2.3 (ii) in [9].

(ii) If we choose $\mu = \frac{1}{2}$, $\lambda = 1$ and $\alpha = 1$, we have the following inequality

$$\begin{aligned}
 & \left| \frac{f(a) + f(mb)}{2} - \frac{1}{mb - a} \int_a^{mb} f(x) dx \right| \\
 & \leq \frac{(mb - a)}{8} \{ |f'(a)| + m |f'(b)| \}
 \end{aligned}$$

which is the same of the trapezoid type inequality in Corollary 2.3 (i) in [9].

(iii) If we choose $\mu = \frac{1}{2}$, $\lambda = 0$ and $\alpha = 1$, we have the following midpoint type inequality

$$\begin{aligned}
 & \left| f\left(\frac{a + mb}{2}\right) - \frac{1}{mb - a} \int_a^{mb} f(x) dx \right| \\
 & \leq \frac{(mb - a)}{8} \{ |f'(a)| + m |f'(b)| \}.
 \end{aligned}$$

Corollary 2.2. Under the assumptions of Theorem 2 with $\mu = \frac{1}{2}$ and $\lambda = \frac{1}{3}$, we have the following Simpson type inequality

$$\left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+mb}{2}\right) + f(mb) \right] - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right|$$

$$\leq (mb-a) \left(\frac{5}{72}\right)^{1-\frac{1}{q}} \left\{ \left(\delta_3^* |f'(a)|^q + m \left(\frac{5}{72} - \delta_3^*\right) |f'(b)|^q \right)^{\frac{1}{q}} \right.$$

$$\left. + \left(\beta_1^* |f'(a)|^q + m \left(\frac{5}{72} - \beta_1^*\right) |f'(b)|^q \right)^{\frac{1}{q}} \right\},$$

where

$$\delta_3^* = \frac{2 + 3^{\alpha+1} (2\alpha + 1)}{6^{\alpha+2} (\alpha + 1) (\alpha + 2)}$$

and

$$\beta_1^* = \frac{2 \times 5^{\alpha+2} + 6^{\alpha+1} (\alpha - 4) - 3^{\alpha+1} (2\alpha + 7)}{6^{\alpha+2} (\alpha + 1) (\alpha + 2)}.$$

Remark 2. In Corollary 2.2, if we take $\alpha = m = 1$, we obtain the following Simpson type inequality

$$\left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{b-a} \int_a^b f(x) dx \right|$$

$$\leq (b-a) \left(\frac{5}{72}\right)^{1-\frac{1}{q}} \left\{ \left(\frac{29}{1296} |f'(b)|^q + \frac{61}{1296} |f'(a)|^q \right)^{\frac{1}{q}} \right.$$

$$\left. + \left(\frac{61}{1296} |f'(b)|^q + \frac{29}{1296} |f'(a)|^q \right)^{\frac{1}{q}} \right\},$$

which is the same of the inequality in Theorem 10 in [11] for $s = 1$.

Corollary 2.3. Under the assumptions of Theorem 2 with $\mu = \frac{1}{2}$ and $\lambda = 1$, we have the following trapezoid type inequality

$$\left| \frac{f(a) + f(mb)}{2} - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right| \leq (mb-a) \left(\frac{1}{8}\right)^{1-\frac{1}{q}}$$

$$\times \left\{ \left(\frac{1}{2^{\alpha+2} (\alpha + 1) (\alpha + 2)} |f'(a)|^q \right. \right.$$

$$\left. + m \left(\frac{1}{8} - \frac{1}{2^{\alpha+2} (\alpha + 1) (\alpha + 2)} \right) |f'(b)|^q \right)^{\frac{1}{q}}$$

$$+ \left(\frac{\alpha 2^{\alpha+1} + 1}{2^{\alpha+2} (\alpha + 1) (\alpha + 2)} |f'(a)|^q \right.$$

$$\left. + m \left(\frac{1}{8} - \frac{\alpha 2^{\alpha+1} + 1}{2^{\alpha+2} (\alpha + 1) (\alpha + 2)} \right) |f'(b)|^q \right)^{\frac{1}{q}} \left. \right\}.$$

Corollary 2.4. Under the assumptions of Theorem 2 with $\mu = \frac{1}{2}$ and $\lambda = 0$, we have the following midpoint type inequality

$$\left| f\left(\frac{a+mb}{2}\right) - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right|$$

$$\leq (mb-a) \left(\frac{1}{8}\right)^{1-\frac{1}{q}} \times \left\{ \left(\frac{1}{2^{\alpha+2} (\alpha + 2)} |f'(a)|^q \right. \right.$$

$$\left. + m \left(\frac{1}{8} - \frac{1}{2^{\alpha+2} (\alpha + 2)} \right) |f'(b)|^q \right)^{\frac{1}{q}}$$

$$+ \left(\frac{2^{\alpha+2} - \alpha - 3}{2^{\alpha+2} (\alpha + 1) (\alpha + 2)} |f'(a)|^q \right.$$

$$\left. + m \left(\frac{1}{8} - \frac{2^{\alpha+2} - \alpha - 3}{2^{\alpha+2} (\alpha + 1) (\alpha + 2)} \right) |f'(b)|^q \right)^{\frac{1}{q}} \left. \right\}$$

Theorem 3. Let $f : I \subseteq [0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a, b]$, where $a, b \in I^\circ$ with $a < b$ and $\lambda, \mu \in [0, 1]$. If $|f'|^q$ is (α, m) -convex on $[a, b]$, for $(\alpha, m) \in (0, 1]^2$, $mb > a$, $q > 1$, then the following inequality holds:

$$\left| \lambda (\mu f(a) + (1-\mu)f(mb)) + (1-\lambda)f(\mu a + m(1-\mu)b) \right.$$

$$\left. - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right|$$

$$(10)$$

$$\leq (mb-a) \left(\frac{1}{p+1}\right)^{\frac{1}{p}}$$

$$\left\{ \begin{array}{l} \left[\vartheta_1^{\frac{1}{p}} A^{\frac{1}{q}} + \vartheta_3^{\frac{1}{p}} B^{\frac{1}{q}} \right], \lambda(1-\mu) \leq \mu \leq 1-\lambda\mu \\ \left[\vartheta_2^{\frac{1}{p}} A^{\frac{1}{q}} + \vartheta_3^{\frac{1}{p}} B^{\frac{1}{q}} \right], \mu \leq \lambda(1-\mu) \leq 1-\lambda\mu \\ \left[\vartheta_2^{\frac{1}{p}} A^{\frac{1}{q}} + \vartheta_4^{\frac{1}{p}} B^{\frac{1}{q}} \right], \lambda(1-\mu) \leq 1-\lambda\mu \leq \mu, \end{array} \right.$$

where

$$A = \mu \times \min \left\{ \frac{|f'(\mu a + m(1-\mu)b)|^q + \alpha m |f'(b)|^q}{\alpha + 1}, \frac{|f'(mb)|^q + \alpha m \left| f' \left(\frac{\mu a + m(1-\mu)b}{m} \right) \right|^q}{\alpha + 1} \right\}$$

$$B = (1-\mu) \times \min \left\{ \frac{|f'(\mu a + m(1-\mu)b)|^q + \alpha m \left| f' \left(\frac{a}{m} \right) \right|^q}{\alpha + 1}, \frac{|f'(a)|^q + \alpha m \left| f' \left(\frac{\mu a + m(1-\mu)b}{m} \right) \right|^q}{\alpha + 1} \right\}$$

$$\vartheta_1 = [\lambda(1-\mu)]^{p+1} + [\mu - \lambda(1-\mu)]^{p+1},$$

$$\vartheta_2 = [\lambda(1-\mu)]^{p+1} - [\lambda(1-\mu) - \mu]^{p+1},$$

$$\vartheta_3 = [1 - \lambda\mu - \mu]^{p+1} + [\lambda\mu]^{p+1},$$

$$\vartheta_4 = [\lambda\mu]^{p+1} - [\mu - 1 + \lambda\mu]^{p+1},$$

and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. From Lemma 1 and by Hölder's integral inequality, we have

$$\left| \lambda(\mu f(a) + (1-\mu)f(mb)) + (1-\lambda)f(\mu a + m(1-\mu)b) - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right|$$

$$\leq (mb-a) \left[\int_0^\mu |-t + \lambda(1-\mu)| |f'(ta + m(1-t)b)| dt + \int_\mu^1 |-t + (1-\mu\lambda)| |f'(ta + m(1-t)b)| dt \right]$$

$$\leq (mb-a) \left\{ \left(\int_0^\mu |-t + \lambda(1-\mu)|^p dt \right)^{\frac{1}{p}} \times \left(\int_0^\mu |f'(ta + m(1-t)b)|^q dt \right)^{\frac{1}{q}} + \left(\int_\mu^1 |-t + (1-\lambda\mu)|^p dt \right)^{\frac{1}{p}} \times \left(\int_\mu^1 |f'(ta + m(1-t)b)|^q dt \right)^{\frac{1}{q}} \right\}$$

(11)

Since $|f'|^q$ is (α, m) -convex on $[a, b]$, for $(\alpha, m) \in (0, 1]^2$ and $\mu \in (0, 1]$ by the inequality (2), we get

$$\int_0^\mu |f'(ta + m(1-t)b)|^q dt = \mu \left[\frac{1}{\mu(mb-a)} \int_{\mu a + m(1-\mu)b}^{mb} |f'(x)|^q dx \right]$$

$$\leq \mu \times \min \left\{ \frac{|f'(\mu a + m(1-\mu)b)|^q + \alpha m |f'(b)|^q}{\alpha + 1}, \frac{|f'(mb)|^q + \alpha m \left| f' \left(\frac{\mu a + m(1-\mu)b}{m} \right) \right|^q}{\alpha + 1} \right\}.$$

(12)

The inequality (12) also holds for $\mu = 0$. Similarly, for $\mu \in [0, 1)$ by the inequality (2), we have

$$\int_\mu^1 |f'(ta + m(1-t)b)|^q dt = (1-\mu) \left[\frac{1}{(1-\mu)(mb-a)} \int_a^{\mu a + m(1-\mu)b} |f'(x)|^q dx \right]$$

$$\leq (1-\mu) \times \min \left\{ \frac{|f'(\mu a + m(1-\mu)b)|^q + \alpha m \left| f' \left(\frac{a}{m} \right) \right|^q}{\alpha + 1}, \frac{|f'(a)|^q + \alpha m \left| f' \left(\frac{\mu a + m(1-\mu)b}{m} \right) \right|^q}{\alpha + 1} \right\}.$$

(13)

The inequality (12) also holds for $\mu = 1$. By simple computation

$$\int_0^\mu |-t + \lambda(1-\mu)|^p dt = \begin{cases} \frac{[\lambda(1-\mu)]^{p+1} + [\mu - \lambda(1-\mu)]^{p+1}}{p+1}, & \lambda(1-\mu) \leq \mu \\ \frac{[\lambda(1-\mu)]^{p+1} - [\lambda(1-\mu) - \mu]^{p+1}}{p+1}, & \lambda(1-\mu) \geq \mu \end{cases},$$

(14)

and

$$\int_\mu^1 |-t + (1-\lambda\mu)|^p dt = \begin{cases} \frac{[1-\lambda\mu-\mu]^{p+1} + [\lambda\mu]^{p+1}}{p+1}, & \mu \leq 1-\lambda\mu \\ \frac{[\lambda\mu]^{p+1} - [\mu-1+\lambda\mu]^{p+1}}{p+1}, & \mu \geq 1-\lambda\mu \end{cases};$$

(15)

thus, using (12)-(15) in (11), we obtain the inequality (10). This completes the proof. \square

Corollary 2.5. *Under the assumptions of Theorem 3 with $\mu = \frac{1}{2}$ and $\lambda = \frac{1}{3}$, we have the following Simpson type inequality*

$$\left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+mb}{2}\right) + f(mb) \right] - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right| \leq \left(\frac{mb-a}{12} \right) \left(\frac{2^{p+1}+1}{3(p+1)} \right)^{\frac{1}{p}} \left(A_1^{\frac{1}{q}} + B_1^{\frac{1}{q}} \right),$$

where

$$A_1 = \min \left\{ \frac{|f'(\frac{a+mb}{2})|^q + \alpha m |f'(b)|^q}{\alpha + 1}, \frac{|f'(mb)|^q + \alpha m |f'(\frac{a+mb}{2m})|^q}{\alpha + 1} \right\},$$

and

$$B_1 = \min \left\{ \frac{|f'(\frac{a+mb}{2})|^q + \alpha m |f'(\frac{a}{m})|^q}{\alpha + 1}, \frac{|f'(a)|^q + \alpha m |f'(\frac{a+mb}{2m})|^q}{\alpha + 1} \right\}.$$

Remark 3. In Corollary 2.5, if we take $\alpha = m = 1$, then we obtain the following Simpson type inequality

$$\left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+b}{2}\right) + f(b) \right] - \frac{1}{b-a} \int_a^b f(x) dx \right| \leq \left(\frac{b-a}{12} \right) \left(\frac{1+2^{p+1}}{3(p+1)} \right)^{\frac{1}{p}} \times \left\{ \left(\frac{|f'(\frac{a+b}{2})|^q + |f'(a)|^q}{2} \right)^{\frac{1}{q}} + \left(\frac{|f'(\frac{a+b}{2})|^q + |f'(b)|^q}{2} \right)^{\frac{1}{q}} \right\},$$

which is the same of the inequality in Corollary 3 in [11].

Corollary 2.6. *Under the assumptions of Theorem 3 with $\mu = \frac{1}{2}$ and $\lambda = 1$, we have the following trapezoid type inequality*

$$\left| \frac{f(a) + f(mb)}{2} - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right| \leq \left(\frac{mb-a}{4} \right) \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(A_1^{\frac{1}{q}} + B_1^{\frac{1}{q}} \right).$$

Corollary 2.7. *Under the assumptions of Theorem 3 with $\mu = \frac{1}{2}$ and $\lambda = 0$, we obtain the following midpoint type inequality*

$$\left| f\left(\frac{a+mb}{2}\right) - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right| \leq \left(\frac{mb-a}{4} \right) \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \left(A_1^{\frac{1}{q}} + B_1^{\frac{1}{q}} \right).$$

Theorem 4. *Let $f : I \subseteq [0, \infty) \rightarrow \mathbb{R}$ be a differentiable mapping on I° such that $f' \in L[a, b]$, where $a, b \in I^\circ$ with $a < b$ and $\lambda, \mu \in [0, 1]$. If $|f'|^q$ is (α, m) -convex on $[a, b]$, for $(\alpha, m) \in (0, 1]^2$, $mb > a$, $q > 1$, then the following inequality holds:*

$$\left| \lambda (\mu f(a) + (1-\mu)f(mb)) + (1-\lambda)f(\mu a + m(1-\mu)b) - \frac{1}{mb-a} \int_a^{mb} f(x) dx \right| \leq (mb-a) \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \times \begin{cases} \left[\vartheta_1^{\frac{1}{p}} A_2^{\frac{1}{q}} + \vartheta_3^{\frac{1}{p}} B_2^{\frac{1}{q}} \right], & \lambda(1-\mu) \leq \mu \leq 1-\lambda\mu \\ \left[\vartheta_2^{\frac{1}{p}} A_2^{\frac{1}{q}} + \vartheta_3^{\frac{1}{p}} B_2^{\frac{1}{q}} \right], & \mu \leq \lambda(1-\mu) \leq 1-\lambda\mu \\ \left[\vartheta_2^{\frac{1}{p}} A_2^{\frac{1}{q}} + \vartheta_4^{\frac{1}{p}} B_2^{\frac{1}{q}} \right], & \lambda(1-\mu) \leq 1-\lambda\mu \leq \mu \end{cases} \quad (16)$$

where

$$A_2 = \frac{\mu^{\alpha+1} |f'(a)|^q + m [\mu(\alpha+1) - \mu^{\alpha+1}] |f'(b)|^q}{\alpha + 1}$$

and

$$B_2 = \frac{(1-\mu^{\alpha+1}) |f'(a)|^q + m [(\mu^{\alpha+1}-1) + (1-\mu)(\alpha+1)] |f'(b)|^q}{\alpha + 1}.$$

Proof. From Lemma 1 and by Hölder's integral inequality, we have inequality (11). Since $|f'|^q$ is (α, m) -convex on $[a, b]$, we know that for $t \in [0, \mu]$ and $t \in [\mu, 1]$

$$|f'(ta + m(1-t)b)|^q \leq t^\alpha |f'(a)|^q + m(1-t^\alpha) |f'(b)|^q.$$

Hence,

$$\begin{aligned} & \left| \lambda (\mu f(a) + (1-\mu)f(mb)) + (1-\lambda)f(\mu a + m(1-\mu)b) - \frac{1}{mb-a} \int_a^{mb} f(x)dx \right| \\ & \leq (mb-a) \left\{ \left(\int_0^\mu | -t + \lambda(1-\mu) |^p dt \right)^{\frac{1}{p}} \left(\int_0^\mu [t^\alpha |f'(a)|^q + m(1-t^\alpha) |f'(b)|^q] dt \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\int_\mu^1 | -t + (1-\alpha\lambda) |^p dt \right)^{\frac{1}{p}} \left(\int_\mu^1 [t^\alpha |f'(a)|^q + m(1-t^\alpha) |f'(b)|^q] dt \right)^{\frac{1}{q}} \right\} \tag{17} \\ & \leq (mb-a) \left\{ \left(\int_0^\mu | -t + \lambda(1-\mu) |^p dt \right)^{\frac{1}{p}} \left(\frac{\mu^{\alpha+1} |f'(a)|^q + m[\mu(\alpha+1) - \mu^{\alpha+1}] |f'(b)|^q}{\alpha+1} \right)^{\frac{1}{q}} \right. \\ & \quad \left. + \left(\int_\mu^1 | -t + (1-\alpha\lambda) |^p dt \right)^{\frac{1}{p}} \left(\frac{(1-\mu^{\alpha+1}) |f'(a)|^q + m[(\mu^{\alpha+1}-1) + (1-\mu)(\alpha+1)] |f'(b)|^q}{\alpha+1} \right)^{\frac{1}{q}} \right\}. \end{aligned}$$

Thus, using (14), (15) in (17), we obtain the inequality (16). This completes the proof. \square

Corollary 2.10. *In Corollary 2.9, if we take $\alpha = 1$ we obtain the following trapezoid type inequality*

Corollary 2.8. *Let the assumptions of Theorem 4 hold. Then for $\mu = \frac{1}{2}$ and $\lambda = \frac{1}{3}$ from the inequality (16), we get the following Simpson type inequality*

$$\begin{aligned} & \left| \frac{1}{6} \left[f(a) + 4f\left(\frac{a+mb}{2}\right) + f(mb) \right] - \frac{1}{mb-a} \int_a^{mb} f(x)dx \right| \\ & \leq \left(\frac{mb-a}{12} \right) \left(\frac{2^{p+1}+1}{3(p+1)} \right)^{\frac{1}{p}} \left(\frac{1}{2^\alpha(\alpha+1)} \right)^{\frac{1}{q}} \left(A_3^{\frac{1}{q}} + B_3^{\frac{1}{q}} \right). \end{aligned}$$

where

$$A_3 = |f'(a)|^q + m[2^\alpha(\alpha+1) - 1] |f'(b)|^q$$

and

$$B_3 = (2^{\alpha+1} - 1) |f'(a)|^q + m[2^\alpha(\alpha+1) + 1 - 2^{\alpha+1}] |f'(b)|^q.$$

Corollary 2.9. *Let the assumptions of Theorem 4 hold. Then for $\mu = \frac{1}{2}$ and $\lambda = 1$ from the inequality (16), we get the following trapezoid type inequality:*

$$\begin{aligned} & \left| \frac{f(a) + f(mb)}{2} - \frac{1}{mb-a} \int_a^{mb} f(x)dx \right| \\ & \leq \left(\frac{mb-a}{4} \right) \left(\frac{1}{(p+1)} \right)^{\frac{1}{p}} \left(\frac{1}{2^\alpha(\alpha+1)} \right)^{\frac{1}{q}} \left(A_3^{\frac{1}{q}} + B_3^{\frac{1}{q}} \right). \end{aligned}$$

$$\begin{aligned} & \left| \frac{f(a) + f(mb)}{2} - \frac{1}{mb-a} \int_a^{mb} f(x)dx \right| \leq (mb-a) \left(\frac{1}{p+1} \right)^{\frac{1}{p}} \\ & \quad \times \left(\frac{1}{4} \right)^{1+\frac{1}{q}} \left[(|f'(a)|^q + 3m|f'(b)|^q)^{\frac{1}{q}} \right. \\ & \quad \left. + (3|f'(a)|^q + m|f'(b)|^q)^{\frac{1}{q}} \right] \\ & \leq (mb-a) \left(\frac{1}{4} \right)^{1+\frac{1}{q}} \left[(|f'(a)|^q + 3m|f'(b)|^q)^{\frac{1}{q}} \right. \\ & \quad \left. + (3|f'(a)|^q + m|f'(b)|^q)^{\frac{1}{q}} \right], \tag{18} \end{aligned}$$

where we have used the fact that $1/2 < (1/(p+1))^{1/p} < 1$. We note that the inequality (18) is the same of the inequality in Corollary 2.7 (i) [9].

Corollary 2.11. *Under the assumptions of Theorem 4 with $\mu = \frac{1}{2}$ and $\lambda = 0$, we have the following midpoint type inequality:*

$$\begin{aligned} & \left| f\left(\frac{a+mb}{2}\right) - \frac{1}{mb-a} \int_a^{mb} f(x)dx \right| \\ & \leq \left(\frac{mb-a}{4} \right) \left(\frac{1}{(p+1)} \right)^{\frac{1}{p}} \left(\frac{1}{2^\alpha(\alpha+1)} \right)^{\frac{1}{q}} \left(A_3^{\frac{1}{q}} + B_3^{\frac{1}{q}} \right). \end{aligned}$$

Competing interest

The author declares that he has no competing interest.

Acknowledgements

The author would like to thank to referees for their valuable suggestion and comments.

Received: 18 March 2013 Accepted: 21 April 2013

Published: 6 May 2013

References

1. Miheşan, VG: A generalization of the convexity. In: *Seminar on Functional Equations, Approximation and Convexity*. Cluj-Napoca, Romania, (1993)
2. Bakula, MK, Ozdemir, ME, Pecaric J: Hadamard type inequalities for m -convex and (α, m) -convex functions. *J. Inequal. Pure Appl. Math.* **9**(4), 12 (2008). [http://www.emis.de/journals/JIPAM/article1032.html]
3. İşcan, İ: Hermite-Hadamard type inequalities for functions whose derivatives are (α, m) -convex. *Int. J. Eng. Appl. Sci.* **2**(3), 69–78 (2013)
4. Ozdemir, ME, Avcı, M, Kavurmacı, H: Hermite-Hadamard-type inequalities via (α, m) -convexity. *Comput Math. Appl.* **61**, 2614–2620 (2011)
5. Ozdemir, ME, Kavurmacı, H, Set, E: Ostrowski's type inequalities for (α, m) -convex functions. *Kyungpook. Math. J.* **50**, 371–378 (2010)
6. Ozdemir, ME, Set, E, Sankaya, MZ: Some new Hadamard's type inequalities for co-ordinated m -convex and (α, m) -convex functions. *Hacetatepe J. Math. Stat.* **40**(2), 219–229 (2011)
7. Set, E, Sardari, M, Ozdemir, ME, Rooin, J: On generalizations of the Hadamard inequality for (α, m) -convex functions. *RGMI Res. Rep. Coll. Article.* **12**(4), 4 (2009)
8. Yildiz, C, Gurbuz, M, Akdemir, AO: The Hadamard type inequalities for m -convex functions. *Konuralp J. Math.* **1**(1), 40–47 (2013)
9. Park, J: Hermite-Hadamard and Simpson-like type inequalities for differentiable (α, m) -convex mappings. *Int. J. Math. and Mathematical Sciences.* **2012** (2012). 12 Article ID 809689
10. Sarikaya, MZ, Aktan, N: On the generalization of some integral inequalities and their applications. *Math. Comput. Model.* **54**, 2175–2182 (2011)
11. Sarikaya, MZ, Set, E, Özdemir, M E: On new inequalities of Simpson's type for s -convex functions. *Comput Math. Appl.* **60**, 2191–2199 (2010)
12. Sarikaya, MZ, Set, E, Özdemir, M E: On new inequalities of Simpson's type for convex functions. *RGMI Res. Rep. Coll. Article.* **13**(2) (2010)

doi:10.1186/2251-7456-7-22

Cite this article as: İşcan: A new generalization of some integral inequalities for (α, m) -convex functions. *Mathematical Sciences* 2013 **7**:22.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- Convenient online submission
- Rigorous peer review
- Immediate publication on acceptance
- Open access: articles freely available online
- High visibility within the field
- Retaining the copyright to your article

Submit your next manuscript at ► springeropen.com