

Letter to Editor

Incidence of imipenem-resistant *Acinetobacter baumannii* in a general intensive care unit (ICU)

Sir

Hospital-acquired infections in critically ill patients remain one of the most important contributors to morbidity and mortality. Hospital-acquired *A. baumannii* has emerged as a serious threat to ICU patient population. The main purpose of our study was to determine the incidence of ICU acquired imipenem-resistant *A. baumannii* and its impact on ICU mortality.

This prospective study was carried from January 2010 to October 2010 in a general ICU of a major teaching hospital in Tehran, Iran. The patients who had positive culture of *A. baumannii* were included in the study. *A. baumannii* isolates were tested for the sensitivity or resistance to imipenem by E-test. Each test contains 10 mcg of imipenem. Sensitivity to imipenem was defined as less than 4 mcg/ml. Sixty one positive cultures from 61 patients were detected during the study period. The most frequent infection was ventilator-associated pneumonia. Fifty (82.5 %) of isolates were detected in sputum, 5 (8%) in wounds, 3 (4.9%) in urine, 2 (3%) from central venous line and 1 (1.6) in blood. All isolates were imipenem resistant. The resistance level determined by the E-test was 12 mcg/ml in 1% of patients, 16 in 4%, 24 in 5%, and 32 in 88% of isolates. In this study, *A. baumannii* displayed 100% resistance to imipenem. Other study has reported imipenem resistance in the range of 11-24% (1). In a report from 5 European countries, *Acinetobacter* species were found to have the highest increase in resistance to antibiotics among all the gram-negative bacilli studied (2). Several risk factors have been identified for developing infection with multidrug-resistant *A. baumannii*, including admission diagnosis of multiple trauma, mechanical ventilation, length of ICU stay, colonization pressure within the unit and exposure to antimicrobials agents (3- 6).

The alarming imipenem resistance in this study can be due to inappropriate choice of antibiotics and doses. In one large multicenter study in US hospitals, *acinetobacter* blood stream infection was associated with a crude mortality rate of 34% (7). The reviewed data suggest that infection with or the acquisition of *A. baumannii* seems to be associated with increased mortality (8).

A. baumannii has been found on inanimate surfaces even after cleaning with antiseptic solutions (9). Healthcare workers play a potential role in epidemic outbreaks through the contamination of the environment and other patients.

However, this very alarming study conveys important messages that strict infection control measures have to be put in place by infection control department. In summary, considering the unacceptably of the high incidence of imipenem resistant *A. baumannii* infections with its significant morbidity, mortality and costs, it is mandatory to improve infection control strategies and optimize its diagnosis and management.

Citation:

Alavi-Moghadam M, Miri M, Mokhtari M, et al. Incidence of Imipenem Resistant *Acinetobacter baumannii* in a General Intensive Care Unit. *Caspian J Intern Med* 2014; 5 (3): 186-187.

Mostafa Alavi-Moghadam (MD)¹

Mirmohammad Miri (MD)²

Majid Mokhtari (MD)²

Mehran Kouchek (MD)²

Reza Goharani (MD)²

Mohammad Sistanzad (Pharm D, PhD)³

Saeed Safari (MD)¹

Mehrdad Solouki (MD)^{*4}

1. Department of Emergency Medicine, Emam Hossein Medical and Educational Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

2. Department of Critical Care Medicine, Emam Hossein Medical and Educational Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

3. Department of Clinical Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

4. Department of Internal and Critical Care Medicine, Emam Hossein Medical and Educational Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran

Correspondence

Mehrdad Solouki, Department of Internal and Critical Care Medicine, Emam Hossein Medical and Educational Center, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Email: mehrdad2672@yahoo.com

Tel and Fax: 0098 21 88681051

Received: 7 June 2013

Revised: 14 Feb 2014

Accepted: 8 April 2014

References

1. Carbonne A, Naas T, Blanckaert K, et al. Investigation of a nosocomial outbreak of extended-spectrum beta-lactamase VEB-1-producing isolates of *Acinetobacter baumannii* in a hospital setting. *J Hosp Infect* 2005; 60: 14-8.
2. Hanberger H, Garcia-Rodriguez JA, Gobernado M, et al. Antibiotic susceptibility among aerobic gram negative bacilli in ICU in 5 European countries. French and Portuguese ICU study groups. *JAMA* 1999; 28: 67-70.
3. Playford EG, Craig JC, Iredell JR. Carbapenem-resistant *Acinetobacter baumannii* in intensive care unit patients: risk factors for acquisition, infection and their consequences. *J Hosp Infect* 2007; 65: 204-11.
4. Corbella X, Montero A, Pujol M, et al. Emergence and rapid spread of carbapenem resistance during a large and sustained hospital outbreak of multiresistant *Acinetobacter baumannii*. *J Clin Microbiol* 2000; 38: 4086-95.
5. del Mar Tomas M, Cartelle M, Pertega S, et al. Hospital outbreak caused by a carbapenem-resistant strain of *Acinetobacter baumannii*: patient prognosis and risk-factors for colonisation and infection. *Clin Microbiol Infect* 2005; 11: 540-46.
6. Lee SO, Kim NJ, Choi SH, et al. Risk factors for acquisition of imipenem-resistant *Acinetobacter baumannii*: a case-control study. *Antimicrob. Agents Chemother* 2004; 48: 224-28.
7. Wisplinghoff H, Bischoff T, Tallent SM, et al. Nosocomial blood stream infections in US hospitals: analysis of 24,179 cases from a prospective nation wide surveillance study. *Clin Infect Dis* 2004; 39: 309-17.
8. Falagas ME, Bliziotis IA, Siempos II. Attributable mortality of *Acinetobacter baumannii* infections in critically ill patients: a systematic review of matched cohort and case-control studies. *Crit Care* 2006; 10: R48.
9. Roberts SA, Findlay R, Lang SD. Investigation of an outbreak of multi-drug resistant *Acinetobacter baumannii* in an intensive care burns unit. *J Hosp Infect* 2001; 48: 228-32.