Dental Research, Dental Clinics, Dental Prospects

Letter

Update on Prevalence of Minor Aphtha and the Involved

Factors in Tabriz, Northwest Iran

Mahmood Sina^{1*} • Mahmood Toorchi² • Sina Ghertasi Oskouei³

¹Assistant professor, Department of Oral Pathology, Faculty of Dentistry, Tabriz University of Medical Sciences, Tabriz, Iran ²Professor, Department of Biotechnology, Faculty of Agriculture, Tabriz University, Tabriz, Iran ³Faculty of Dentistry, Tabriz University of Medical Sciences, Iran *Corresponding Author; E-mail: sinam@tbzmed.ac.ir

Received: 08 March 2011; Accepted: 12 May 2011 J Dent Res Dent Clin Dent Prospect 2011; 5(3):106-107 This article is available from: http://dentistry.tbzmed.ac.ir/joddd

© 2011 The Authors; Tabriz University of Medical Sciences

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/3.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

In the article entitled "Two-year prevalence of mi-nor aphtha in Tabriz, Northwest Iran" which was published in the winter issue of the third volume of the journal, a 0.3% prevalence rate for minor recurrent aphthous stomatitis (RAS) was reported.¹ We attempted to continue studying the archives of Tabriz University of Medical Sciences Faculty of Dentistry until June 2008, to allow for a consecutive 15000 records. In addition, the clear influence of involved factors in minor RAS was also assessed. The same methods were used in the assessments. Since the response variable of RAS is assumed to be categorical with two response levels. Yes and No, chi-square test was used to determine statistical association of lesions prevalence with gender, age decade, familial involvement, and cigarette smoking. Furthermore, logistic regression model was adapted to investigate the aggregate relationship of RAS with four independent covariates; gender, age decade, family history, and cigarette smoking using stepwise selection method by SAS software (1998, SAS institute, Inc., Cary, USA). The confounding effect of gender was analyzed later by multivariate analysis to minimize selection bias.²

The prevalence of minor aphtha in the new sample

was found to be 0.66%. Aphthous lesion was more prevalent in the third decade of life (28.64%, P < 0.05). Positive history of smoking was seen in 5.85% of the patients who had a lower occurrence of aphthous lesions compared with non-smokers (P = 0.0001). Family history covariate, although significantly associated with RAS prevalence, was omitted from further analysis, since 25% of the contingency table cells had expected counts less than 5 and chisquare test was not valid. The likelihood ratio chisquare for the remaining three categorical covariates including cigarette smoking, gender, and age decade were 57.88 (P = 0.0001), 1.10 (P = 0.2934), and 3.56 (P = 0.0593), respectively. The maximum likelihood estimates of model parameters are given in Table 1.

A recent study in Saudi Arabia found the prevalence of aphthous ulcer to be 0.4% over a three-year period.³ In two surveys in the US children and youth, the prevalence of recurrent aphthous stomatitis was found to be 1.51% and 1.21%.⁴ However, another study on Jordanian population reported a high prevalence rate of 78% for recurrent aphthous ulcerations.⁵ The inconsistency between the results of different studies might be attributed to genetic predisposition and environmental factors affecting different popula-

Table 1. Bivariate and multiple logistic regression models for RAS prevalence

	Bivariate analysis		Multivariate analysis			
Covariate	Parameter estimates	Pr > Chi-square	Stepwise variables entered	Overall fit	Wald Chi-square	Pr
Cigarette	-1.97	0.0001	Cigarette smoking	57.88	79.46	0.0001
smoking						
Gender	0.23	0.3001	Gender	64.71	6.59	0.0102
Age decade	-0.12	0.0591	Age decade	67.78	3.09	0.0788

tions.6

A prevalence rate of 64.73% of minor aphtha in females in our results is consistent with the previous findings which have reported higher incidence of minor aphtha in females.^{7,8}

Among three independent covariates, cigarette smoking was the first variable entered in the model and its relationship with RAS was statistically significant at P < 0.01.Gender was the second categorical covariate entered in the model and its inclusion improved model fit in the presence of cigarette smoking. These results indicate the synergistic interaction effect of cigarette smoking and gender on each other in association with RAS, as indicated with the likelihood ratio chi-square of 64.71 (P < 0.01). Age decade was the last covariate entered in the model and its inclusion could improve the model fit. All the evaluated covariates including age decade, gender, and cigarette smoking seemed to have affected RAS prevalence in the population under investigation.

References

- Sina M, Toorchi M, Vosoughhosseini S, Zenouz AT, Mehdipour M. Two-year prevalence of minor aphtha in Tabriz, Northwest Iran. J Dent Res Dent Clin Dent Prospect 2009;3:17–9.
- Mumcu G, Ergun T, Inanc N, Fresko I, Atalay T, Hayran O, et al. Oral health is impaired in Behçet's disease and is associated with disease severity. *Rheumatology (Oxford)* 2004;43:1028–33.
- Al-Mobeeriek A, AlDosari AM. Prevalence of oral lesions among Saudi dental patients. Ann Saudi Med 2009;29:365–8.
- Shulman JD. An exploration of point, annual, and lifetime prevalence in characterizing recurrent aphthous stomatitis in USA children and youths. *J Oral Pathol Med* 2004;33:558– 66.
- 5. Safadi RA. Prevalence of recurrent aphthous ulceration in Jordanian dental patients, *BMC Oral Health* 2009;9:31.
- 6. Messadi DV, Younai F. Aphthous ulcers. Dermatol Ther 2010; 23:281–90.
- 7. Rivera-Hidalgo F, Shulman JD, Beach MM. The association of tobacco and other factors with recurrent aphthous stomatitis in an US adult population. *Oral Dis* 2004;10:335–45.
- 8. Davatchi F, Tehrani-Banihashemi A, Jamshidi AR, Chams-Davatchi C, Gholami J, Moradi M, et al. The prevalence of oral aphthosis in a normal population in Iran: a WHO-ILAR COPCORD study. *Arch Iran Med* 2008;11:207–9.