

Weighted Sequence Spaces and Cyclicity

J. Doroodgar

Islamic Azad University-Shiraz Branch
Shiraz Farzanegan Pre-University

B. Yousefi

Shiraz University

Abstract: In this paper we investigate the cyclicity of the multiplication operator M_z acting on the weighted Hardy spaces of formal Laurent series.

AMS Subject Classification: Primary 47B37; Secondary 47A16.

Keywords and Phrases: Banach space of Laurent series associated with a sequence β , cyclic vector, multiplication operator, disc algebra.

1. Introduction

Suppose that $1 < p < \infty$ and $\{\beta(n)\}_{n=-\infty}^{\infty}$ denotes a sequence of positive numbers with $\beta(0) = 1$. For a sequence $f = \{\hat{f}(n)\}_{n=-\infty}^{\infty}$, we define

$$\|f\| = \|f\|_p = \left(\sum_{n=-\infty}^{\infty} |\hat{f}(n)|^p |\beta(n)|^p \right)^{\frac{1}{p}}.$$

Furthermore, we shall use the notation $f(z) = \sum_{n=-\infty}^{\infty} \hat{f}(n)z^n$ regardless whether the series converges for any complex value of z . Throughout

this article, by the space $L^p(\beta)$ we mean

$$L^p(\beta) = \left\{ f : f(z) = \sum_{n=-\infty}^{\infty} \hat{f}(n)z^n, \|f\|_p < \infty \right\}$$

which is called a weighted Hardy space of formal Laurent series (note that when n ranges on $\mathbb{N} \cup \{0\}$, it is called a weighted Hardy space of formal power series and is denoted by $H^p(\beta)$). These are reflexive Banach spaces with the norm $\|\cdot\|_\beta$. Let $\hat{f}_k(n) = \delta_k(n)$. So $f_k(z) = z^k$ and then $\{f_k\}_{k=-\infty}^{\infty}$ is a basis for $L^p(\beta)$ such that $\|f_k\| = \beta(k)$. Now consider M_z , the operator of multiplication by z on $L^p(\beta)$:

$$(M_z f)(z) = \sum_{n=-\infty}^{\infty} \hat{f}(n)z^{n+1}$$

where

$$f(z) = \sum_{n=-\infty}^{\infty} \hat{f}(n)z^n \in L^p(\beta).$$

In other words $(\widehat{M_z f})(n) = \hat{f}(n-1)$ for all $n \in \mathbb{Z}$. Clearly M_z shifts the basis $\{f_k\}_k$. The operator M_z is bounded if and only if $\{\beta(k+1)/\beta(k)\}_k$ is bounded and in this case

$$\|M_z^n\| = \sup_k [\beta(k+n)/\beta(k)]$$

for all $n \in \mathbb{N} \cup \{0\}$.

By the same method used in [3] we can see that $L^p(\beta)^* = L^q(\beta^{\frac{p}{q}})$, where $\frac{1}{p} + \frac{1}{q} = 1$. Also if

$$f(z) = \sum_{n=-\infty}^{\infty} \hat{f}(n)z^n \in L^p(\beta)$$

and

$$g(z) = \sum_{n=-\infty}^{\infty} \hat{g}(n)z^n \in L^q(\beta^{\frac{p}{q}}),$$

then clearly

$$\langle f, g \rangle = \sum_{n=-\infty}^{\infty} \hat{f}(n)\overline{\hat{g}(n)}\beta(n)^p$$

and

$$\begin{aligned} \|g\|_q^q &= \sum_{n=-\infty}^{\infty} |\hat{g}(n)|^q (\beta(n)^{\frac{p}{q}})^q \\ &= \sum_{n=-\infty}^{\infty} |\hat{g}(n)|^q \beta(n)^p \end{aligned}$$

(see [3]). Here for simplicity we used $\|g\|_q$ instead of $\|g\|_{L^q(\beta^{\frac{p}{q}})}$. For some topics on these spaces see [2–16].

Let X be a Banach space. We denote by $B(X)$, the set of bounded operators on the Banach space X . Let $A \in B(X)$ and $x \in X$. We say that x is a cyclic vector of A if X is equal to the closed linear span of the set

$$\{A^n x : n = 0, 1, 2, \dots\}.$$

An operator $A \in B(X)$ is called cyclic if it has a cyclic vector.

If X is a Banach space, it is convenient and helpful to introduce the notation (x, x^*) to stand for $x^*(x)$, for $x \in X$ and $x^* \in X^*$.

In [3] and [5] we studied the cyclicity of the multiplication operator M_z on $H^p(\beta)$ and here we want to investigate the cyclicity of the multiplication operator M_z on the both spaces $H^p(\beta)$ and $L^p(\beta)$.

2. Main Results

First we note that the multiplication operator M_z on $L^p(\beta)$ ($H^p(\beta)$) is unitarily equivalent to an injective bilateral (unilateral) weighted shift and conversely, every injective bilateral (unilateral) weighted shift is unitarily equivalent to M_z acting on $L^p(\beta)$ ($H^p(\beta)$) for a suitable choice of β (the proof is similar to the case $p=2$ that was proved in [2]).

We will use the following notations:

$$r_0 = \overline{\lim} \beta(-n)^{-1/n},$$

$$r_1 = \underline{\lim} \beta(n)^{1/n},$$

$$\Omega_0 = \{z \in \mathbf{C} : |z| > r_0\},$$

$$\Omega_1 = \{z \in \mathbf{C} : |z| < r_1\},$$

$$\Omega = \Omega_0 \cap \Omega_1.$$

From now on we consider that M_z is bounded on $L^p(\beta)$.

Theorem 1. *Let $0 < r_0 < r_1 = 1$ and $\frac{1}{p} + \frac{1}{q} = 1$. If*

$$\sum_{n<0} \frac{r_0^{nq}}{\beta(n)^q} < \infty \quad ; \quad \sum_{n \geq 0} \frac{1}{\beta(n)^q} < \infty,$$

then M_z has no cyclic vector on $L^p(\beta)$.

Proof. Note that Ω is an annulus with the unit disc as an outer boundary. Now for any function

$$f = \sum_{n=-\infty}^{\infty} \hat{f}(n) f_n$$

in $L^p(\beta)$, by the Holder inequality we have

$$\begin{aligned} \sum_{n=-\infty}^{\infty} |\hat{f}(n)| |z|^n &\leq \left(\sum_{n=-\infty}^{\infty} |\hat{f}(n)|^p \beta(n)^p \right)^{1/p} \left(\sum_{n=-\infty}^{\infty} \frac{|z|^{nq}}{\beta(n)^q} \right)^{1/q} \\ &= \|f\|_p \left[\left(\sum_{n<0} \frac{|z|^{nq}}{\beta(n)^q} \right)^{1/q} + \left(\sum_{n \geq 0} \frac{|z|^{nq}}{\beta(n)^q} \right)^{1/q} \right] \\ &\leq \|f\|_p \left[\left(\sum_{n<0} \frac{r_0^{nq}}{\beta(n)^q} \right)^{1/q} + \left(\sum_{n \geq 0} \frac{1}{\beta(n)^q} \right)^{1/q} \right] \end{aligned}$$

for all z in Ω . Since

$$\sum_{n<0} \frac{r_0^{nq}}{\beta(n)^q} < \infty \quad ; \quad \sum_{n \geq 0} \frac{1}{\beta(n)^q} < \infty,$$

by a similar method used in the proof of Theorem 3 in [3] and Theorem 1 in [7], we get $H^p(\beta) \subset H(\Omega) \cap C(T)$ where $H(\Omega)$ is the set of analytic

functions on Ω and $C(T)$ is the set of continuous functions on the unit circle T .

Now define the operator

$$L : L^p(\beta) \longrightarrow C(T) \quad ; \quad L(f) = f|_T.$$

Clearly L maps the set of all Laurent polynomials onto the set of all polynomials in z and \bar{z} which is dense in $C(T)$ by the Stone-Weierstrass theorem. Thus L has dense range. Now if g is a cyclic vector for M_z as an operator on $L^p(\beta)$, then $g|_T$ is a cyclic vector for M_z as an operator on $C(T)$. Thus g has no zero on T and this implies that the operator M_g is invertible on $C(T)$. Let $V\{.\}$ denotes the uniform closed linear span of the set $\{.\}$ in $C(T)$. Clearly $V\{M_z^n g|_T : n \geq 0\}$ is equal to the uniform closure of the set

$$\{pg|_T : p \text{ is an analytic polynomial}\}.$$

Thus, we get

$$V\{M_z^n g|_T : n \geq 0\} = M_g A$$

where A is the disc algebra of analytic functions in $C(T)$. Indeed

$$A = \text{uniform-closure } \{p|_T : p \text{ is an analytic polynomial}\}$$

(see [1]). But A is a proper closed subspace of $C(T)$, so $M_g A$ is also a proper closed subspace of $C(T)$, since M_g is invertible on $C(T)$. This says that $g|_T$ can not be a cyclic vector for M_z as an operator on $C(T)$, hence g can not be a cyclic vector for M_z as an operator on $L^p(\beta)$ that is a contradiction. Now the proof is complete. \square

Theorem 2. *i) If a function f in $H^p(\beta)$ is cyclic, then the zeros of f can not belong to Ω_1 .*

ii) If the zeros of a polynomial P are not belong to Ω_1 , then P is a cyclic vector for M_z .

Proof. See [3]. \square

By the same method we can have a similar result for the spaces $L^p(\beta)$ and in this case we should use Ω instead of Ω_1 .

Theorem 3. *Let $1 \leq p < \infty$. Suppose that $\beta(n)$ is in the form $\beta(n) = \alpha(n)\gamma(n)$ where $\{\alpha(n)\}$ and $\{\gamma(n)\}$ satisfies:*

i) There exists a positive number M , such that

$$\sup\left\{\left|\frac{\gamma(n+i)}{\gamma(n)\gamma(i)}\right| : i, n = 0, 1, 2, \dots\right\} \leq M$$

ii) There exists a positive integer m_0 such that:

$$L_{m_0} = \sup\left\{\left|\frac{\alpha(n+i)\alpha(m_0)}{\alpha(n+m_0)\alpha(i)}\right| : n > 0, i \geq m_0\right\} < \infty$$

and

$$\left\{ \frac{\alpha(n + m_0)}{\alpha(n)} \right\}_n \in \ell^q,$$

where $\frac{1}{p} + \frac{1}{q} = 1$.

If $x = \sum_m x_m f_m$ belongs to $H^p(\beta)$ and $x_0 \neq 0$, then x is a cyclic vector of M_z as an operator on $H^p(\beta)$.

Proof. See [5].□

Corollary 4. Let $\frac{1}{p} + \frac{1}{q} = 1$, M_z be power bounded and $f = \sum_{m=0}^{\infty} \hat{f}(m)z^m \in H^p(\beta)$ be such that $\hat{f}(0) \neq 0$. If we have

$$\left\{ \frac{\beta(n + j_0)}{\beta(n)} \right\}_n \in \ell^q$$

for some $j_0 \in \mathbb{N}$ and $\beta(n) > 0$ for all n , then f is a cyclic vector of M_z on $H^p(\beta)$.

By a similar method the above results can be extended from the formal power sequence spaces $H^p(\beta)$ to the formal Laurent sequence spaces $L^p(\beta)$.

References

- [1] R. G. Douglas, *Banach algebras techniques in operator theory*, Academic Press, INC., 1972.

- [2] A. L. Shields, Weighted shift operators and analytic functions theory, *Math. Surveys, A.M.S. Providence*, 13, (1974), 49-128.
- [3] B. Yousefi, On the space $\ell^p(\beta)$, *Rendiconti del Circolo Matematico di Palermo*, 49 (2000), 115-120.
- [4] B. Yousefi and K. Jahedi, Application the Rosenthal-Dor theorem on Banach spaces of formal power series, *Islamic Azad University Journal of Sciences*, Fall (2001), 3147-3168.
- [5] B. Yousefi, Unicellularity of the multiplication operator on Banach spaces of formal power series, *Studia Mathematica*, 147 (3)(2001), 201-209.
- [6] B. Yousefi and M. Mehrpoor, On the direct sum of multiplication operators on Banach spaces of analytic functions, *International Journal of Applied Mathematics*, 10 (4) (2002), 353-360.
- [7] B. Yousefi, Bounded analytic structure of the Banach space of formal power series, *Rendiconti del Circolo Matematico di Palermo, Serie II, Tomo LI* (2002), 403-410.
- [8] B. Yousefi and S. Jahedi, Composition operators on Banach spaces of formal power series, *Bollettino Della Unione Matematica Italiano* (8) 6-B (2003), 481-487.
- [9] B. Yousefi, Strictly cyclic algebra of operators acting on Banach spaces $H^p(\beta)$, *Czechoslovak Mathematical Journal*, 54 (129) (2004), 261-266.
- [10] B. Yousefi and R. Soltani, On the Hilbert space of formal power series, *Honam Mathematical Society*, 26 (3) (2004), 299-308.
- [11] B. Yousefi and Y. N. Dehghan, Reflexivity on weighted Hardy spaces, *Southeast Asian Bulletin of Mathematics*, 28 (2004), 587-593.
- [12] B. Yousefi, on the eighteenth question of Allen Shields, *International Journal of Mathematics*, 16 (1) (2005), 37-42.
- [13] B. Yousefi and A. I. Kashkuli, Cyclicity and unicellularity of the differentiation operator on Banach spaces of formal power series, *Mathematical Proceedings of the Royal Irish Academy*, 105A (1) (2005), 1-7.
- [14] B. Yousefi and A. Farrokhinia, Hypercyclicity and supercyclicity on Banach sequence spaces, *Far East Journal of Mathematical Sciences*, 18 (1) (2005), 65-72.
- [15] B. Yousefi and H. Rezaei, Hypercyclicity and supercyclicity on $L^p(\beta)$, *Institute of Mathematics and Computer Sciences*, 19 (1) (2006), 17-20.

- [16] B. Yousefi and A. Farrokhinia, Composition operators on Banach spaces $H^p(\beta)$, *Korean Annals of Mathematics*, 23 (1) (2006), 1-8.

Jinalo Doroodgar

Department of Mathematics
Islamic Azad University-Shiraz Branch
Shiraz, Iran

Shiraz Farzanegan Pre-University
Shiraz Training-Education
Shiraz, Iran
E-mail: jinalollo_dorodgar@yahoo.com

Bahman Yousefi

Department of Mathematics
College of Sciences
Shiraz University
Shiraz 71454, Iran
E-mail: byousefi@shirazu.ac.ir

Archive of SID