

Supercyclicity with Respect to a Sequence on Special Sequence Spaces

S. Foroutan, and B. Yousefi
Shiraz University

J. Doroodgar
Islamic Azad University-Shiraz Branch
Shiraz Farzanegan Pre-University

Abstract. Let $\{\beta(n)\}_{n=-\infty}^{\infty}$ be a sequence of positive numbers such that $\beta(0) = 1$ and let $1 < p < \infty$. We consider the space of all formal Laurent series $f(z) = \sum_{n=-\infty}^{\infty} \hat{f}(n)z^n$ such that

$$\sum_{n=-\infty}^{\infty} |\hat{f}(n)|^p \beta(n)^p < \infty.$$

We investigate the supercyclicity with respect to a sequence on the Banach spaces of formal Laurent series.

AMS Subject Classification: 47B37; 47A16.

Keywords and Phrases: Banach space of Laurent series associated with a sequence β , supercyclic vector, shift operator.

1. Introduction

Let $\{\beta(n)\}_{n=-\infty}^{\infty}$ be a sequence of positive numbers with $\beta(0) = 1$ and $1 < p < \infty$. Consider the space of $f(z) = \sum_{n=-\infty}^{\infty} \hat{f}(n)z^n$ such that

$$\|f\|^p = \|f\|_{\beta}^p = \sum_{n=-\infty}^{\infty} |\hat{f}(n)|^p \beta(n)^p < \infty.$$

They are called formal Laurent series and the space of such formal Laurent series is denoted by $L^p(\beta)$. These are reflexive Banach spaces with the norm $\|\cdot\|_\beta$. The operator B on $L^p(\beta)$ is defined by $Bf_j = f_{j-1}$ for all $j \in \mathbb{Z}$. Clearly B is bounded if and only if the sequence $\{\beta(k)/\beta(k+1)\}_k$ is bounded.

Let X be a complex Banach space and $B(X)$ be the set of bounded linear operators from X into itself. If $T \in B(X)$, then the orbit of a vector $x \in X$ is the set

$$Orb(T, x) = \{T^n x : n \in \mathbb{N} \cup \{0\}\}.$$

A vector $x \in X$ is called hypercyclic for T if $Orb(T, x)$ is dense in X . The operator T is called hypercyclic if it has a hypercyclic vector. A vector $x \in X$ is said to be cyclic for an operator $T \in B(X)$ if the linear span of $Orb(T, x)$ is dense in X . Also a vector $x \in X$ is called a supercyclic vector for an operator $T \in B(X)$ if the set

$$\{\lambda y : y \in Orb(T, x), \lambda \in \mathbb{C}\}$$

is dense in X . An operator $T \in B(X)$ is cyclic (supercyclic) if it has a cyclic (a supercyclic) vector. It is evident that hypercyclicity implies supercyclicity and this, in turn, implies cyclicity.

Sources on formal series include [7, 11, 12, 14, 17]. Also, hypercyclicity and supercyclicity have been studied in several works (see [1, 2, 3, 5, 7, 8, 9, 10, 13, 15, 16, 18, 19, 20]).

We will investigate the supercyclicity with respect to a sequence on the Banach spaces of formal Laurent series.

2. Main Result

Supercyclicity was introduced by Hilden and Wallen ([6]). They showed that all unilateral backward weighted shifts are supercyclic, but there does not exist a vector that is supercyclic vector for all the unilateral backward weighted shifts. H. Salas ([10]) gives a condition for supercyclicity in Frechet spaces.

We can extend the notions to sequences of linear operators; let $\{n_k\}$ be an increasing sequence of nonnegative integers. Then the sequence $\{T_{n_k}\}_{k \geq 0}$ of bounded linear operators from a complex Banach space X into itself is hypercyclic (supercyclic) if there exists $x \in X$ such that the orbit $\{T_{n_k}x\}_{k \geq 0}$ ($\{\lambda T_{n_k}x : k \in \mathbb{N} \cup \{0\}, \lambda \in \mathbb{C}\}$) is dense in X . In the special case when $T \in B(X)$ and the sequence $\{T^{n_k}\}_{k \geq 0}$ is hypercyclic (supercyclic), we say that the operator T is hypercyclic (supercyclic) with respect to the sequence $\{n_k\}$. Here we will investigate the supercyclicity of the operator B with respect to a sequence on the Banach spaces of formal Laurent series.

Suppose that B is bounded on $L^p(\beta)$ and $\{n_k\}$ is an increasing sequence of nonnegative integers. For investigation about the supercyclicity of the sequence $\{B^{n_k}\}_k$, we need the following lemma.

Lemma 1. *Let E be a normed space and T be a bounded linear operator on E . Then the sequence $\{T^{n_k}\}$ is supercyclic if and only if the set*

$$\{(x, \lambda T^{n_k} x) : x \in E, \lambda \in Q + iQ, k \in \mathbb{N}\}$$

is dense in $E \times E$.

Proof. The proof is similar to the proof of Theorem 1.2.2 in [4, page 11] and so we omit it. \square

Theorem 2. *The sequence $\{B^{n_i}\}_i$ is supercyclic on $L^p(\beta)$ if and only if*

$$\liminf_{i \rightarrow \infty} \max \left\{ \frac{\beta(j - n_i)\beta(k + n_i)}{\beta(j)\beta(k)} : |j| \leq n_m, |k| \leq n_m \right\} = 0$$

for all $m \in \mathbb{N}$.

Proof. Let $0 < \varepsilon < 1$ and $m \in \mathbb{N}$. Choose $\alpha > 0$ such that $\frac{\alpha}{1-\alpha} < \varepsilon^{\frac{1}{2}}$.

Let

$$y = w = \sum_{|j| \leq n_m} f_j / \beta(j)$$

be in $L^p(\beta)$. Suppose $\{B_{n_i}\}_i$ is supercyclic. Then by Lemma 1 there exists an arbitrary large $i > m$, a vector

$$x = \sum_n \hat{x}(j) f_j$$

in $L^p(\beta)$, and a complex number λ such that $\|x - w\| < \alpha$ and $\|\lambda B^{n_i} x -$

$y\| < \alpha$. Note that $\lambda \neq 0$. Therefore,

$$\begin{aligned} \|x - w\|^p &= \sum_{|j| \leq n_m} |\hat{x}(j)\beta(j) - 1|^p + \sum_{|j| > n_m} |\hat{x}(j)|^p \beta(j)^p \\ &< \alpha^p. \end{aligned}$$

Thus

$$|\hat{x}(j)|\beta(j) > 1 - \alpha, \quad |j| \leq n_m \tag{1}$$

$$|\hat{x}(j)|\beta(j) < \alpha, \quad |j| > n_m. \tag{2}$$

Also since

$$\begin{aligned} \|\lambda B^{n_i} x - y\|^p &= \sum_{|k| \leq n_m} |\lambda \hat{x}(k + n_i)\beta(k) - 1|^p \\ &+ \sum_{|k| > n_m} |\lambda|^p |\hat{x}(k + n_i)|^p \beta(k)^p < \alpha^p, \end{aligned}$$

we have

$$|\lambda \hat{x}(k + n_i)\beta(k) - 1| < \alpha, \quad |k| \leq n_m \tag{3}$$

$$|\lambda| |\hat{x}(k + n_i)|\beta(k) < \alpha, \quad |k| > n_m. \tag{4}$$

Note that $j - n_i < -n_m$ for $|j| \leq n_m$, so by (1) and (4) we have

$$\frac{\beta(j - n_i)}{\beta(j)} < \frac{1}{|\lambda|} \frac{\alpha}{1 - \alpha}$$

for $|j| \leq n_m$. Also since $k + n_i > n_m$ for $|k| \leq n_m$, by (2) the relation

$$|\hat{x}(k + n_i)| < \frac{\alpha}{\beta(k + n_i)}$$

is consistent and so by (3) we get

$$\frac{\beta(k+n_i)}{\beta(k)} < |\lambda| \frac{\alpha}{1-\alpha}$$

for $|k| \leq n_m$. Therefore

$$\frac{\beta(j-n_i)\beta(k+n_i)}{\beta(j)\beta(k)} < \left(\frac{\alpha}{1-\alpha}\right)^2 < \varepsilon$$

for all $-n_m \leq j, k \leq n_m$ and $i > m$ arbitrarily large enough.

Conversely suppose that $\varepsilon > 0$ is given and consider

$$y = \sum_{|j| \leq n_m} \hat{y}(j) f_j$$

and

$$w = \sum_{|j| \leq n_m} \hat{w}(j) f_j$$

in $L^p(\beta)$ such that both are different from zero. By Lemma 1, it is sufficient to find $x \in L^p(\beta)$ and $i \in \mathbb{N}$ such that $\|x - y\| \leq \varepsilon$ and $\|\lambda B^{n_i} x - w\| \leq \varepsilon$ for some $\lambda \in \mathbb{C}$. Let

$$S^{n_i} w = \sum_{|k| \leq n_m} \hat{w}(k) f_{k+n_i},$$

where $i \in \mathbb{N}$. Also let

$$x = y + \frac{1}{\lambda} S^{n_i} w$$

with i to be determined but $\|\frac{1}{\lambda} S^{n_i} w\| = \varepsilon$. Note that

$$B^{n_i} x = B^{n_i} y + \frac{1}{\lambda} w.$$

Thus it suffices to find i such that $\|\lambda B^{n_i} y\| < \varepsilon$. We have

$$\begin{aligned} \|\lambda B^{n_i} x - w\|^p &= \|\lambda B^{n_i} y\|^p \\ &= \|B^{n_i} y\|^p \|S^{n_i} w\|^p / \varepsilon^p \\ &= \left\| \sum_{|j| \leq n_m} \hat{y}(j) f_{j-n_i} \right\|^p \cdot \left\| \sum_{|k| \leq n_m} \hat{w}(k) f_{k+n_i} \right\|^p / \varepsilon^p \\ &= \left(\sum_{|j| \leq n_m} |\hat{y}(j)|^p \beta(j - n_i)^p \right) \\ &\quad \times \left(\sum_{|k| \leq n_m} |\hat{w}(k)|^p \beta(k + n_i)^p \right) / \varepsilon^p. \end{aligned}$$

So we get

$$\|\lambda B^{n_i} x - w\| \leq \max \left\{ \frac{\beta(j - n_i) \beta(k + n_i)}{\beta(j) \beta(k)} : |k| \leq n_m, |j| \leq n_m \right\} \cdot \|y\| \|w\| / \varepsilon$$

and consequently by our hypothesis there exists i large enough such that

$\|\lambda B^{n_i} x - w\| < \varepsilon$. This completes the proof. \square

Acknowledgment

The third author, thanks the Research Council of Islamic Azad University-Shiraz Branch.

References

- [1] N. S. Feldman, T. L. Miller, and V. G. Miller, Hypercyclic and supercyclic cohyponormal operators, *Acta. Sci. Math.*, (Szeged) 68 (2002), 303-328.
- [2] N. S. Feldman, Hypercyclicity and supercyclicity for invertible bilateral weighted shifts, *Proc. Amer. Math. Soc.*, 131 (2003), 479-485.
- [3] G. Godefroy and J. H. Shapiro, Operators with dense, invariant cyclic vector manifolds, *J. Func. Anal.*, 98 (1991), 229-269.
- [4] K. G. Grosse-Erdmann, Holomorphic Monster und universelle Funktionen, *Mitt. Math. Sem. Giessen*, 176 (1987), 1-84.
- [5] G. Herzog, On linear having supercyclic operators, *Studia Math.*, 103 (1992), 295-298.
- [6] H. M. Hilden and L. J. Wallen, Some cyclic and non-cyclic vectors of certain operators, *Indiana Univ. Math. Journal*, 24 (1974), 557-565.
- [7] K. Jahedi and B. Yousefi, On the weighted Hardy spaces, *Journal of Mathematical Extension*, 1 (1) (2006), 45-55.
- [8] C. Kitai, *Invariant closed sets for linear operators*, Dissertation, University of Toronto, 1982.
- [9] A. Montes-Rodriguez and H. N. Salas, Supercyclic subspaces, spectral theory and weighted shifts, *Adv. Math.*, 163 (2001), 74-134.
- [10] H. N. Salas, Supercyclicity and weighted shifts, *Studia Math.*, 135 (1999), 55-74.
- [11] A. L. Shields, Weighted shift operators and analytic functions theory, *Math. Surveys*, Amer. Math. Soc. Providence, 13(1974), 49-128.
- [12] B. Yousefi, On the space $\ell^p(\beta)$, *Rend. Circ. Mat. Palermo*, 49 (2000), 115-120.
- [13] B. Yousefi and H. Rezaei, Hypercyclicity on the algebra of Hilbert-Schmidt operators, *Journal of Results in Mathematics*, 46 (2004), 174-180.
- [14] B. Yousefi, On the eighteenth question of Allen Shields, *International Journal of Mathematics*, 16, No. 1 (2005) 37-42.
- [15] B. Yousefi and H. Rezaei, Some necessary and sufficient conditions for Hypercyclicity Criterion, *Proc. Indian Acad. Sci. (Math. Sci.)*, 115, (2) (2005), 209-216.

- [16] B. Yousefi and A. Farrokhinia, Hypercyclicity and supercyclicity on Banach sequence spaces, *Far East Journal of Mathematics*, 18 (1) (2005), 65-72.
- [17] B. Yousefi, Z. Kamali, and L. Bagheri, Isometries of weighted Hardy spaces among composition operators, *Korean Annals of Math.*, 23 (2) (2006), 119-125.
- [18] B. Yousefi and A. Farrokhinia, On the hereditarily hypercyclic vectors, *Journal of the Korean Mathematical Society*, to appear.
- [19] B. Yousefi and H. Rezaei, Hypercyclic property of weighted composition operators, *Proc. Amer. Math. Soc.*, to appear.
- [20] B. Yousefi, H. Rezaei, and J. Doroodgar, Supercyclicity in the operator algebra using Hilbert-Schmidt operators, *Rend. Circ. Mat. Palermo*, to appear.

Sedigheh Foroutan

Department of Mathematics
 College of Sciences
 Shiraz University
 Shiraz 71454, Iran
 E-mail: foroutan@shirazu.ac.ir

Bahmann Yousefi

Department of Mathematics
 College of Sciences
 Shiraz University
 Shiraz 71454, Iran
 E-mail: byousefi@shirazu.ac.ir

Jinalo Doroodgar

Department of Mathematics
 Islamic Azad University-Shiraz Branch
 Shiraz, Iran
 Farzanegan Pre-University, Fars Educational Organization
 Shiraz, Iran
 E-mail: jinalollo-dorodgar@yahoo.com