

# Exact Solutions for the Modified KdV and the Generalized KdV Equations via Exp-Function Method

J. Manafian Heris\* \*

Islamic Azad University-Ahar Branch

M. Bagheri

Islamic Azad University-Ahar Branch

**Abstract.** An application of the Exp-function method (EFM) to search for exact solutions of nonlinear partial differential equations is analyzed. This method is used for the modified KdV equation and the generalized KdV equation. The EFM was used to construct periodic wave and solitary wave solutions of nonlinear evolution equations (NLEEs). This method is developed for searching exact travelling wave solutions of nonlinear partial differential equations. It is shown that the Exp-function method, with the help of symbolic computation, provides a straightforward and powerful mathematical tool for solving nonlinear evolution equations in mathematical physics and applied mathematics.

**AMS Subject Classification:** 2015.

**Keywords and Phrases:** Exp-function method, modified KdV equation, generalized KdV equation, periodic wave and solitary wave solutions, nonlinear partial differential equations (NLPDEs).

## 1. Introduction

Nonlinear phenomena plays a fundamental role in applied mathematics and physics. Recently, the study of nonlinear partial differential equations in modelling physical phenomena, has become an important tool.

---

Received January 2010; Final Revised March 2010

\*\*Corresponding author

## *Archive of SID*

The investigation of the traveling wave solutions plays an important role in nonlinear sciences. A variety of powerful methods have been presented, such as the inverse scattering transform ([3]), Hirota's bilinear method ([17, 23]), homotopy analysis method ([1, 7]), variational iteration method ([8, 9]), Adomian decomposition method ([10]), homotopy perturbation method ([11, 12, 19, 21, 22]), sine-cosine method ([24, 32]), tanh-function method ([4, 13, 25]), tanh-coth method ([26, 27]), Bäcklund transformation ([18, 20]) and so on. Here, we use an effective method for constructing a range of exact solutions for the following nonlinear partial differential equations which was first presented by J. H. He ([14]). A new method called the Exp-function method (EFM) is presented to look for traveling wave solutions of nonlinear evolution equations (NLEEs). The Exp-function method has successfully been applied to many situations. For example, He et al. ([15]) have solved the nonlinear wave equations by using the Exp-function method. Wu and He ([30]) have used the Exp-function method to give new periodic solutions for nonlinear evolution equations. He and Abdou ([16]) have examined the Exp-function method to find generalized solitary solutions and compacton-like solutions of the Jaulent-Miodek equations. Abdou ([2]) has solved generalized solitary and periodic solutions for nonlinear partial differential equations by the Exp-function method. Boz and Bekir ([5]) have applied the Exp-function method for (3+1)-dimensional nonlinear evolution equations. Chun ([6]) has obtained the solitons and periodic solutions for the fifth-order KdV equation by using the Exp-function method. The Exp-function method along with Hirota's and tanh-coth methods have been applied for computing solitary wave solutions of the generalized shallow water wave equation by Wazwaz ([28]). Wu and He ([31]) have applied the Exp-function method to nonlinear equations. The EFM has recently been generalized by Zhang ([33]) to high-dimensional nonlinear evolution equation. The KdV equation, that plays an important role in the solitary wave theory, is given by

$$u_t + 6uu_x + u_{xxx} = 0, \quad (1)$$

The balance between the weak nonlinear steepening of  $uu_x$  and the dispersion effect of  $u_{xxx}$  in Eq. (1) generates solitons. The KdV equation

## Archive of SID

is a spatially one dimensional model. The KdV equation is completely integrable ([29]), admits multiple-soliton solutions and exhibits an infinite number of conservation laws of energy. In this article, we have used the Exp-function method to investigate the modified KdV equation and the generalized KdV equation given by [25]

$$u_t + 3au^2u_x + bu_{xxx} = 0, \quad (2)$$

and

$$u_t + a(n+1)u^n u_x + bu_{xxx} = 0. \quad (3)$$

By using the Exp-function method we obtained various solutions for the modified KdV equation and the generalized KdV equation and some new results are formally developed in this article. Our aim of this paper is to obtain analytical solutions of the modified KdV equation and the generalized KdV equation, and to determine the accuracy of the Exp-function method in solving these kinds of problems. The remainder of the paper is organized as follows: In Section 2, a brief discussions for the Exp-function method is presented and exact solutions of Eqs. (2) and (3) are obtained. In Section 3, we describe this method briefly and apply this technique to the modified KdV equation. In Section 4, exact results are considered for the generalized KdV equation by the aforementioned method. Section 5, ends this report with a brief conclusion.

## 2. Basic Idea of the Exp-Function Method

We first consider the nonlinear equation of the form:

$$\mathcal{N}(u, u_t, u_x, u_{xx}, u_{tt}, u_{tx}, \dots) = 0, \quad (4)$$

and introduce a transformation

$$u(x, t) = u(\eta), \quad \eta = kx + \omega t, \quad (5)$$

where  $k$  and  $\omega$  are constant to be determined later. Therefore Eq. (4) is reduced to an ODE as follows

$$\mathcal{M}(u, \omega u', ku', k^2 u'', \dots) = 0. \quad (6)$$

## Archive of SID

The EFM is based on the assumption that travelling wave solutions as in ([15]) can be expressed in the form

$$u(\eta) = \frac{\sum_{n=-c}^d a_n \exp(n\eta)}{\sum_{m=-p}^q b_m \exp(m\eta)}, \quad (7)$$

where  $c$ ,  $d$ ,  $p$  and  $q$  are positive integers which could be freely chosen and  $a_n$ 's and  $b_m$ 's are unknown constants to be determined. To determine the values of  $c$  and  $p$ , we balance the linear term of highest order in Eq. (6) with the highest order nonlinear term. Also to determine the values of  $d$  and  $q$ , we balance the linear term of lowest order in Eq. (6) with the lowest order nonlinear term.

### 3. The Modified KdV Equation

In this section we employ the Exp-function method to the modified KdV equation

$$u_t + 3au^2u_x + bu_{xxx} = 0, \quad (8)$$

and by using the wave variable  $\eta = \mu(x - ct)$  reduces it to an ODE [25],

$$-c\mu u' + 3a\mu u^2 u' + b\mu^3 u''' = 0. \quad (9)$$

Then by integrating Eq. (9) and neglecting the constant of integration we obtain

$$-cu + au^3 + b\mu^2 u'' = 0. \quad (10)$$

In order to determine the values of  $c$  and  $p$ , we balance the linear term of the highest order  $u''$  with the highest order nonlinear term  $u^3$  in Eq. (10) and get

$$u'' = \frac{c_1 \exp((c + 3p)\eta) + \dots}{c_2 \exp(4p\eta) + \dots} \quad (11)$$

$$u^3 = \frac{c_3 \exp(3c\eta) + \dots}{c_4 \exp(3p\eta) + \dots} = \frac{c_3 \exp((3c + p)\eta) + \dots}{c_4 \exp(4p\eta) + \dots} \quad (12)$$

*Archive of SID*

respectively. Balancing highest order of the Exp-function in (11) and (12) and get

$$c + 3p = 3c + p, \quad (13)$$

which leads to the result  $c = p$ .

Similarly, to determine the values of  $d$  and  $q$ , for the terms  $u''$  and  $u^3$  in Eq. (10) by simple calculation, we obtain

$$u'' = \frac{\dots + d_1 \exp(-(d + 3q)\eta)}{\dots + d_2 \exp(-4q\eta)}, \quad (14)$$

$$u^3 = \frac{\dots + d_3 \exp(-3d\eta)}{\dots + d_4 \exp(-3q\eta)} = \frac{\dots + d_3 \exp(-(3d + q)\eta)}{\dots + d_4 \exp(-4q\eta)}, \quad (15)$$

respectively. Balancing lowest order of the Exp-function in (14) and (15) gives us

$$-(d + 3q) = -(3d + q), \quad (16)$$

which leads to the result  $d = q$ .

Now we consider the following cases:

**Case I:**  $p = c = 1$  and  $q = d = 1$ .

For simplicity, we set  $b_1 = 1$ ,  $p = c = 1$  and  $d = q = 1$ . Then (7) reduces to

$$u(\eta) = \frac{a_1 \exp(\eta) + a_0 + a_{-1} \exp(-\eta)}{b_1 \exp(\eta) + b_0 + b_{-1} \exp(-\eta)}. \quad (17)$$

Substituting (17) into Eq. (10), and using the well-known Maple software, we have

$$\frac{1}{A} [C_3 \exp(3\eta) + C_2 \exp(2\eta) + C_1 \exp(\eta) + C_0 + C_{-1} \exp(-\eta) + \quad (18)$$

$$C_{-2} \exp(-2\eta) + C_{-3} \exp(-3\eta)] = 0,$$

where

$$A = [b_{-1} \exp(-\eta) + b_0 + \exp(\eta)]^3, \quad (19)$$

## Archive of SID

and  $C_n$ ' are coefficients of  $\exp(n\eta)$ '. Equating the coefficients of  $\exp(n\eta)$  to zero, we obtain the following set of algebraic equations for  $a_1, a_0, a_{-1}, b_0, b_{-1}$  and  $c$ , as

$$\begin{cases} C_3 = 0, & C_2 = 0, & C_1 = 0, \\ C_0 = 0, \\ C_{-3} = 0, & C_{-2} = 0, & C_{-1} = 0. \end{cases} \quad (20)$$

Solving the system of algebraic equations with the help of Maple gives the following set of non-trivial solutions

(I) The first set is:

$$a_{-1} = 0, \quad b_0 = 0, \quad b_{-1} = \frac{1}{8} \frac{aa_0^2}{\mu^2 b}, \quad a_0 = a_0, \quad a_1 = 0, \quad c = \mu^2 b, \quad \mu = \mu, \quad (21)$$

which gives:

$$u_1(x, t) = \frac{a_0}{\frac{1}{8} \frac{aa_0^2}{\mu^2 b} \exp[-\mu(x - \mu^2 bt)] + \exp[(\mu(x - \mu^2 bt))]} \quad (22)$$

If we choose  $a_0 = 2\sqrt{\frac{2\mu^2 b}{a}}$  or  $a_0 = 2\sqrt{-\frac{2\mu^2 b}{a}}$ , then the Solution (22) becomes to

$$\begin{aligned} u_{1,1}(x, t) &= \sqrt{\frac{2\mu^2 b}{a}} \operatorname{sech}[\mu(x - \mu^2 bt)], \\ u_{1,2}(x, t) &= \sqrt{-\frac{2\mu^2 b}{a}} \operatorname{csch}[\mu(x - \mu^2 bt)]. \end{aligned}$$

(II) The second set is:

$$a_{-1} = -a_1 b_{-1}, \quad b_0 = 0, \quad b_{-1} = b-1, \quad a_0 = 0, \quad a_1 = a_1, \quad c = aa_1^2, \quad \mu = \sqrt{-\frac{a}{2b}} a_1, \quad (23)$$

which gives:

$$u_2(x, t) = \frac{-a_1 b_{-1} \exp\left[-\sqrt{-\frac{a}{2b}} a_1 (x - aa_1^2 t)\right] + a_1 \exp\left[\sqrt{-\frac{a}{2b}} a_1 (x - aa_1^2 t)\right]}{b_{-1} \exp\left[-\sqrt{-\frac{a}{2b}} a_1 (x - aa_1^2 t)\right] + \exp\left[\sqrt{-\frac{a}{2b}} a_1 (x - aa_1^2 t)\right]} \quad (24)$$

If we choose  $b_{-1} = 1$  or  $b_{-1} = -1$ , then the Solution (24) becomes to

$$\begin{aligned} u_{2,1}(x, t) &= a_1 \tanh\left[\sqrt{-\frac{a}{2b}} a_1 (x - aa_1^2 t)\right], \\ u_{2,2}(x, t) &= a_1 \coth\left[\sqrt{-\frac{a}{2b}} a_1 (x - aa_1^2 t)\right]. \end{aligned}$$

### Archive of SID

Respectively, where  $a_1$  is an arbitrary constant. In the case  $\mu$  is an imaginary number, each of the obtained solitary solutions above can be converted into a periodic solution or compact-like solution. Here, we only discuss the solution given by (24). If  $\mu = i\bar{\mu}$  in (24), then we obtain

$$\exp[i\bar{\mu}(x - 2b\bar{\mu}^2t)] = \cos[\bar{\mu}(x - 2b\bar{\mu}^2t)] + i \sin[\bar{\mu}(x - 2b\bar{\mu}^2t)], \quad (25)$$

and

$$\exp[-i\bar{\mu}(x - 2b\bar{\mu}^2t)] = \cos[\bar{\mu}(x - 2b\bar{\mu}^2t)] - i \sin[\bar{\mu}(x - 2b\bar{\mu}^2t)]. \quad (26)$$

and (24) becomes:

$$u_2(x, t) = a_1 \frac{(1 - b_{-1})\cos[\bar{\mu}(x - 2b\bar{\mu}^2t)] + i(1 + b_{-1})\sin[\bar{\mu}(x - 2b\bar{\mu}^2t)]}{(1 + b_{-1})\cos[\bar{\mu}(x - 2b\bar{\mu}^2t)] + i(1 - b_{-1})\sin[\bar{\mu}(x - 2b\bar{\mu}^2t)]}. \quad (27)$$

or

$$u_2(x, t) = a_1 \frac{(1 - b_{-1}) + i(1 + b_{-1})\tan[\bar{\mu}(x - 2b\bar{\mu}^2t)]}{(1 + b_{-1}) + i(1 - b_{-1})\tan[\bar{\mu}(x - 2b\bar{\mu}^2t)]}. \quad (28)$$

If we choose  $b_{-1} = 1$  or  $b_{-1} = -1$ , then the complex Solution (28) respectively gives:

$$u_{2,3}(x, t) = a_1 i \tan[\bar{\mu}(x - 2b\bar{\mu}^2t)], \quad u_{2,4}(x, t) = a_1 i \cot[\bar{\mu}(x - 2b\bar{\mu}^2t)],$$

where  $a_1$  is an arbitrary constant.

**Case II:**  $p = c = 2$  and  $q = d = 1$ .

For simplicity, we set  $b_2 = 1$ ,  $p = c = 2$  and  $d = q = 1$ . Then (7) reduces to

$$u(\eta) = \frac{a_2 \exp(2\eta) + a_1 \exp(\eta) + a_0 + a_{-1} \exp(-\eta)}{\exp(2\eta) + b_1 \exp(\eta) + b_0 + b_{-1} \exp(-\eta)}. \quad (29)$$

Substituting (29) into Eq. (10), and using the well-known Maple software, we get

$$\frac{1}{A} [C_6 \exp(6\eta) + C_5 \exp(5\eta) + C_4 \exp(4\eta) + C_3 \exp(3\eta) + C_2 \exp(2\eta)] \quad (30)$$

*Archive of SID*

$$C_1 \exp(\eta) + C_0 + C_{-1} \exp(-\eta) + C_{-2} \exp(-2\eta) + C_{-3} \exp(-3\eta)] = 0,$$

where

$$A = [b_{-1} \exp(-\eta) + b_0 + b_1 \exp(\eta) + \exp(2\eta)]^3, \quad (31)$$

and  $C_n$ ' are coefficients of  $\exp(n\eta)$ '. Equating the coefficients of  $\exp(n\eta)$  to zero, we obtain the following set of algebraic equations for  $a_1, a_0, a_{-1}, b_1, b_0, b_{-1}$  and  $c$ , as

$$\begin{cases} C_6 = 0, & C_5 = 0, & C_4 = 0, & C_3 = 0, & C_2 = 0, & C_1 = 0, \\ C_0 = 0, \\ C_{-3} = 0, & C_{-2} = 0, & C_{-1} = 0. \end{cases} \quad (32)$$

Solving the system of algebraic equations gives the following sets of non-trivial solutions

(I) The first set is:

$$a_{-1} = -a_2 b_{-1}, \quad b_{-1} = b_{-1}, \quad a_0 = 0, \quad b_0 = 0, \quad a_2 = a_2, \quad c = a a_2^2, \quad (33)$$

$$a_1 = 0, \quad b_1 = 0, \quad \mu = \sqrt{-\frac{2a}{9b}} a_2,$$

which gives:

$$u_1(x, t) = \frac{-a_2 b_{-1} \exp \left[ -\sqrt{-\frac{2a}{9b}} a_2 (x - a a_2^2 t) \right] + a_2 \exp \left[ 2\sqrt{-\frac{2a}{9b}} a_2 (x - a a_2^2 t) \right]}{b_{-1} \exp \left[ -\sqrt{-\frac{2a}{9b}} a_2 (x - a a_2^2 t) \right] + \exp \left[ 2\sqrt{-\frac{2a}{9b}} a_2 (x - a a_2^2 t) \right]}. \quad (34)$$

If we choose  $b_{-1} = 1$  or  $b_{-1} = -1$ , then the solution (34) respectively gives

$$\begin{aligned} u_{1,1}(x, t) &= a_2 \tanh \left[ \sqrt{-\frac{2a}{9b}} a_2 (x - a a_2^2 t) \right], \\ u_{1,2}(x, t) &= a_2 \coth \left[ \sqrt{-\frac{2a}{9b}} a_2 (x - a a_2^2 t) \right]. \end{aligned}$$

where  $a_2$  is an arbitrary constant.

(II) The second set is:

$$a_{-1} = \frac{1}{8} \frac{(a_2^2 b_1^2 - 4a_2^2 b_0 - a_1^2)(a_2 b_1 - a_1)}{a_2^2}, \quad a_0 = -\frac{1}{2} \frac{a_2^2 b_1^2 - 2a_2^2 b_0 - a_1^2}{a_2}, \quad (35)$$



*Archive of SID*

$$b_{-1} = -\frac{1}{8} \frac{(a_2^2 b_1^2 - 4a_2^2 b_0 - a_1^2)(a_2 b_1 - a_1)}{a_2^3}, \quad b_0 = b_0, \quad a_2 = a_2, \quad c = aa_2^2,$$

$$a_1 = a_1, \quad b_1 = b_1, \quad \mu = \sqrt{-\frac{a}{2b}} a_2, \quad \eta = \sqrt{-\frac{a}{2b}} a_2 (x - aa_2^2 t),$$

which gives:

$$u_2(x, t) = \frac{\frac{1}{8} \frac{(a_2^2 b_1^2 - 4a_2^2 b_0 - a_1^2)(a_2 b_1 - a_1)}{a_2^3} e^{(-\eta)} - \frac{1}{2} \frac{a_2^2 b_1^2 - 2a_2^2 b_0 - a_1^2}{a_2} + a_1 e^{(\eta)} + a_2 e^{(2\eta)}}{-\frac{1}{8} \frac{(a_2^2 b_1^2 - 4a_2^2 b_0 - a_1^2)(a_2 b_1 - a_1)}{a_2^3} e^{(-\eta)} + b_0 + b_1 e^{(\eta)} + e^{(2\eta)}}.$$

**Case III:**  $p = c = 2$  and  $q = d = 2$ .

Since the values of  $c$  and  $d$  can be freely chosen, we set  $p = c = 2$  and  $d = q = 2$  and then the trial function (7), becomes

$$u(\eta) = \frac{a_2 \exp(2\eta) + a_1 \exp(\eta) + a_0 + a_{-1} \exp(-\eta) + a_{-2} \exp(-2\eta)}{b_2 \exp(2\eta) + b_1 \exp(\eta) + b_0 + b_{-1} \exp(-\eta) + b_{-2} \exp(-2\eta)}. \quad (36)$$

There are some free parameters in (36). We set  $b_2 = 1$  and  $b_1 = b_{-1} = 0$  for simplicity, the trial function, Eq. (36) is simplified as follows

$$u(\eta) = \frac{a_2 \exp(2\eta) + a_1 \exp(\eta) + a_0 + a_{-1} \exp(-\eta) + a_{-2} \exp(-2\eta)}{\exp(2\eta) + b_0 + b_{-2} \exp(-2\eta)}. \quad (37)$$

Substituting (37), into Eq. (10), and using the well-known Maple software, we will have

$$\begin{aligned} & \frac{1}{A} [C_6 \exp(6\eta) + C_5 \exp(5\eta) + C_4 \exp(4\eta) + C_3 \exp(3\eta) + C_2 \exp(2\eta) \\ & + C_1 \exp(\eta) + C_0 + C_{-1} \exp(-\eta) + C_{-2} \exp(-2\eta) + C_{-3} \exp(-3\eta) + \\ & C_{-4} \exp(-4\eta) + C_{-5} \exp(-5\eta) + C_{-6} \exp(-6\eta)] = 0, \end{aligned} \quad (38)$$

where

$$A = [b_{-1} \exp(-\eta) + b_0 + b_1 \exp(\eta) + \exp(2\eta)]^3, \quad (39)$$

and  $C_n$ ' are coefficients of  $\exp(n\eta)$ '. Equating the coefficients of  $\exp(n\eta)$  to zero, we obtain the following set of algebraic equations for  $a_2, a_{-2}, a_1, a_0, a_{-1}, b_0, b_{-2}$  and  $c$ , as

$$\begin{cases} C_6 = 0, & C_5 = 0, & C_4 = 0, & C_3 = 0, & C_2 = 0, & C_1 = 0. \\ C_0 = 0, \\ C_{-6} = 0, & C_{-5} = 0, & C_{-4} = 0, & C_{-3} = 0, & C_{-2} = 0, & C_{-1} = 0. \end{cases} \quad (40)$$

# Archive of SID

Solving the system of algebraic equations we get

(I) The first set is:

$$a_1 = a_1, \quad a_{-2} = 0, \quad a_{-1} = -\frac{1}{8} \frac{a_1(aa_1^2 - 8b\mu^2b_0)}{b\mu^2}, \quad b_0 = b_0, \quad a_0 = 0, \quad (41)$$

$$c = b\mu^2, \quad b_{-2} = -\frac{1}{64} \frac{aa_1^2(aa_1^2 - 8b\mu^2b_0)}{b^2\mu^4}, \quad a_2 = 0, \quad \mu = \mu,$$

which gives:

$$u_1(x, t) = \frac{a_{-1}e^{-A} + a_1e^A}{b_{-2}e^{-2A} + b_0 + e^{2A}},$$

where  $A = \mu(x - b\mu^2t)$ ,  $a_0$ ,  $b_0$ ,  $a_1$  and  $a_2$  are arbitrary constants.

(II) The second set is:

$$a_1 = 0, \quad a_{-2} = 0, \quad a_{-1} = 0, \quad b_0 = 0, \quad a_0 = a_0, \quad a_2 = 0, \quad (42)$$

$$c = 4b\mu^2, \quad b_{-2} = \frac{1}{32} \frac{aa_0^2}{b\mu^2}, \quad \mu = \mu,$$

which gives:

$$u_2(x, t) = \frac{a_0}{\frac{1}{32} \frac{aa_0^2}{b\mu^2} \exp[-2\mu(x - 4b\mu^2t)] + \exp[2\mu(x - 4b\mu^2t)]}. \quad (43)$$

If we choose  $a_0 = 4\sqrt{\frac{2\mu^2b}{a}}$  or  $a_0 = 4\sqrt{\frac{-2\mu^2b}{a}}$ , then the solution (43) respectively gives

$$u_{2,1}(x, t) = \sqrt{\frac{8b\mu^2}{a}} \operatorname{sech} [2\mu(x - 4b\mu^2t)],$$

$$u_{2,2}(x, t) = \sqrt{\frac{-8b\mu^2}{a}} \operatorname{csch} [2\mu(x - 4b\mu^2t)].$$

Obtained results in above are the exact solutions of the modified KdV equation.

## 4. The Generalized KdV Equation

We next apply the Exp-function method to the generalized KdV equation

$$u_t + a(n+1)u^n u_x + bu_{xxx} = 0, \quad (44)$$

and by using the wave variable  $\eta = \mu(x - ct)$  reduce it to an ODE [25].

$$-c\mu u' + a(n+1)\mu u^n u' + b\mu^3 u''' = 0, \quad (45)$$

where by integrating (45) and neglecting the constant of integration we obtain

$$-cu + au^{n+1} + b\mu^2 u'' = 0. \quad (46)$$

For various values of  $n$  we introduce the transformation

$$u = v^{\frac{1}{n}}, \quad (47)$$

that carries Eq. (46) to

$$-cn^2 v^2 + an^2 v^3 + bn\mu^2 vv'' + b(1-n)\mu^2 (v')^2 = 0. \quad (48)$$

In order to determine the values of  $c$  and  $p$ , we balance  $v^3$  with  $vv''$  in Eq. (48), to get

$$vv'' = \frac{c_1 \exp((2c+3p)\eta) + \dots}{c_2 \exp(5p\eta) + \dots}, \quad (49)$$

$$v^3 = \frac{c_3 \exp(3c\eta) + \dots}{c_4 \exp(3p\eta) + \dots} = \frac{c_3 \exp((3c+2p)\eta) + \dots}{c_4 \exp(5p\eta) + \dots}, \quad (50)$$

respectively. Balancing highest order of the Exp-function in (49) and (50), we obtain

$$c = p. \quad (51)$$

Similarly, to determine the values of  $d$  and  $q$ , we balance  $v^3$  with  $vv''$  in Eq. (48), we obtain

$$vv'' = \frac{\dots + d_1 \exp(-(2d+3q)\eta)}{\dots + d_2 \exp(-5q\eta)}, \quad (52)$$

*Archive of SID*

$$v^3 = \frac{\dots + d_3 \exp(-3d\eta)}{\dots + d_4 \exp(-3q\eta)} = \frac{\dots + d_3 \exp(-(3d + 2q)\eta)}{\dots + d_4 \exp(-5q\eta)}, \quad (53)$$

respectively. Balancing lowest order of the Exp-function in (52) and (53), we obtain

$$d = q. \quad (54)$$

**Case I:**  $p = c = 1$  and  $q = d = 1$ .

For simplicity, we set  $b_1 = 1$ ,  $p = c = 1$  and  $d = q = 1$ . Then (7) reduces to

$$v(\eta) = \frac{a_1 \exp(\eta) + a_0 + a_{-1} \exp(-\eta)}{b_1 \exp(\eta) + b_0 + b_{-1} \exp(-\eta)}. \quad (55)$$

Substituting (55) into Eq. (48) and using the well-known Maple software, we will have

$$\frac{1}{A} [C_4 \exp(4\eta) + C_3 \exp(3\eta) + C_2 \exp(2\eta) + C_1 \exp(\eta) + C_0 + C_{-1} \exp(-\eta) + C_{-2} \exp(-2\eta) + C_{-3} \exp(-3\eta) + C_{-4} \exp(-4\eta)] = 0 \quad (56)$$

where

$$A = [b_{-1} \exp(-\eta) + b_0 + \exp(\eta)]^4, \quad (57)$$

and  $C_n$ ' are coefficients of  $\exp(n\eta)$ '. Equating the coefficients of  $\exp(n\eta)$  to zero, we obtain the following set of algebraic equations for  $a_1, a_0, a_{-1}, b_0, b_{-1}$  and  $c$ , as

$$\begin{cases} C_4 = 0, & C_3 = 0, & C_2 = 0, & C_1 = 0, \\ C_0 = 0, \\ C_{-4} = 0, & C_{-3} = 0, & C_{-2} = 0, & C_{-1} = 0. \end{cases} \quad (58)$$

Solving the system of algebraic equations we get

(I) The first set is:

$$a_{-1} = 0, \quad b_0 = b_0, \quad b_{-1} = \frac{1}{4}b_0^2, \quad a_0 = \frac{b\mu^2 b_0(n+2)}{an^2}, \quad a_1 = 0, \quad c = \frac{\mu^2 b}{n^2}, \quad \mu = \mu, \quad (59)$$

which gives:

$$v_1(x, t) = \frac{\frac{b\mu^2 b_0(n+2)}{an^2}}{\frac{1}{4}b_0^2 \exp[-\mu(x - \frac{\mu^2 b}{n^2}t)] + b_0 + \exp[(\mu(x - \frac{\mu^2 b}{n^2}t)]},$$

*Archive of SID*

$$u_1(x, t) = \left\{ \frac{\frac{b\mu^2 b_0(n+2)}{an^2}}{\frac{1}{4}b_0^2 \exp[-\mu(x - \frac{\mu^2 b}{n^2}t)] + b_0 + \exp[(\mu(x - \frac{\mu^2 b}{n^2}t)]} \right\}^{\frac{1}{n}} \quad (60)$$

If we choose  $b_0 = 2$  or  $b_0 = -2$ , then the solution (60) respectively gives (cf. Eqs. (38) and (37) in [25])

$$u_{1,1}(x, t) = \left\{ \frac{b\mu^2(n+2)}{2an^2} \operatorname{sech}^2 \left[ \frac{\mu}{2} \left( x - \frac{\mu^2 b}{n^2}t \right) \right] \right\}^{\frac{1}{n}},$$

$$u_{1,2}(x, t) = \left\{ -\frac{b\mu^2(n+2)}{2an^2} \operatorname{csch}^2 \left[ \frac{\mu}{2} \left( x - \frac{\mu^2 b}{n^2}t \right) \right] \right\}^{\frac{1}{n}}.$$

If  $\mu = i\bar{\mu}$  in (60) then we obtain

$$\exp \left[ i\bar{\mu} \left( x + \frac{\bar{\mu}^2 b}{n^2}t \right) \right] = \cos \left[ \bar{\mu} \left( x + \frac{\bar{\mu}^2 b}{n^2}t \right) \right] + i \sin \left[ \bar{\mu} \left( x + \frac{\bar{\mu}^2 b}{n^2}t \right) \right], \quad (61)$$

and

$$\exp \left[ -i\bar{\mu} \left( x + \frac{\bar{\mu}^2 b}{n^2}t \right) \right] = \cos \left[ \bar{\mu} \left( x + \frac{\bar{\mu}^2 b}{n^2}t \right) \right] - i \sin \left[ \bar{\mu} \left( x + \frac{\bar{\mu}^2 b}{n^2}t \right) \right]. \quad (62)$$

and (60) becomes

$$u_1(x, t) = \left\{ -\frac{\frac{4b\bar{\mu}^2 b_0(n+2)}{an^2}}{(4 + b_0^2)\cos \left[ \bar{\mu} \left( x + \frac{\bar{\mu}^2 b}{n^2}t \right) \right] + 4b_0 + i(4 - b_0^2)\sin \left[ \bar{\mu} \left( x + \frac{\bar{\mu}^2 b}{n^2}t \right) \right]} \right\}^{\frac{1}{n}}. \quad (63)$$

It should be pointed out that the transformation  $\mu = i\bar{\mu}$  was first used by He and Wu in [16] to find periodic solutions or compact-like solutions from the obtained solitary solutions. If we search for a periodic solution or compact-like solution, the imaginary part in the denominator of (63) must be zero, that requires that

$$b_0 = \pm 2. \quad (64)$$

by applying (64) then (63) reduces to a compact-like solution, which reads

$$u_1(x, t) = \left\{ \mp \frac{\frac{b\bar{\mu}^2(n+2)}{an^2}}{\cos \left[ \bar{\mu} \left( x + \frac{\bar{\mu}^2 b}{n^2}t \right) \right] \pm 1} \right\}^{\frac{1}{n}}. \quad (65)$$

*Archive of SID*

Also (65) is further simplified to obtain a periodic solution respectively (cf. Eqs. (40) and (39) in [25])

$$u_{1,3}(x, t) = \left\{ -\frac{b\bar{\mu}^2(n+2)}{an^2} \sec^2 \left[ \frac{\bar{\mu}}{2} \left( x + \frac{\bar{\mu}^2 b}{n^2} t \right) \right] \right\}^{\frac{1}{n}}, \quad (66)$$

$$u_{1,4}(x, t) = \left\{ -\frac{b\bar{\mu}^2(n+2)}{an^2} \csc^2 \left[ \frac{\bar{\mu}}{2} \left( x + \frac{\bar{\mu}^2 b}{n^2} t \right) \right] \right\}^{\frac{1}{n}}. \quad (67)$$

**Case II:**  $p = c = 2$  and  $q = d = 1$ .

For simplicity, we set  $b_2 = 1$ ,  $p = c = 2$  and  $d = q = 1$ . Then (7) reduces to

$$v(\eta) = \frac{a_2 \exp(2\eta) + a_1 \exp(\eta) + a_0 + a_{-1} \exp(-\eta)}{\exp(2\eta) + b_1 \exp(\eta) + b_0 + b_{-1} \exp(-\eta)}. \quad (68)$$

Substituting (68) into Eq. (48), we have

$$\begin{aligned} \frac{1}{A} [C_8 \exp(8\eta) + C_7 \exp(7\eta) + C_6 \exp(6\eta) + C_5 \exp(5\eta) + C_4 \exp(4\eta) + \\ C_3 \exp(3\eta) + C_2 \exp(2\eta) + C_1 \exp(\eta) + C_0 + C_{-1} \exp(-\eta) + \\ C_{-2} \exp(-2\eta) + C_{-3} \exp(-3\eta) + C_{-4} \exp(-4\eta)] = 0, \end{aligned} \quad (69)$$

where

$$A = [b_{-1} \exp(-\eta) + b_0 + b_1 \exp(\eta) + \exp(2\eta)]^4, \quad (70)$$

and  $C_n$ ' are coefficients of  $\exp(n\eta)$ '. Equating the coefficients of  $\exp(n\eta)$  to zero, we obtain the following set of algebraic equations for  $a_1, a_0, a_{-1}, b_1, b_0, b_{-1}$  and  $c$ , as

$$\begin{cases} C_8 = 0, C_7 = 0, C_6 = 0, C_5 = 0, C_4 = 0, C_3 = 0, C_2 = 0, C_1 = 0, \\ C_0 = 0, \\ C_{-4} = 0, C_{-3} = 0, C_{-2} = 0, C_{-1} = 0. \end{cases} \quad (71)$$

By the same manipulation as illustrated above, we get

(I) The first set is:

$$a_{-1} = 0, \quad b_{-1} = b_{-1}, \quad a_0 = a_0, \quad a_1 = \frac{4b^2\mu^4 b_{-1}(n+2)^2}{a^2 n^4 a_0}, \quad (72)$$

*Archive of SID*

$$c = \frac{\mu^2 b}{n^2}, \quad \mu = \mu, \quad a_2 = 0,$$

$$b_0 = \frac{a_0^3 a^3 n^6 + 4b^3 \mu^6 b_{-1}^2 n^3 + 24b^3 \mu^6 b_{-1}^2 n^2 + 48b^3 \mu^6 b_{-1}^2 n + 32b^3 \mu^6 b_{-1}^2}{b \mu^2 a^2 n^4 (n+2) a_0^2},$$

$$b_1 = \frac{a_0^3 a^3 n^6 + 16b^3 \mu^6 b_{-1}^2 n^3 + 96b^3 \mu^6 b_{-1}^2 n^2 + 192b^3 \mu^6 b_{-1}^2 n + 128b^3 \mu^6 b_{-1}^2}{4b^2 \mu^4 a a_0 n^2 (n+2)^2 b_{-1}},$$

which gives:

$$v_1(x, t) = \frac{a_0 + \frac{4b^2 \mu^4 b_{-1} (n+2)^2}{a^2 n^4 a_0} e^{\left[ \mu \left( x - \frac{\mu^2 b}{n^2} t \right) \right]}}{b_{-1} e^{\left[ -\mu \left( x - \frac{\mu^2 b}{n^2} t \right) \right]} + b_0 + b_1 e^{\left[ \mu \left( x - \frac{\mu^2 b}{n^2} t \right) \right]} + e^{\left[ 2\mu \left( x - \frac{\mu^2 b}{n^2} t \right) \right]}},$$

$$u_1(x, t) = \left\{ \frac{a_0 + \frac{4b^2 \mu^4 b_{-1} (n+2)^2}{a^2 n^4 a_0} e^{\left[ \mu \left( x - \frac{\mu^2 b}{n^2} t \right) \right]}}{b_{-1} e^{\left[ -\mu \left( x - \frac{\mu^2 b}{n^2} t \right) \right]} + b_0 + b_1 e^{\left[ \mu \left( x - \frac{\mu^2 b}{n^2} t \right) \right]} + e^{\left[ 2\mu \left( x - \frac{\mu^2 b}{n^2} t \right) \right]}} \right\}^{\frac{1}{n}}.$$

**Case III:**  $p = c = 2$  and  $q = d = 2$ .

Since the values of  $c$  and  $d$  can be freely chosen, we set  $p = c = 2$  and  $d = q = 2$ . Then the trial function, (7) becomes

$$v(\eta) = \frac{a_2 \exp(2\eta) + a_1 \exp(\eta) + a_0 + a_{-1} \exp(-\eta) + a_{-2} \exp(-2\eta)}{b_2 \exp(2\eta) + b_1 \exp(\eta) + b_0 + b_{-1} \exp(-\eta) + b_{-2} \exp(-2\eta)}. \quad (73)$$

There are some free parameters in (73), we set  $b_2 = 1, b_1 = b_{-1} = 0$  for simplicity, the trial function, (73) is simplified as follows

$$u(\eta) = \frac{a_2 \exp(2\eta) + a_1 \exp(\eta) + a_0 + a_{-1} \exp(-\eta) + a_{-2} \exp(-2\eta)}{\exp(2\eta) + b_0 + b_{-2} \exp(-2\eta)}. \quad (74)$$

Substituting (74) into Eq. (48), to get

$$\frac{1}{A} [C_8 \exp(8\eta) + C_7 \exp(7\eta) + C_6 \exp(6\eta) + C_5 \exp(5\eta) + C_4 \exp(4\eta) + \quad (75)$$

*Archive of SID*

$$C_3 \exp(3\eta) + C_2 \exp(2\eta) + C_1 \exp(\eta) + C_0 + C_{-1} \exp(-\eta) + \\ C_{-2} \exp(-2\eta) + C_{-3} \exp(-3\eta) + C_{-4} \exp(-4\eta) + C_{-5} \exp(-5\eta) + \\ C_{-6} \exp(-6\eta) + C_{-7} \exp(-7\eta) + C_{-8} \exp(-8\eta) = 0,$$

where

$$A = [b_{-1} \exp(-\eta) + b_0 + b_1 \exp(\eta) + \exp(2\eta)]^4, \quad (76)$$

and  $C_n$ 's are coefficients of  $\exp(n\eta)$ . Equating the coefficients of  $\exp(n\eta)$  to zero, we obtain the following set of algebraic equations for  $a_2, a_{-2}, a_1, a_0, a_{-1}, b_0, b_{-2}$  and  $c$ , as

$$\begin{cases} C_8 = 0, C_7 = 0, C_6 = 0, C_5 = 0, C_4 = 0, C_3 = 0, C_2 = 0, C_1 = 0. \\ C_0 = 0, \\ C_{-8} = 0, C_{-7} = 0, C_{-6} = 0, C_{-5} = 0, C_{-4} = 0, C_{-3} = 0, C_{-2} = 0, C_{-1} = 0. \end{cases} \quad (77)$$

By the same manipulation as illustrated above, we get

(I) The first set is:

$$a_1 = 0, \quad a_{-2} = 0, \quad a_{-1} = 0, \quad b_0 = b_0, \quad a_0 = \frac{4b_0\mu^2b(n+2)}{an^2}, \\ c = \frac{4b\mu^2}{n^2}, \quad b_{-2} = \frac{1}{4}b_0^2, \quad a_2 = 0, \quad \mu = \mu, \quad (78)$$

which gives:

$$v_1(x, t) = \frac{\frac{4b_0\mu^2b(n+2)}{an^2}}{\frac{1}{4}b_0^2 \exp \left[ -2\mu \left( x - \frac{4\mu^2b}{n^2}t \right) \right] + b_0 + \exp \left[ 2\mu \left( x - \frac{4\mu^2b}{n^2}t \right) \right]}, \\ u_1(x, t) = \left\{ \frac{\frac{4b_0\mu^2b(n+2)}{an^2}}{\frac{1}{4}b_0^2 \exp \left[ -2\mu \left( x - \frac{4\mu^2b}{n^2}t \right) \right] + b_0 + \exp \left[ 2\mu \left( x - \frac{4\mu^2b}{n^2}t \right) \right]} \right\}^{\frac{1}{n}}. \quad (79)$$

If we choose  $b_0 = 2$  or  $b_0 = -2$  and then the solution (79) respectively gives (cf. Eqs. (38) and (37) in [25])

$$u_{1,1}(x, t) = \left\{ \frac{b\mu^2(n+2)}{2an^2} \operatorname{sech}^2 \left[ \mu \left( x - \frac{4\mu^2b}{n^2}t \right) \right] \right\}^{\frac{1}{n}},$$



*Archive of SID*

$$u_{1,2}(x, t) = \left\{ -\frac{b\mu^2(n+2)}{2an^2} \operatorname{csch}^2 \left[ \mu \left( x - \frac{4\mu^2 b}{n^2} t \right) \right] \right\}^{\frac{1}{n}}.$$

As illustrated in the previous case, the obtained solitary solutions can be converted into periodic solutions or compact-like solutions if  $\mu$  is chosen as an imaginary number. Here, we only discuss the solution given by (79). If  $\mu = i\bar{\mu}$ , then it becomes

$$u_1(x, t) = \left\{ -\frac{\frac{4b\bar{\mu}^2 b_0(n+2)}{an^2}}{(4+b_0^2)\cos(B) + 4b_0 + i(4-b_0^2)\sin(B)} \right\}^{\frac{1}{n}}. \quad (80)$$

where  $B = 2\bar{\mu} \left( x + \frac{4\bar{\mu}^2 b}{n^2} t \right)$ . Elimination of the imaginary part requires that

$$b_0 = \pm 2. \quad (81)$$

We therefore, by applying (81) obtain from (80) the periodic solutions

$$u_1(x, t) = \left\{ \mp \frac{\frac{b\bar{\mu}^2(n+2)}{an^2}}{\cos \left[ 2\bar{\mu} \left( x + \frac{4\bar{\mu}^2 b}{n^2} t \right) \right] \pm 1} \right\}^{\frac{1}{n}}. \quad (82)$$

and (82) is further simplified to obtain a periodic solution respectively (cf. Eqs. (40) and (39) in [25])

$$u_{1,3}(x, t) = \left\{ -\frac{b\bar{\mu}^2(n+2)}{an^2} \sec^2 \left[ \bar{\mu} \left( x + \frac{4\bar{\mu}^2 b}{n^2} t \right) \right] \right\}^{\frac{1}{n}}, \quad (83)$$

$$u_{1,4}(x, t) = \left\{ -\frac{b\bar{\mu}^2(n+2)}{an^2} \csc^2 \left[ \bar{\mu} \left( x + \frac{4\bar{\mu}^2 b}{n^2} t \right) \right] \right\}^{\frac{1}{n}}. \quad (84)$$

Obtained results in above are the exact solutions of the generalized KdV equation.

## 5. Conclusion

Based on the Exp-function method, some nonlinear evolution equations are solved exactly. In this article we investigated the modified KdV

## Archive of SID

equation and the generalized KdV equation. The Exp-function method is a useful method for finding travelling wave solutions of nonlinear evolution equations. This method has been successfully applied to obtain some new generalized solitary solutions to the modified KdV equation and the generalized KdV equation. The Exp-function method is more powerful in searching for exact solutions of NLPDEs. Comparing our results and Wazwaz's results [25] it can be seen that the results are the same. Also, new results are formally developed in this article. It can be concluded that the this method is a very powerful and efficient technique in finding exact solutions for wide classes of problems.

## Acknowledgment

The authors would like to thank the referees for their valuable suggestions and comments.

## References

- [1] S. Abbasbandy, Approximate solution for the nonlinear model of diffusion and reaction in porous catalysts by means of the homotopy analysis method, *Chem. Eng. J.* 136 (2008), 144-150.
- [2] M.A. Abdou, Generalized solitary and periodic solutions for nonlinear partial differential equations by the Exp-function method, *Nonlinear Dyn.* 52 (2008), 1-9.
- [3] M.J. Ablowitz and P.A. Clarkson, *Solitons, nonlinear evolution equations and inverse scattering*, Cambridge University Press, 1991.
- [4] C. L. Bai and H. Zhao, Generalized extended tanh-function method and its application, *Chaos Solitons Fractals.* 27 (2006), 1026-1035.
- [5] A. Boz and A. Bekir, Application of Exp-function method for (3+1)-dimensional nonlinear evolution equations, *Comput. Math. Appl.* 56 (2008), 1451-1456.
- [6] C. Chun, Solitons and periodic solutions for the fifth-order KdV equation with the Exp-function method, *Phys. Lett.* 372 (2008), 2760-2766.
- [7] M. Dehghan, J. Manafian, and A. Saadatmandi, Solving nonlinear fractional partial differential equations using the homotopy analysis method, *Num. Meth. Partial Differential Eq. J.* 26 (2010), 448-479.

*Archive of SID*

- [8] M. Dehghan, J. Manafian, and A. Saadatmandi, Application of semi-analytic methods for the Fitzhugh–Nagumo equation, which models the transmission of nerve impulses, *Math. Meth. Appl. Sci.* 33 (2010), 1384-1398.
- [9] M. Dehghan and M. Tatari, Identifying an unknown function in a parabolic equation with overspecified data via He's variational iteration method, *Chaos Solitons and Fractals.* 36 (2008), 157-166.
- [10] M. Dehghan and F. Shakeri, The use of the decomposition procedure of Adomian for solving a delay differential equation arising in electrodynamics, *Phys. Scr.* 78 (2008), 1-11.
- [11] M. Dehghan and J. Manafian, The solution of the variable coefficients fourth-order parabolic partial differential equations by homotopy perturbation method, *Z. Naturforsch.* 64a (2009), 420-430.
- [12] M. Dehghan and F. Shakeri, Solution of an integro-differential equation arising in oscillating magnetic fields using He's homotopy perturbation method, *Progress In Electromagnetics Research, PIER.* 78 (2008), 361-376.
- [13] E. Fan, Extended tanh-function method and its applications to nonlinear equations, *Phys. Lett. A.* 277 (2000), 212-218.
- [14] J. H. He, *Non-perturbative method for strongly nonlinear problems.* Dissertation, De-Verlag im Internet GmbH, Berlin, 2006.
- [15] J.H. He and X.H. Wu, Exp-function method for nonlinear wave equations, *Chaos Solitons Fractals.* 30 (2006), 700-708.
- [16] J.H. He and M.A. Abdou, New periodic solutions for nonlinear evolution equations using Exp-function method, *Chaos Solitons Fractals.* 34 (2007), 1421-1429.
- [17] R. Hirota, *The Direct Method in Soliton Theory*, Cambridge Univ. Press, 2004.
- [18] X. Lü, H. W. Zhu, X. H. Meng, Z. C. Yang, and B. Tian, Soliton solutions and a Bäcklund transformation for a generalized nonlinear Schrödinger equation with variable coefficients from optical fiber communications, *J. Math. Anal. Appl.* 336 (2007), 1305-1315.

*Archive of SID*

- [19] X. Ma, L. Wei, and Z. Guo, He's homotopy perturbation method to periodic solutions of nonlinear Jerk equations, *J. Sound Vibration*. 314 (2008), 217-227.
- [20] X. H. Menga, W. J. Liua, H. W. Zhua, C. Y. Zhang, and B. Tian, Multi-soliton solutions and a Ba kklund transformation for a generalized variable-coefficient higher-order nonlinear Schrö dinger equation with symbolic computation, *Phys. A*. 387 (2008), 97-107.
- [21] A. Saadatmandi, M. Dehghan, and A. Eftekhari, Application of He's homotopy perturbation method for non-linear system of second-order boundary value problems, *Nonlinear Anal. Real World Appl.* 10 (2009), 1912-1920.
- [22] F. Shakeri and M. Dehghan, Solution of delay differential equations via a homotopy perturbation method, *Math. Comput. Model.* 48 (2008), 486-498.
- [23] A. M. Wazwaz, Multiple-soliton solutions for the KP equation by Hirota's bilinear method and by the tanh-coth method, *Appl. Math. Comput.* 190 (2007), 633-640.
- [24] A. M. Wazwaz, Travelling wave solutions for combined and double combined sine-cosine-Gordon equations by the variable separated ODE method, *Appl. Math. Comput.* 177 (2006), 755-760.
- [25] A. M. Wazwaz, The tanh method for generalized forms of nonlinear heat conduction and Burgers-Fisher equations, *Appl. Math. Comput.* 169 (2005), 321-338.
- [26] A. M. Wazwaz, The tanh-coth method for new compactons and solitons solutions for the  $K(n, n)$  and the  $K(n+1, n+1)$  equations, *Chaos Solitons Fractals*. 188 (2007), 1930-1940.
- [27] A. M. Wazwaz, Multiple-front solutions for the Burgers-Kadomtsev-Petviashvili equation, *Appl. Math. Comput.* 200 (2008), 437-443.
- [28] A.M. Wazwaz, Solitary wave solutions of the generalized shallow water wave (GSWW) equation by Hirota's method, tanh-coth method and Exp-function method, *Appl. Math. Comput.* 202 (2008), 275-286.
- [29] A.M. Wazwaz, Multiple-soliton solutions of two extended model equations for shallow water waves, *Appl. Math. Comput.* 201 (1/2) (2008), 790-799.

*Archive of SID*

- [30] X.H. Wu and J.H. He, Solitary solutions, periodic solutions and compacton-like solutions using the Exp-function method, *Comput. Math. Appl.* 54 (2007), 966-986.
- [31] X.H. Wu and J.H. He, Exp-function method and its application to non-linear equations, *Chaos Solitons Fractals.* 38 (2008), 903-910.
- [32] E. Yusufoglu, A. Bekir, and M. Alp, Periodic and solitary wave solutions of Kawahara and modified Kawahara equations by using Sine-Cosine method, *Chaos Solitons Fractals.* 37 (2008), 1193-1197.
- [33] S. Zhang, Application of Exp-function method to high-dimensional non-linear evolution equation, *Chaos Solitons Fractals.* 38 (2008), 270-276.

**Jalil Manafian Heris**

Department of Mathematics  
Ph.D. student of Mathematics  
Islamic Azad University-Ahar Branch  
Ahar, Iran  
E-mail: manafeian2@gmail.com

**Majid Bagheri**

Department of Mathematics  
Instructor of Mathematics  
Islamic Azad University-Ahar Branch  
Ahar, Iran  
E-mail: majidbagheri53@yahoo.com