

ORIGINAL ARTICLE

Effectiveness of a Social Cognitive Theory Intervention on Promoting Personal Protective Equipment (PPEs) Use among Iranian Aluminum Company Workers

HAMID REZA YAGHOTI¹, AZAR TOL^{1*}, ADEL MAZLOUMI³, ROYA SADEGHI^{1*}, ABBAS RAHIMI FOROUSHANI³, ARMAN GHADERI⁴

¹ Department of Health Education and Promotion, School of Public health, Tehran University of Medical Sciences, Tehran, Iran

² Department of Occupational Health Engineering, School of Public health, Tehran University of Medical Sciences, Tehran, Iran

³ Department of Epidemiology and Biostatistics, School of Public health, Tehran University of Medical Sciences, Tehran, Iran

⁴ Occupational Health Expert, Industrial Health Unit, Iran Aluminum Company (IRALCO), Arak, Iran

Received April 18, 2019; Revised August 22, 2019; Accepted August 29, 2019

This paper is available on-line at <http://ijoh.tums.ac.ir>

ABSTRACT

Workers in industrial workplaces are at increased risk of developing, occupational hazards, work-related accidents, and diseases. Personal Protective Equipment use (PPE) may protect the user against accident or safety risks in the working environment. Educational interventions may improve knowledge and practice related to prevention, but little is known about their knowledge and exposure prevention. This study was aimed to determine the effect of educational intervention based on social cognitive theory (SCT) on knowledge and (PPEs) use among Iranian workers of Aluminum Company (IRALCO). This parallel randomized trial was conducted in a sample of 188 workers who were randomly assigned as the intervention (n=94) and a control (n=94) groups. The intervention group received an educational intervention of six sessions lasting 45-60 minutes augmented with the electronic-learning whereas; the control group attended a routine education program of the company. A researcher-made questionnaire based on SCT constructs and a practice/checklist was used to collect data at baseline and six months after the intervention. Mann-Whitney Test and Wilcoxon Signed Ranks Test analysis were used for data analysis using SPSS version 24.0. $P < 0.05$ was considered statistically significant. After the intervention, knowledge, practices (checklist) items, and all SCT constructs (except emotional adaptability ($P = 0.077$)) revealed significant differences in the intervention group ($P \leq 0.001$) whereas in the control group were no significant differences ($P > 0.05$). This study indicated that the application of educational intervention based on SCT constructs can reflect a positive impact on knowledge and appropriate use of PPEs to reduce occupational-related injuries. This theory is a recommended method to improve workers' personal protective behaviors.

Keywords: Knowledge, Personal Protective Equipment (PPE), Social Cognitive Theory (SCT), Occupational Health, Iranian Aluminum Company Worker

INTRODUCTION

It is estimated that 2.78 million deaths related to occupational accidents occurring annually as a work-related death. Based on the Global Burden of Disease Study, work-related mortality caused 5% of the global total deaths [1]. Also, 68 million cases of work-related illnesses annually around the world and about 170 million working days are lost every year due to work-related accidents [2]. Ignoring safety rules, carelessness, and inappropriate or non-use of PPEs played an important role in event of these accidents; the most important causes of work-related deaths are falling, tensile, electric shock, and burns [1].

Personal Protective Equipment (PPE) consists of protective clothing, helmets, goggles, and other garments or equipment designed to protect the user's body from injury or infection. Protective Equipment may be worn for job-related occupational safety and health purposes [3].

PPE is equipment worn by a worker to diminish exposure to specific hazards. Any item of PPE imposes a barrier between the wearer/user in the workplace. This can create extra pressure on the users; weaken their ability to do their work and make significant discomfort levels [4].

Technical feasibility, low quality of some PPEs, feeling uncomfortable with the existing PPEs, inappropriate training, stereotypical instructions, and inattention to effective incentives for PPEs use lead workers even supervisors to have a little interest in PPEs use in the workplace. Any of these can discourage users from using PPEs correctly, therefore workers are more vulnerable to *be injured*, or *ill-health* over the course or under extreme circumstances may cause death [4].

In this way, in order to correct the existing situation based on the staff's opinions, revision in the selection of appropriate approaches seems necessary [5]. In many industries, the use of engineering techniques is not feasible to reduce harmful factors, and PPEs use is the best way to maintain individuals' health [6]. Concerning the reduction of accidents, the role of education as the axis of advancement and development has a very important role in different fields [2]. Safety training is one of the effective tools

in preventing occupational diseases and accidents. Safety education defined as a process by which workers acquire knowledge, learn new skills and acquire the motivation to do things [7].

Some studies revealed that education as a good prevention strategy for safety and health Industry members should more extensively consider their best practices to protect workers' safety and health [8-9]. But despite the training provided to workers, workers' knowledge of personal protective equipment is still low. Study results on the knowledge and use of personal protective equipment by Welding Workers showed that 31.4% of workers had low knowledge of personal protective equipment and 50.3% of them had moderate knowledge [10].

Preventive behavioral change studies in the safe behavioral field had also demonstrated that the knowledge of potential hazards is insufficient to change behavior and addressing other determinants of behavior, such as attitudes, perceived social norms, and environmental factors that inhibit or encourage behavioral necessity perceive [3-11].

In a study, Amidi Mazaheri et al. found that workers' knowledge of occupational hazards improvement may encourage them to work safely using health education theories [12].

Social Cognitive Theory (SCT) has been widely used in education and health promotion interventions and its effectiveness in improving lifestyle have been confirmed in several studies [13-18]. This theory describes human behavior by triadic mutual causation (behavior, environment, and personal) factors [19-20]. In the theoretic framework of the present study, some constructs as knowledge (learning details and insight obtaining related to PPEs use) outcome expectations (expectation of the possible outcomes which follow as a result of engaging in PPEs use), outcome expectancies (value a worker on the probable outcomes that would result from performing PPEs use), environment (physical or social circumstances of IRALCO), self-efficacy (worker's ability confidence to follow PPEs use), self-efficacy in overcoming impediments (self-confidence that a worker feels to overcome barriers while doing PPEs use), goal setting/self-controlling (setting goals and developing plans to achieve PPEs use), and emotional coping (techniques engaged by a worker to control

Corresponding author: Roya Sadeghi

E-mail: sadeghir@tums.ac.ir

emotional and physiological states associated with gaining of PPEs use) were used [20]. Based on this theory, learning is performed by observation, imitation, and modeling. Moreover, learning is considered an individual as an active creature and confirms the role of thinking and cognitive processes in human learning [20- 21]. This study was grounded in SCT constructs as a conceptual framework for behavioral management to give tailored health education and promotion interventions focusing on knowledge and PPEs use as individual-level factors for behavior change. The purposes of this study were: (1) to increase knowledge of IRALCO workers about PPEs use, (2) improving the mean scores of SCT constructs, and (3) promoting healthy practices for PPEs.

MATERIALS AND METHODS

This randomized intervention trial was conducted in three phases on IRALCO workers at Arak city, south-west of Tehran province, Iran. At the baseline, knowledge and PPEs use was assessed using a self-administered researcher-made questionnaire based on SCT constructs, which served as a need assessment for the other phases of the study. In the second phase, educational intervention was conducted based on SCT constructs during a month and in four educational sessions. In the third phase, six months after the intervention, the questionnaires were completed again by the intervention and control groups and data analysis was performed. Inclusion criteria were workers who live in Arak city, employment as fixed-job in the production line of IRALCO, lack of confirmed occupational diseases and interest in study participating. Participants who did not attend more than two educational sessions were excluded. The protocol was ethically approved by TUMS educational board with application number IR.TUMS.SPH.REC.1396 .4136 and registered Iranian registry of clinical trials (IRCT ID: IRCT20130213012460N17).

In the present study, data gathering instrument was a four-part researcher-made questionnaire based on SCT constructs including:

- Participants' demographic information including age, work experience, level of education, occupational incident history, marital status, occupational disease history, insurance coverage

status, job satisfaction, income, Wife's job, employment status, and shift work (12 items).

- Knowledge about PPEs use (38 items) on a three-point Likert scale from e.g. correct answer (2), I do not know (1), false answer (0).
- SCT constructs (56 items) consist of knowledge (8 questions), outcome expectations (8 questions), outcome expectancies (8 questions), self-efficacy (7 questions), self-control/setting goals (7 questions), environment (6 questions), self-efficacy in overcoming impediments (7 questions), emotional coping (5 questions). All items were rated based on a five-point Likert scale ranging from zero = totally disagree to 4 = totally agree.
- Practice/checklist (43 items) was rated based on a three-points Likert-type scaling { Yes (2), somewhat (1), No (0)}.

The content validity of the instrument was assessed. The scale was reviewed by a panel consisted of ten experts in health, education, promotion, and occupational health. The panel was asked to judge the necessity and relevance of the items in order to calculate the content validity ratio (CVR) and content validity index (CVI). The result of CVR was confirmed by the experts' opinions. The CVR values according to constructs (knowledge, SCT, and checklist) were calculated 0.84, 0.79, and 0.83, respectively. The CVR acceptable domain for ten experts' panel was 0.62, which was satisfied [22]. Using the Waltz and Bausell method based on the experts' opinions, CVI was calculated. The CVI of the questionnaire items were calculated for knowledge items (0.83), SCT constructs (0.83), and checklist items (0.82). In this method, the results of questions with a score of more than 0.79 are satisfied. The results between 70.0 and 79.0 need to be corrected and less than 0.7 are unacceptable and should be removed. Therefore, it could be concluded that the CVI of the scales was accepted [23]. Internal consistency was evaluated by Cronbach's alpha. A coefficient value of more than 0.75 was significantly correlated. The total Cronbach's alpha of SCT questionnaire and Practice were 0.98 and 0.97, respectively.

Workers were invited to participate in the study through oral invitation and advertisements were placed in the main halls of the performed factory by research teams. All of the participants were given detailed information about the study and signed written consent before participation. To ensure privacy

and confidentiality, no names and identification number was written on the questionnaire. Eligible participants were randomly selected among a list of 2700 workers working in different shifts at IRANLCO. Intervention and control groups were selected in somehow to minimize the information bias. Following written informed consent, a group of eligible workers was randomly assigned as intervention and control groups using four-block randomizations. Participants in both groups were invited to complete the questionnaires at the baseline and six months after the intervention. Both intervention and control groups attended in routine educational programs based on safety and EPP use. The practice/checklist was completed by operational managers of the production line during standard operating procedures for data collection. Furthermore, the intervention group had received educational safety and EPP use program based on SCT constructs in six sessions lasting 45-60 min twice a week during a month. After analyzing pre-test results, the educational program was run based on SCT constructs such as knowledge, outcome expectations, outcome expectancies, self-efficacy overcoming impediments, goal setting in self-control, environment, emotional coping for workers within a month in the form of blended training with a specific educational strategy based on each construct.

Lecture and providing fact sheets were utilized for improving knowledge construct; setting achievable goals, self-monitoring, and propose rewards provide for goal setting/self-control construct; provide some opportunities to overcome workers and situational obstacles, provide access to the health system to improve environment construct; using educational films, motivational interviewing, question

& answer and discussion benefits, role play and virtual training via mobile to enhance outcome expectations and outcome expectancies constructs. To improve self-efficacy and self-efficacy in overcoming impediments constructs using different equipment of PPEs in small steps, utilizing persuasion and reinforcement and decreasing stress and hardship related of using PPEs were performed.

The sessions consisted of information on occupational diseases, occupational accidents, PPE use, PPE demonstrations, fit testing, exposure reduction strategies, and barriers to PPEs use. Appropriate educational strategies were established based on SCT constructs to promote appropriate PPE use. After six months of the educational intervention, two groups filled the questionnaire. Knowledge, SCT constructs, and PPEs use were considered as the primary and secondary outcome measurements, respectively. In this study, the sample size was calculated based on the mean and standard deviation data, SCT constructs, and checklist scores with 95% confidence intervals. The sample size was selected through simple random sampling and finally 188 participants out of 94 persons were selected as the intervention and control group. Data were analyzed via SPSS software version 24 using descriptive methods (mean and standard deviation). To assess data description Mann-Whitney Test, Cross-tabs and Wilcoxon Signed Ranks Test, demographics, SCT constructs, knowledge, and practices/checklist were applied and the relationship between the components before and after intervention in two groups was determined. $P < 0.05$ was considered statistically significant. The consort flow diagram of the participants has been presented in Figure 1.




Fig. 1. Consort flow diagram of the study participants

A total of 188 participants were selected as sample size and among them, 94 workers attended the educational programs and 94 workers formed the control group with a response rate of 100 % (n=188). All the participants were male.

At the baseline, the randomized groups did not show any significant difference in terms of demographic variables and work-related information, SCT constructs, knowledge, and practice (checklist) items. In order to find the relationship between variables crosstabs were used which there was no significant difference in demographic variables between intervention and control groups (age, $P=0.255$; work experience $P=0.680$; education level $P=0.315$; occupational incident history $P=0.766$; marital status $P=0.114$; occupational disease history $P=0.315$; insurance coverage status (All the workers

were covered by insurance), job satisfaction $P=0.052$; income $P=0.219$; Wife's job $P=0.592$; employment status $P=0.023$; and shift work $P=0.467$).

According to the results of the Kolmogorov–Smirnov test, knowledge, SCT constructs, and practice (checklist) items were not normally distributed ($p=0.014$, $p=0.008$, and $p=0.017$, respectively). Therefore, non-parametric tests were used to analyze data. Mann-Whitney test was used to compare the mean of knowledge, SCT constructs, and practice (checklist) items of two groups, and the Wilcoxon Signed Ranks Test was applied to compare the results of before and after the intervention in each group. The comparison of the mean and SD of knowledge scores (Table 1) in the intervention group using Wilcoxon's test before and six months after intervention showed a statistically significant increase in the mean score ($P <$

0.001). Before the intervention, there was no significant difference between the two groups about knowledge scores using the Mann-Whitney test, but after the intervention, these differences were significant ($p < 0.001$); whereas, no significant difference was found in the control group ($P > 0.05$).

Applying Wilcoxon Signed Ranks Test, it was also found that after implementing the intervention in the intervention group, total scores of knowledge were remarkably improved ($p < 0.001$); whereas, no significant improvement was found in the control group ($p=0.109$) (Table 1).

Table 1. Knowledge variable before and six months after intervention among participants

Knowledge	Groups	Before	Six months after	P value
		Intervention mean(SD)	Intervention mean(SD)	
Overall	Intervention	6.15(2.18)	7.91(2.00)	<0.001
	Control	6.16(1.68)	6.24(1.55)	0.180
	<i>p</i>	0.32	<0.001	-
Apron	Intervention	4.28(2.00)	5.20(1.08)	<0.001
	Control	4.46(1.40)	4.47(1.41)	0.317
	<i>p</i>	0.772	<0.001	-
Helmet	Intervention	7.51(2.97)	8.50(1.72)	<0.001
	Control	7.48(1.97)	7.52(1.98)	0.317
	<i>p</i>	0.103	<0.001	-
Eye goggles	Intervention	7.50(2.58)	8.31(1.46)	<0.001
	Control	7.32(1.70)	7.36(1.61)	0.317
	<i>p</i>	0.055	<0.001	-
Hearing protection device	Intervention	5.84(2.26)	6.68(1.43)	<0.001
	Control	5.73(1.51)	5.79(1.47)	0.180
	<i>p</i>	0.129	<0.001	-
Hand glove and hand shield	Intervention	5.94(2.26)	6.65(1.52)	<0.001
	Control	5.96(1.70)	6.02(1.68)	0.157
	<i>p</i>	0.462	<0.001	-
Safety boot and gaiter	Intervention	8.03(2.93)	9.26(1.22)	<0.001
	Control	8.01(2.12)	8.05(2.07)	0.157
	<i>p</i>	0.107	<0.001	-
Respiratory protection	Intervention	6.18(2.02)	6.76(1.34)	<0.001
	Control	5.98(1.59)	6.02(1.56)	0.317
	<i>p</i>	0.082	<0.001	-
Fall arrest belt assembly	Intervention	4.83(1.47)	5.34(0.97)	<0.001
	Control	4.70(1.15)	4.74(1.16)	0.317
	<i>p</i>	0.10	<0.001	-
Total	Intervention	56.25(16.92)	64.73(9.34)	<0.001
	Control	56.73(9.02)	57.26(8.29)	0.109
	<i>p</i>	0.104	<0.001	-

The comparison of mean and SD of SCT constructs (see Table 2) in the intervention group using Wilcoxon's test before and after six months intervention showed a statistically significant increase in the mean score ($P < 0.05$). Before the intervention,

there was no significant difference between the two groups in SCT constructs score using the Mann-Whitney test but after the intervention, these differences were significant ($p < 0.001$) except emotional compatibility ($p=0.077$) (refer to Table 2).

Table 2. Changes in SCT constructs before and six months after the intervention among participants

SCT constructs		Before Intervention mean(SD)	Six months after Intervention mean(SD)	<i>P value</i>
Knowledge	Intervention	24.99(7.6)	27.20(4.98)	0.001
	Control	25.03(5.59)	25.63(5.23)	0.109
	<i>p</i>	0.361	0.033	-
Outcome of expectations	Intervention	24.91(6.94)	27.36(4.11)	<0.001
	Control	25.46(5.37)	26.03(4.95)	0.109
	<i>p</i>	0.839	0.043	-
Outcome of expectancies	Intervention	26.68(6.97)	26.46(4.78)	0.003
	Control	24.39(5.95)	24.97(5.68)	0.102
	<i>p</i>	0.321	0.042	-
Self-efficacy	Intervention	20.01(6.44)	21.64(5.03)	0.002
	Control	19.63(5.21)	19.78(5.07)	0.180
	<i>p</i>	0.175	0.002	-
Self-control /Setting goals	Intervention	20.66(6.13)	22.47(4.32)	<0.001
	Control	21.02(5.13)	21.18(4.92)	0.317
	<i>p</i>	0.672	0.019	-
Environment	Intervention	14.71(5.60)	16.28(4.55)	<0.001
	Control	14.28(4.04)	14.23(4.03)	0.655
	<i>p</i>	0.365	0.001	-
Self-efficacy in overcoming impediments	Intervention	17.69(5.93)	20.50(3.82)	<0.001
	Control	19.07(4.04)	19.19(3.80)	0.180
	<i>p</i>	0.207	0.012	-
Emotional coping	Intervention	14.17(4.71)	15.78(3.08)	<0.001
	Control	14.73(3.98)	14.84(3.78)	0.157
	<i>p</i>	0.659	0.077	-

The comparison of mean and SD of Practice (checklist) items (see Table 3) in the intervention group using Wilcoxon's test before and six months after the intervention showed a statistically significant increase in the mean score ($P < 0.05$). Before the intervention, there was no significant difference between the two groups about Practice (checklist) items scores using the Mann-Whitney test, but after the intervention, these differences were significant

($p < 0.001$); whereas, no significant difference was found in the control group ($P > 0.05$). Applying Wilcoxon Signed Ranks Test, it was also found that after implementing the intervention in the intervention group, a total score of practice was remarkably improved ($p < 0.001$); whereas, no significant improvement was found in the control group ($p=0.164$) (refer to Table 3).

Table 3. Practice (checklist) items before and six months after the intervention among participants

Practice(checklist) items		Before	Six months after	<i>p</i>
		Intervention mean(SD)	Intervention mean(SD)	
Eye goggles	Intervention	10.21(4.81)	11.38(4.02)	<0.001
	Control	9.57(3.39)	9.66(3.44)	0.180
	<i>p</i>	0.094	0.001	-
Hearing protection	Intervention	10.21(4.49)	11.60(3.64)	<0.001
	Control	9.84(3.27)	9.91(3.25)	0.180
	<i>p</i>	0.161	0.001	-
Respiratory protection	Intervention	8.30(3.55)	10.18(2.58)	<0.001
	Control	8.51(2.35)	8.60(2.29)	0.180
	<i>p</i>	0.637	<0.001	-
Safety boot and gaiter	Intervention	10.30(4.05)	11.39(3.16)	<0.001
	Control	9.45(3.38)	9.60(3.26)	0.180
	<i>p</i>	0.022	<0.001	-
Helmet	Intervention	5 (2.77)	5.95(2.09)	<0.001
	Control	4.85(2.04)	4.89(2.07)	0.180
	<i>p</i>	0.196	<0.001	-
Overall and apron	Intervention	7.82(3.41)	8.69(2.86)	<0.001
	Control	7.65(2.68)	7.73(2.56)	0.180
	<i>p</i>	0.643	0.012	-
Hand glove and hand shield	Intervention	4.40(1.91)	4.86(1.49)	0.001
	Control	4.27(1.58)	4.31(1.53)	0.317
	<i>p</i>	0.196	0.006	-
Total	Intervention	56.34(21.98)	64.07(15.60)	<0.001
		54.10(14.10)	54.73(13.66)	0.164
		0.10	<0.001	-

DISCUSSION

The purpose of this study was to evaluate the knowledge and PPEs use in an SCT based intervention among Iranian workers at Aluminum Company. The use of workplace intervention to provide education may be particularly helpful in the industrial sector whereas the workplace as an appropriate setting for health-promoting behavior [20]. As Honda et al. in a study, revealed that adherence to appropriate PPE use can be a challenge related to inadequate education on its usage, technical difficulties, and tolerability of PPE

in the workplace but redesigning PPE is a key to improve the safety at the workplace [24].

The study findings revealed that after the intervention, knowledge in the intervention group significantly increased which was in line with studies of Umoren et al. and Adewoye et al. They indicated that the intervention led to knowledge promotion about using personal protective equipment in different workplaces [25-26]. Our finding was also in agreement with the different studies done in Ethiopia and Nigeria among woodworkers which showed

health education intervention improved the use of PPEs [27].

To the best of our knowledge, there was no interventional study based on SCT to compare at the workplace. So, in the current study, some SCT based studies in the other fields were evaluated first. In future studies, it is recommended that this study framework be applied.

The results of this study showed that the mean scores of all SCT constructs (except emotional coping) were increased in the intervention group after the education compared with the control group, with the increase being considered as statistically significant.

These findings were consistent with that of Shojaei et al., findings that showed after the educational program, the mean scores of SCT constructs in the intervention group were increased significantly compared to the pre-intervention and the control groups [28]. This comparison revealed that SCT can be feasible in different settings in a workplace or in the individuals in the community.

In the present study, after the intervention, the mean score of knowledge construction in the intervention group was significantly higher than the control group. These findings were in approved the results of Duncan-Whaley and Shirvani et al., studies [29-30]. Based on Navidian et al. study results, safety educational intervention programs could have a constructive impact on the increasing workers' knowledge, attitude, and adopting safe behaviors. The result of this study was in line with our study and the similarity of developing motivational interviewing as an educational strategy was notable [31].

The difference in the mean scores of outcome expectations and outcome expectancies in the intervention groups was statistically significant in comparison to the control groups after the intervention. The findings by Vanderhoof [6] were consistent with the current study. The results of Abbasian et al., and Hashemi et al., studies among Iranian students in this construction were not consistent with the findings of the present study [32-33]. The reasons mentioned for the difference in the results may be the difference in workplace conditions and different levels of knowledge. Contrary to the findings of this study, in studies by Abbasian et al., and Hashemi et al., the mean score of outcome expectancies did not change significantly [32-33]. It

seems that the use of blended education strategies can be effective to change this structure significantly after the intervention. Also, in Poddar et al. study educational intervention had no effect on outcome expectancies, which could be due to education via Email and superficially education [34]. The study results showed that the mean scores of outcome expectancies and outcome expectation increased significantly after the intervention, which was consistent with the results of Taymoori et al., and Shirvani et al., studies [35-30].

The findings of this study showed a significant increase in self-efficacy in the intervention group in comparison to the control group. The findings of a study entitled "The effect of educational program based on self-efficacy theory on occupational stress management of nurses" showed that the difference between the mean score of self-efficacy of the intervention group compared to the pre-intervention and the control group was statistically significant, which was consistent with the results of this study [36-37]. The results of Stacey et al. and Chan studies showed that educational intervention in the intervention group caused a significant difference in outcome expectancies and self-efficacy compared to the control group [38-39] which was consistent with the results of this study.

The difference in the result of environment construction in the intervention group was statistically significant in comparison to the control group after the intervention. In the study of Lubans and Sylva which was done using a social cognitive theory (SCT), there was a significant difference between the intervention and the control groups at the end of the educational program [40] and it was consistent with the results of the present study. Also, similar results were found by Shirvani et al. who worked using Social Cognitive Theory [30]. The results of two studies conducted by Pirasteh et al., and Taymoori et al., [35-37] were consistent with the findings of the present study.

Similar to the findings, results of Shojaei et al, which evaluated the effects of an educational program based on the Social Cognitive Theory (SCT) on low back pain severity in healthcare workers showed the mean differences between the baseline and 6-month follow-up of four constructs of SCT including self-efficacy, self-control, outcome reinforcement, and emotional coping were significant. However, a significant difference was found in the

intervention group after the educational intervention, and the total mean scores of all predictor constructs were higher in the intervention group than the control group following the intervention [28]. Moreover, six months after the intervention, a significant difference was found between the mean scores of practice (checklist) items between the two groups. Studies carried out among welders in Nigeria [25-26] reported similar findings.

Furthermore, in a study, Quach et al. suggested that the operational managers should continuously be updated about PPE use and its importance. Therefore, researchers may broadly be able to investigate both managers and workers who can have a monitoring control in their workplaces [41].

Overall, the effective implementation of the educational program combined with the high follow up rate, indicated that the intervention was feasible. The study questionnaires were designed based on self-reporting whereas it could increase the likelihood of Socially Desirable Bias. Moreover, this study was conducted in a limited statistical society, so precaution should be exercised when generalizing these results to all workers. Also, in the future researches more detailed questions to define which barriers are associated with individual PPEs use in occupational exposures is recommended. Future efforts may examine long time frame compared to this study to assess the result of sustainability. The study novelties were the application of SCT for the industrial safety intervention and a comprehensive review of the impact of the intervention program on all PPEs use.

CONCLUSION

To the best of our knowledge, in the previous studies, the effect of educational intervention based on social cognitive theory (SCT) on knowledge and PPEs use among Iranian workers of Aluminum Company (IRALCO) was not evaluated. Overall, target group knowledge and practice on PPEs use could be improved through educational efforts. The current study results indicated that the intervention based on SCT can promote knowledge and PPEs use among studied workers. It seems that establishing appropriate workshops to enhance the knowledge and practice of workers about PPEs use. Future research about industrial workers will optimistically fill gaps between occupational risk factors and PPEs use to prevent

adverse health outcomes among workers. According to the important role of managers to audit PPEs use among workers, it is recommended that managers be involved in all the safety and health services delivered by the workplace.

ACKNOWLEDGMENTS

The authors enthusiastically thank all IRALCO staff. We would also like to thank educational chancellor of TUMS for partially funding as a part of MSPH thesis.

CONFLICTS OF INTEREST

The authors have no conflict of interests to declare.

REFERENCES:

1. Hämäläinen P, Takala J, Kiar TB. Global Estimates of Occupational Accidents and Work-related Illnesses, World Congress on Safety and Health at Work., 3-4 September 2017. Singapore.
2. Vahedian Shahroodi M, Mohammadi, A, Tehrani H. A review of studies in the field of knowledge and safe behaviors of workers in Iran. *Journal of Health Literacy*. 2016;1(1): 25-38.
3. Wright T, Adhikari A, Yin J, Vogel R, Smallwood S, Shah G. Issue of compliance with use of personal protective Equipment among wastewater workers across the southeast region of the United States. *Int J Environ Res Public Health*. 2019. 5; 16(11). pii: E2009. DOI: 10.3390/ijerph16112009.
4. Flynn MA, Keller B, DeLaney SC. Promotion of alternative-sized personal protective Equipment. *J Safety Res*. 2017; 63:43-46. DOI: 10.1016/j.jsr.2017.08.004.
5. Shamsi M, Shams M, Nilipour Tabatabaei A. Attitude and behaviors related to using personal protective equipment: Audience analysis to design a behavior change intervention for employees in constructing underground stations in Esfahan. *Iran Occupational Health*. 2013; 10 (3):20-28.

- URL: <http://ioh.iums.ac.ir/article-1-760-en.html>
6. Vanderhoof W. How to get workers to wear it. *Occup Health Saf.* 2004; 73(12):42-44.
 7. Kiani F, Samavatyan H, Poorabdian S, Mansournejad Z, Jafari E. Effectiveness of a safety training course in changing employees' attitude toward safety issues and its dimensions: a pathological study. *SJSPH.* 2011; 9 (2) :53-68. [In Persian].
URL: <http://sjsph.tums.ac.ir/article-1-53-en.html>
 8. O'Connor T, Flynn M, Weinstock D, Zaroni J Occupational safety and health education and training for underserved populations. *New Solut.* 2014; 24(1):83-106. DOI: 10.2190/NS.24.1.d.
 9. Syron LN, Bovbjerg VE, Mendez-Luck CA, Kincl LD. Safety and Health Programs in Alaska's Seafood Processing Industry: Interviews with Safety and Health Managers. *J Agro medicine.* 2019; 24(4):449-461. DOI: 10.1080/1059924X.2019.1639578.
 10. Taghdisi MH, Madadzadeh N, Shadzi SH, Hassanzadeh A. Effects of Education Interventions on the Coke workers' Immune Performances on Baznef Model Basis at Isfahan Melting Factory, 2005. *sjimu.* 2008, 16(3): 1-10. [In Persian].
 11. Balkhyour MA, Ahmad I, Rehan M. Assessment of personal protective Equipments use and occupational exposures in small industries in Jeddah: Health implications for workers. *Saudi J Biol Sci.* 2019; 26(4):653-659. DOI: 10.1016/j.sjbs.2018.06.011
 12. Amidi Mazaheri M, Heydarnia AR, Ghofranipour FA, Shafiei A. The effect of theory-based intervention on promote workers' safe behavior in Isfahan steel company. *Journal of Isfahan medical school (I.U.M.S).* 2010, 27(102):798-808. [In Persian].
 13. Schroer S, Haupt J, Pieper C. Evidence-based lifestyle interventions in the workplace-an overview. *Occupational Medicine. Narnia.* 2014, 64(1):8-12. DOI: 10.1093/occmed/kqt136.
 14. Van Wier MF, Ariëns GA, Dekkers JC, Hendriksen IJ, Pronk NP, Smid T, van Mechelen W. ALIFE@Work: a randomized controlled trial of a distance counselling lifestyle programme for weight control among an overweight working population. *BMC Public Health.* 2006 May 24; 6: 140. DOI:10.1186/1471-2458-6-140
 15. Van Wier, MF, Ariëns GAM, Dekkers JC, Hendriksen IJM, Smid T, Mechelen WV. Phone and e-mail counselling are effective for weight management in an overweight working population: a randomized controlled trial. *BMC Public Health.* 2009; 9(1): 6. DOI: 10.1186/1471-2458-9-6.
 16. JangY, Yoo H. Self-management programs based on the social cognitive theory for Koreans with chronic disease: A systematic review', *Contemporary Nurse Routledge.* 2012; 40(2):147-159. DOI: 10.5172/conu.2012.40.2.147.
 17. Hatchett A, Hallam JS, Ford, MA. Evaluation of a social cognitive theory-based email intervention designed to influence the physical activity of survivors of breast cancer', *Psycho-Oncology.* 2013; 22(4):829-836. DOI: 10.1002/pon.3082
 18. Dennis EA, Potter KL, Estabrooks PA, Davy BM. Weight Gain Prevention for College Freshmen: Comparing Two Social Cognitive Theory-Based Interventions with and without Explicit Self-Regulation Training', *Journal of Obesity. Hindawi Publishing Corporation Journal of Obesity.* 2012; 1-10. DOI: 10.1155/2012/803769.
 19. Stajkovic AD, Luthans F. Social cognitive theory and self-efficacy: Going beyond traditional motivational and behavioral approaches. *Organizational Dynamics. Pergamon,* 1998; 26 (4): 62-74. DOI: 10.1016/S0090-2616(98)90006-7.
 20. Sharma M. Theoretical foundations of health education and health promotion. World Headquarters Jones & Bartlett Learning, third edition, USA, 2016.
 21. Lubans DR, Plotnikoff RC, Morgan PJ, Dewar D, Costigan S, Collins CE. Explaining dietary intake in adolescent girls from disadvantaged secondary schools. A test of Social Cognitive Theory. *Appetite.* 2012; 58(2):517-24. DOI: 10.1016/j.appet.2011.12.012.
 22. Lawshe CH. A quantitative approach to content validity. *Personnel Psychology. John Wiley & Sons, Ltd .*1975, 28(4), pp. 563-575. DOI: 10.1111/j.1744-6570.1975.

23. Pazargadi M, Ashktorab T, Alavimajd H, Khosravi S. Developing an Assessment Tool for Nursing Students` General Clinical Performance. *Iranian Journal of Medical Education*. 2013; 12 (11) :877-887
URL: <http://ijme.mui.ac.ir/article-1-2182-en.html>
24. Honda H, Iwata K. Personal protective equipment and improving compliance among healthcare workers in high-risk settings. *Curr Opin Infect Dis*. 2016; 29(4):400-6.
DOI: 10.1097/QCO.0000000000000280.
25. Umoren Q, Ekanem US, Johnson OE, Olugbemi MO. An assessment of the effect of health education on the use of personal protective Equipments among small scale welders (panel beaters) in Akwa Ibom state, Nigeria`, *International Journal of Community Medicine and Public Health*. 2016; 3(11):3220-3228.
DOI: 10.18203/2394-6040.ijcmph20163940.
26. Adewoye KR, Awoyemi AO, Babatunde OA, Atoyebi OA, Salami SK, Issa FY. Effect of health education intervention on the knowledge and use of personal protective Equipments among small scale electric arc welders in Ilorin, Nigeria. *Indian J Occup Environ Med*. 2014; 18(1):3-8.
DOI:10.4103/0019-5278.134945
27. Ezeugwu L, Aguwa EN, Arinze-Onyia SU, Okeke TA. Health education: Effect on knowledge and practice of workplace personal hygiene and protective measures among woodworkers in Enugu, Nigeria. *Niger J Clin Pract*. 2017; 20(7):867-872.
DOI: 10.4103/njcp.njcp_258_16).
28. Shojaei S, Tavafian SS, Jamshidi A R, Wagner J, Sepahvandi MR. Social Cognitive Theory-Based Intervention and Low Back Pain among Health Care Workers in Qom Hospitals of Iran. *International Journal of Musculoskeletal Pain prevention*. 2017; 2(1): 193-201.
29. Duncan-Whaley DO. Effect of Nutrition and Physical Activity Counseling on Knowledge and Behavior of Elementary Students in a Rural, Coastal Community. Master thesis of Science in Nursing, 2005.
30. Shirvani H, Sanaeinasab H, Tavakoli R, Saffari M, Khalaji K, Me`mar S. The Effect of a Social Cognitive Theory-based Educational Intervention on the Physical Activity of Female Adolescents. *Iran J Health Educ Health Promot*. 2017; 4(4):309-318.
DOI: 10.18869/acadpub.ihepsaj.4.4.309). [In Persian].
31. Navidian A, Rostami Z, Rozbehani N. Effect of motivational group interviewing-based safety education on Workers' safety behaviors in glass manufacturing. *BMC Public Health*. 2015 19; 15: 929. DOI: 10.1186/s12889-015-2246-8.
32. Abbasian F, Omidvar N, Bondarianzadeh D, Rashidkhani B, Shakibazadeh E, Hashemi B. Effect of a School-based Intervention Based on Social Cognitive Theory on Fruit and Vegetable Consumption in Middle School Students in Tehran. *Hayat*. 2012; 17 (4) :73-84
URL: <http://hayat.tums.ac.ir/article-1-48-en.html>[In Persian].
33. Hashemi B, Omidvar N, Bondarianzadeh D, Shakibazadeh E, Rashidkhani B, Abbasian F. Effect of a Family-Based Intervention Based on Social-Cognitive Theory on Fruit and Vegetable Intake of Middle School Female Students in a District of Tehran. *Hakim Health Sys Res*. 2012; 15 (1) :44-52
URL: <http://hakim.hbi.ir/article-1-970-en.html>[In Persian].
34. Poddar KH, Hosig KW, Anderson-Bill ES, Nickols-Richardson SM, Duncan SE. Dairy intake and related self-regulation improved in college students using online nutrition education. *J Acad Nutr Diet*. 2012; 112(12):1976-86.
DOI: 10.1016/j.jand.2012.07.026.
35. Taymoori P, Niknami SH, Tanya Berry T, Lubans D, Ghofranipour F, Kazemnejad A. A school-based randomized controlled trial to improve physical activity among Iranian high school girls. *International Journal of Behavioral Nutrition and Physical Activity*. 2008; 5(1):18.
DOI: 10.1186/1479-5868-5-18.
36. Peyman N, Ezzati Rastegar KH. Effect of an educational program on job tension management in nurses, based on self-efficacy theory. *Modern Care Journal*. 2012; 9 (33) :1-9.
37. Pirasteh A, Asghari A, Faghihzadeh S, Ghofranipour F. Development and validation of psychosocial determinants measures of physical activity among Iranian adolescent girls. *BMC Public Health*. 2008; 8(1): 50.
DOI: 10.1186/1471-2458-8-150.

38. Stacey FG, James EL, Chapman K, Lubans DR. Social cognitive theory mediators of physical activity in a lifestyle program for cancer survivors and carers: findings from the ENRICH randomized controlled trial. *Int J Behav Nutr Phys Act*. 2016 Apr 14; 13:49.
DOI: 10.1186/s12966-016-0372-z.
39. Chan SY. Social cognitive theory based physical activity intervention targeting non-working time physical activity of workers with intellectual disabilities', Open Access Theses and Dissertations. 2014. Available at: https://repository.hkbu.edu.hk/etd_oa/53 (Accessed: 13 June 2019).
40. Lubans, D, Sylva K. Controlled Evaluation of a Physical Activity Intervention for Senior School Students: Effects of the Lifetime Activity Program. *Journal of Sport and Exercise Psychology*. 2006; 28(3):252-268.
DOI: 10.1123/jsep.28.3.252.
41. Quach T, Von Behren J, Tsoh J, Reynolds P, Fu L, Nguyen T, Le M, Nguyen TT. Improving the knowledge and behavior of workplace chemical exposures in Vietnamese-American nail salon workers: a randomized controlled trial. *Int Arch Occup Environ Health*. 2018; 91(8):1041-1050.
DOI: 10.1007/s00420-018-1343-2.