15(2) 886-897

The effects of different dietary vitamin C and iron levels on the growth, hematological and immunological parameters of rainbow trout *Oncorhynchus mykiss* fingerlings

Adel A.¹; Khara H.^{2*}

Received: August 2013

Accepted: September 2014

Abstract

A 60-days growth trial was conducted in a flow-through culture system to examine the effects of different dietary vitamin C and iron levels on the growth, hematological and immunological parameters of rainbow trout (*Oncorhynchus mykiss*) fingerlings (with average initial weight of 5.12 ± 0.02 g). Three levels of vitamin C (150, 200 and 250 mg kg⁻¹ diet) and iron (5, 7 and 9 mg kg⁻¹ diet) (as ferrous sulfate) and their combination were used to prepare diets used in nine experimental treatments with three replicates. The results revealed that the supplementation of basal diet with vitamin C and iron significantly affected the final body weight (FBW), total length (TL), feed conversion ratio (FCR) and specific growth rate (SGR). Among the haematological parameters haematocrit and mean corpuscular volume (MCV) were significantly influenced by vitamin C and iron. Results also showed that lysozyme and IgM concentrations significantly changed among dietary treatments. Based on the results of this study, it can be conclude that addition of vitamin C and iron to the basal diet of cultured rainbow trout will improve the growth rate and well-being of this fish.

Keywords: Growth performance, Hematological and immunological parameters, Vitamin C, Iron, Rainbow trout

¹⁻ Department of Fishery, Guilan Scince and Research Branch, Islamic Azad University, Rasht, Iran

²⁻Department of Fishery, Lahijan Branch, Islamic Azad University, lahijan, Iran, P.O.Box: 1616 *Corresponding author's Email: h.khara1974@yahoo.com

Introduction

The aquaculture industry is growing rapidly worldwide, but continues to contend with some issues such as disease. product quality, feed contamination and environmental impacts. Fish nutrition and its impact on animal welfare is an important aquaculture issue. Poor understanding of the fish dietary requirements in formulation and development of formulated artificial diets for intensive aquaculture have resulted in mortality rates (Halver, 2002). Vitamins are important essential nutrients for aquatic species. Several morphological and functional abnormalities have been various reported in fish species deprived of vitamins. In a research Lall and Ollivier (1993) concluded that in rainbow trout the disease resistance and hormonal antibody production was directly and positively related to the levels of vitamin C. It has been proven that dietary levels of vitamins C enhance the antibody production and immune memory in juvenile of milk fish (Azad et al., 2007). Additionally, previous research findings some indicate that dietary supplementation with immunomodulatory vitamins such as vitamins Can improve the immune response and disease resistance of a variety of cultured fish species (Durve and Lovell, 1982; Verhlac et al., 1998), while some other studies have failed to show positive responses with over fortification of such vitamins (Hardie et al., 1990; Li et al., 1998). So, it is clear that the biological role played by vitamins C is very vital for the sustained growth and health of many living organisms as well as fish. not Nutrients do function as independent units and have some interactions with other nutrients in terms of function and metabolism. Thus the requirement level of a particular nutrient may be affected by the level of another nutrient in either the diet or metabolically in the animal. There is some evidence that iron may be affecting the metabolism of ascorbic acid in the trout (Desjardins, 1985). However, it should be noted that this apparent interaction between iron and ascorbic acid metabolism most be due to the effects of iron supplementation on both the diet rancidity and on the stability of dietary ascorbic acid. However, there is still a paucity of information on the interaction of these two nutrients in fish. The objective of the present study was to assess the effects of different levels of dietary vitamin C and iron on the growth, hematological and immunological parameters of rainbow trout fingerlings.

Materials and methods

Experimental diets

The basal diet formula applied in this study is given in Table 1. Supplemental levels of vitamin C (L-ascorbyl-2polyphosphate (LAPP) (STAY-C-35, 350 g kg⁻¹ ascorbic acid equivalent, DSM, Netherland) at 150, 200 and 250 mg kg⁻¹ and iron (ferrous sulfate, FeSO₄H₂O) at 5, 7 and 9 mg kg⁻¹ were added to the basal diet to prepare nine experimental diets as follow: T₁; 5 mg Fe, T₂: 7 mg Fe, T₃: 9 mg Fe , T₄:150 mg vitamin C, T₅: 200 mg vitamin C, T₆: 250 mg vitamin C, T₇: 5 mg Fe + 150 vitamin C, T₈: 7 mg Fe + 200 mg vitamin C and T₉: 9 mg Fe + 250 mg vitamin C equivalent kg⁻¹ diet. Also, a treatment without addition of vitamin C and iron was used as control. In preparing the diet, dry ingredients were first ground to small particle sizes in a mill. Ingredients were thoroughly mixed, and then fish oil and water were added to obtain a 25% moisture level. Following that, the diets were coldpelleted into 3-mm diameter size using a pasta maker and then were dried in cool drier for 24 h. After drying, the diets were broken up and sieved into appropriate pellet sizes, and were stored in a cool place until used, and then from each diet one sample was taken for chemical analysis.

 Table 1: Composition of the experimental diet (dry weight).

Ingredients	g kg ⁻¹ dry weight
Fish meal	620
Meat powder	60
Wheat flour	100
Soybean cake	50
Fish oil	40
Soybean oil	40
Lecithin	30
Vitamin mixture (Vitamin E free) ^a	15
Mineral mixture ^b	10
Salt	2
Proximate composition	%
Crude protein	49.18
Crude lipid	14.12
Moisture	14.25
Ash	20.70
Crude Energy (kcal kg ⁻¹)	3012

^a Vitamin mixture was manually provided according to feed requirements of the fish and ingredients were obtained from Science Laboratories (Ghazvin, Iran); which each 1000 g vitamin mixture provides vitamin A, 1,600,000 I.U; vitamin D3, 400 000 I.U; thiamin, 6 g; riboflavin, 8 g; niacin, 12 g; pantothenic acid, 40 g; pyridoxine, 4 g; folic acid, 2 g; cyanocobalamin, 8 mg; vitamin C, 60 g; vitamin K3, 2 g; biotin, 240 mg and inositol, 20 g.
 ^bMineral premix (mg kg)1 diet): NaCl, 500; MgSO4.7H2O, 7500; NaH2PO4.2H2O, 12 500; KH2PO4, 15 500; Ca(H2PO4)2*i*H2O, 10 000; (CH2CHCOO)2Ca.5H2O, 1650; ZnSO4.7H2O, 176.5; MnSO4.4H2O, 81; CuSO4.5H2O, 16.5; CoCl2.6H2O, 0.53; KI, 1.59; starch, 147.5.

889 Adel and Khara, The effects of different dietary vitamin C and iron levels on the growth,...

Experimental fish and feeding trial

The experiment was designed in nine treatments with three replicates for 8 weeks. A total of 810 Rainbow trout fingerlings (average weight: 5.12 ± 0.02 g; mean \pm SD) were stocked into 27 fiberglass tanks (200L) at the Rainbow trout farm in Sari city (Mazandran, Iran). Fish were fed four times a day at 2% of body weight with adjustments made in quantity of feed supplied every week. Total fish weight in each tank was measured every 2 weeks for more accurate feeding rate adjustment. A diurnal 12-h l light: dark cycle was provided by fluorescent light, and dissolved oxygen, temperature and pH of the water were monitored throughout the experiment.

Growth performance

At the end of the feeding trial, fish were fasted for 24 hours and then weight end and total length (TL), weight gain (WG), final weight (FW), feed conversion ratio (FCR), specific growth rate (SGR), and condition factor (CF) and survival rate were calculated according to Huang *et al.* (2003).

FCR = dry feed intake (g)/wet WG (g) SGR (% day⁻¹) = (Ln W_f - Ln W_i) × 100/t

 $CF = 100 \times [wet weht (g)/TL (cm)^3]$

Where W_f and W_i were final and initial fish weights, respectively; TL was total length and t is the experimental duration in day.

Hematological parameters analysis

Fishes were anaesthetized in 100 ppm clove powder solution, and then blood

samples were collected via venipuncture and aspirated into a microcentrifuge tube. The first sample was transferred to an eppendorf tube coated with heparin as anticoagulant and was used for hematological indices determination including hematocrit (Ht), number of red blood cell (RBC) and total leukocyte count (WBC). Red blood cell (RBC, $x10^6$) and total leukocyte counts (WBC) were determined with a Neubauer using Rees diluting solution. То obtain the differential counts of leukocytes, that is, the number of lymphocyte, neutrophil, eosinophil and monocytes, we followed the following procedure. The prepared blood smears were first air dried, fixed in 96% ethanol for 30 minutes, stained by Giemsa staining solution for 30 minutes, and were examined under light microscope (Klontz, 1994). Hemoglobin concentration (Hb) was determined with Drabkin's reagent and read at absorbance at 540 nm (Jain, 1993). According to procedure of Rehulka (2000), haematocrit (Ht) was measured in microhaematocrit capillaries, heparinized using а microhematocrit centrifuge (13,000 rpm for 3 min). Mean corpuscular volume (MCV) and mean corpuscular hemoglobin (MCH) were obtained according to the method described by Haney et al. (1992).

Biochemical analysis

Blood was centrifuged at 3000 rpm for 15 minutes in cooling centrifuge for separation of plasma which was stored at -18°C till used for biochemical analysis. Lysozyme level was determined by turbidometric assav according to the method of Sankaran Gurnani, (1972) with slight and modifications. Aliquots (1.75 mL^{-1}) of Micrococcus lysodeikticus suspension (Sigma) (0.375 mg mL⁻¹, 0.05 M PBS, pH 6.2) were mixed with 250 ml⁻¹ of each sample and the optical density was measured after 15 and 180 s by spectrophotometer (Biophotometer Eppendorf) at 670 nm. PBS was used as the blank and results were expressed in amounts of lysozyme (mg) per 1 mg of sample calibrated using a standard curve determined with hens egg white lysozyme (Sigma) in sterile sodium phosphate buffer. IgM content was determined following the method of Puangkaew et al., 2004.

Data analysis

All data were subjected to a one-way analysis of variance (ANOVA) after confirmation of normality and homogeneity of variance. Significance of the differences between means was tested using Duncan's multiple range test (p<0.05). All assays were carried out in triplicates and data are shown as mean ± SD for each dietary group.

Results

The addition of different dietary vitamin C and iron levels to the basal diets significantly affected the FBW, TL, FCR and SGR; as these values were higher in fish fed diet containing 5 mg Fe + 150 vitamin C (T₇) than those fish fed the other treatment (Table 2). Red blood cells (RBC), White blood

hemoglobin cell (WBC), (Hb), lymphocyte, monocyte, mean corpuscular hemoglobin (MCH), and corpuscular mean hemoglobin concentration (MCHC) were not affected by the supplemental vitamin C and iron levels (Tables 3 and 4). However, hematocrit (Hct) and mean corpuscular volume (MCV) were significantly influenced by different dietary vitamin C and iron levels. At the end of the experiment, significant differences were also observed in IgM (Fig. 1) and lysozyme (Fig. 2) in rainbow trout fed with diets containing different levels of vitamin C and iron.

Discussion

The results of the present study showed significant differences on FBW, TL, FCR and SGR among treatments with different dietary vitamin C and iron levels. Similarly, studies with common carp, (Gouillou-Coustans et al., 1998) hybrid tilapia, Oreochromis niloticus (Shiau and Hsu, 2002), Japanese seabass, Lateolabrax japonicus (Ai et 2004), juvenile al., grouper, Epinephelus malabaricus (Lin et al., 2005) and juvenile cobia, Rachycentron canadum (Zhou et al., 2012) indicate positive effect of vitamin C on the growth. Our results indicated that dietary vitamin C could improve the growth performance of iuvenile rainbow trout. However, Ai et al., (2006) reported that dietary vitamin C levels didn't influence the growth performance of the juveniles of large yellow croaker.

Treatments	Initial weight (g)	Final weight (g)	WG (%)	TL (cm)	FCR	SGR (% day ⁻¹)	Condition factor	Survival rate (%)
Control	5.12 ± 0.02	$24.6\pm5.3~^{\rm c}$	27.6 ± 0.71	13.5 ± 1.08	$1.5\pm0.04\ ^{ab}$	$2.06\pm0.25~^{de}$	1.04 ± 0.31	100
T_1	5.13 ± 0.02	$25.8\pm7.1\ ^{d}$	28.7 ± 1.1	14.1 ± 1.17	$1.4\pm~0.03^{~bc}$	$2.12\pm0.33~^d$	1.04 ± 0.05	100
T ₂	5.14 ± 0.01	$27.1\pm3.5~^{\text{c}}$	29.7 ± 1.3	14.2 ± 1.10	$1.3\pm0.01\ ^{bcd}$	$2.18\pm0.15~^{cd}$	0.96 ± 0.02	100
T ₃	5.12 ± 0.06	$28.2\pm7.5~^{\rm f}$	30.7 ± 1.8	14.9 ± 0.77	$1.2\pm0.02\ ^{cd}$	2.23 ± 0.32 °	$1.04\pm\ 0.09$	100
T_4	5.15 ± 0.02	$21.6\pm4.9^{\ a}$	25.3 ± 1.4	13.6 ± 0.17	$1.6\pm0.04~^a$	$1.91 \pm 0.25 \ {\rm f}$	1.03 ± 0.11	100
T ₅	5.10 ± 0.05	23 ± 9.9 b ^g	26.3 ± 1.6	13.4 ± 1.53	$1.4\pm0.04^{\ bc}$	$1.99\pm0.49~^{\text{c}}$	1.06 ± 0.16	100
T ₆	5.11 ± 0.04	$31{\pm}~7.9~b^{~ef}$	33 ± 1.2	15.3 ± 0.60	$1.1\pm0.02~^{e}$	$2.36\pm0.32~^a$	1.12 ± 0.06	100
T ₇	5.12 ± 0.03	$29.4\pm4.1~^d$	31.5 ± 1.09	14.6 ± 0.78	$1.2\pm0.01^{\ cd}$	2.28 ± 0.17 ^b	1.13 ± 0.02	100
T ₈	5.13 ± 0.02	$27.6\pm1.9\ ^{d}$	30.1 ± 0.61	14.5 ± 0.96	$1.3\pm0.02^{\ bcd}$	2.21± 0.08 °	0.93 ± 0.03	100
T ₉	5.13 ± 0.01	$25.8\pm3.3\ ^{c}$	28.7 ± 1.1	13.8 ± 0.96	$1.3\pm0.01^{\ bcd}$	$2.12\pm0.15\ ^{d}$	1.04 ± 0.07	100

Table 2: Final body weight (FBW), weight gain (WG) (g/fish), total length (TL), feed conversion ratio (FCR), specific growth rate (SGR) and condition factor (CF) of rainbow trout fed the experimental diets for 8 weeks.

Within a column, means with different superscripts are significantly different (p < 0.05).

Treatments	RBC (×10 ⁶)	WBC (×10 ³)	Hematocrit (%)	Hemoglobi n (gr/dl)	MCV (fl)	MCH (pg)	MCHC (gr/dl)
Control	101 ± 21	67.6 ± 15	36 ± 2.6 °	6.3 ± 0.49	$299\pm21.1~^{bcd}$	52.6 ± 6.3	17.5 ± 1.6
T_1	97 ± 16.2	69.6 ± 8.9	$33.3\pm2.8~^{cde}$	6.5 ± 0.42	289 ± 38.2^{bcd}	56.4 ± 7.1	19.5 ± 0.2
T_2	102 ± 21.5	69.3 ± 14	31.6 ± 1.1 ^{cd}	5.7 ± 1.2	303.8 ± 18.7 ^d	59.1 ± 6.2	17.9 ± 2.2
T ₃	101 ± 10.3	64 ± 15	$28.3\pm1.5~^{ab}$	5.9 ± 0.13	$316\pm49.5~^{cd}$	59.5 ± 10.3	18.8 ± 0.36
T_4	103 ± 12.7	64 ± 16.2	$32 \pm 1^{\ cd}$	5.9 ± 0.81	$244\pm11.5~^{ab}$	51.1 ± 6.7	20.9 ± 2.01
T ₅	101 ± 21.7	64.6 ± 15.5	$30.6\pm1.1^{\ bcd}$	6.1 ± 0.11	$300\pm25.7~^{bcd}$	56.1 ± 8.7	18.6 ± 1.8
T ₆	100 ± 13.9	61 ± 12.2	29 ± 4 ^{abc}	5.8 ± 0.49	$273\pm10.2~^{abc}$	54.2 ± 0.09	19.9 ± 0.76
T ₇	102 ± 16	62.6 ± 9.2	$28\pm1~^a$	5.7 ± 0.04	$298\pm51.6^{\text{bcd}}$	61.1 ± 15.8	20.2 ± 2.02
T_8	103 ± 10.3	66 ± 7.5	$33.3\pm2.3~^{cde}$	6.2 ± 1.4	$232\pm20.2~^{a}$	47.9 ± 2.4	20.6 ± 0.75
T ₉	102 ± 17.3	64 ± 13.7	36 ± 2.8 °	6.3 ± 0.49	$254\pm14.1\ ^{ab}$	46.9 ± 9.1	18.6 ± 4.7

Table 3: Hematological parameters of rainbow rout fingerling fed 8 weeks with diets containing different levels of vitamin C and iron.

Within a column, means with different superscripts are significantly different (p < 0.05).

Treatments	Monocyte (%)	Neutrophil (%)
Control	99 ± 1	1 ± 1
T_1	99.5 ± 0.58	0.67 ± 0.57
T_2	98 ± 1.03	1 ± 1
T ₃	98.6 ± 1.5	1.3 ± 0.53
T_4	99.6 ± 0.58	0.33 ± 0.05
T ₅	98.3 ± 2.08	2 ± 0.73
T ₆	99.3 ± 1.1	0.67 ± 0.16
T ₇	98.3 ± 1.5	1.67 ± 0.53
T_8	99 ± 1	1 ± 1
Т9	99.6 ± 0.58	0.33 ± 0.58

Table 4: Differential count of leukocyteof of rainbow rout fingerling fed 8
weeks with diets containing different levels of vitamin C and iron.

Within a column, means with different superscripts are significantly different (p < 0.05).



Figure 1: Total IgM concentration of concentration of rainbow rout fingerling fed 8 weeks with diets containing different levels of vitamin C and iron.



Figure 2: Lysozyme activity of rainbow rout fingerling fed 8 weeks with diets containing different levels of vitamin C and iron.

Studies with common carp (Sakamoto and Yone, 1978a), red sea bream (Sakamoto and Yone, 1978b), Atlantic salmon (Andersen et al., 1996), channel catfish (Barros et al., 2002) and juvenile gibel carp, Carassius auratus gibelio (Pan et al., 2009) did not indicate an adverse effect of iron deficiency on the growth performance. However, Shiau and Su, (2002)reported that weight gain of Nile tilapia increased with dietary iron while significantly decreasing at iron levels. These discrepancies might be due to differences in individual size. development stage, and cultivation environment. There were significant differences in Hct and MCV values among our treatments. Similar results were observed in pirarucu (Arapaima gas) and cobia juvenile's R. canadum (Zhou et al., 2012) fed with diet not supplemented with vitamin C. In agreement with our results Pan et al.,

(2009) also have reported that addition of iron to the basal diet significantly affected the Hct in juvenile gibel carp, C. auratus gibelio. Vitamin C is a powerful antioxidant protecting against oxidative damage to various tissues of fish including red blood cells (Sahoo and Mukherjee, 2003). Lower red blood cells were regarded as the sun of anemia which has been reported in most studies with fish where vitamin C deficiency was observed (NRC, 2011). Moreover. Adham et al. (2000)demonstrated that feeds with insufficient vitamin C cause anemia, characterized by a decrease in the hemoglobin, reduction in the number of erythrocytes and hematocrit. Many researchers have shown that ascorbic acid is involved in the metabolism of iron in fish and a deficiency of ascorbic acid has been observed to cause a reduction in serum iron levels and a redistribution of tissue iron stores in

rainbow trout (Hilton et al., 1989) and a reduction in both hemoglobin and hematocrit levels in catfish, trout and snakehead fish (Lim and Lovell, 1978; Agrawal and Mahajan, 1980). There are also some as that may indicate the effect of iron on ascorbic acid metabolism in trout. Despite the interaction of ascorbic acid and iron metabolism in fish, it is interesting to note that increasing the levels of ascorbic acid do not appear to affect the absorption of dietary iron in fish (Lanno et al., 1985). The innate immune system of fish is regarded to be the first line of defense against pathogens and is more important for fish as compared with mammals. Lysozyme is released by leukocytes and plays a crucial role in antimicroorganism activity. In the present study, fish fed with different dietary vitamin C and iron levels had lower lysozyme activity than fish fed with the basal diets. Previous studies indicated that the lysozyme activity have been found to be positively correlated with supplementation of ascorbic acid (Roberts et al., 1995; Ortuno et al., 1999; Anbarasu and Chandran, 2001; Ai et al., 2004; Ai et al., 2006). In the present study, rainbow trout fed the basal diet had higher concentration than fish fed the other diets. Contrary to our results, Zhou et al., (2012) showed that cobia, R. with diet canadum fed not supplemented with vitamin C had lower concentration than fish fed the other diets. In conclusion, more studies using basal diets containing different vitamins and minerals should be carried out to

examine the effect of dietary vitamins and minerals on juvenile rainbow trout growth performance and haematological parameters.

Acknowledgements

"Financial support by Lahijan Branch, Islamic Azad University Grant No. 17.20.5.3511 is gratefully acknowledged. "

References

- Adham, K.G., Hashem, H.O. and Abu-Shabana, M.B., 2000. Vitamin C deficiency in the catfish *Clarias* gariepinus. Aquaculture Nutrition, 6(2), 129-139.
- Agrawal N.K. and Mahajan, C.L., 1980. Hematological changes due to vitamin C deficiency in Channapunctut Block. *Journal of Nutrition*, 110, 2172-2181.
- Ai, Q.H., Mai, K.S., Zhang, C.X., Xu,
 W., Duan, Q.Y. and Tan, B.P.,
 2004. Effects of dietary vitamin C on growth and immune response of Japanese seabass, *Lateolabrax japonicus*. Aquaculture, 242, 489-500.
- Ai, Q.H., Mai, K.S., Tan, B.P., Xu,
 W., Zhang, W.B. and Ma, H.M.,
 2006. Effects of dietary vitamin C on survival, growth, and immunity of large yellow croaker, *Pseudosciaena crocea*. Aquaculture, 261, 327-36.
- Anbarasu, K. and Chandran, M.R., 2001. Effect of ascorbic acid on the immune response of the catfish, *Mystus gulio* (Hamilton), to different bacterins of *Aeromonas hydrophila*.

895 Adel and Khara, The effects of different dietary vitamin C and iron levels on the growth,...

Fish and Shellfish Immunology, 11, 347-55.

- Andersen, F., Maage, A. and Julshamn K., 1996. An estimation of dietary iron requirement of Atlantic salmon, Salmo salar L., parr. Aquaculture Nutrition, 2, 41– 47.
- Azad, I.S., J.S. Dayal, M. Poornima and S.A. Ali., 2007. Supra dietary levels of vitamins C and E enhance antibody production and immune memory in juvenile milkfish *Chanos chanos* (Forsskal) to formalin-killed Vibrio vulnificus. *Fish and Shellfish Immunolog*, 23, 154-163.
- Barros, M.M., Lim, C. and Klesius P.K., 2002. Effect of soybean meal replacement cottonseed meal and iron supplementation growth, immune response and resistance of channel catfish (*Ictalurus puctatus*) to *Edwardsiella ictaluri* challenge. *Aquaculture*, 207, 263–279.
- **Desjardins, L.M., 1985.** The effect of iron supplementation on diet rancidity and on the growth and physiological response of rainbow trout. MSc thesis, University of Guelph, Ontario, Canada.
- **Durve, V.S. and Lovell R.T., 1982.** Vitamin C and disease resistance in channel catfish (*Ictalurus punctatus*). *Canadian Journal of Fisheries and Aquatic Science*, 39, 948-951.
- Gouillou-Coustans, M.F., Bergot, P. and Kaushik, S.J., 1998. Dietary ascorbic acid needs of common carp (*Cyprinus carpio*) larvae. *Aquaculture*, 161, 453-461.

- Halver J.E., 2002. The vitamins. In: Halver, J.E., Hardy R.W., (Eds), Fish Nutrition. Academic press, NewYork, 62-141.
- Klontz, G.W., 1994. Fish Hematology.In: Techniques in Fish Immunology,Stolen, J. S.; Flecher, T. C.; Rowely,A. F.; Zelikoff, T. C.; Kaattari S. L.and Smith S. A. (Eds.). Vol. 2, SOSPublications, USA. pp. 121-132.
- Haney, D.C., Hursh, D.A., Mix, M.C.
 and Winton, J.R., 1992.
 Physiological and haematological changes in chum salmon artificially infected with erythrocytic necrosis virus. *Journal of Aquatic Animal Health*, 4, 48-57.
- Hardie, L.J., Fletcher, T.C. and Secombes, C.J., 1990. The effect of vitamin E on the immune response of the Atlantic salmon (*Salmo salar* L.). *Aquaculture*, 87, 1–13.
- Hilton, J.W., 1989. The interaction of vitamins, minerals and diet composition in the diet of fish. *Aquaculture*, 79, 223-244.
- Huang, C.H., Chang, R.J., Huang, S.L. and Chen, W., 2003. Dietary vitamin E supplementation affects tissue lipid peroxidation of hybrid tilapia, *Oreochromis niloticus* × *O. aureus. Comparative Biochemistry and Physiology*, 134, 265-70.
- Jain NC.,1993. Essentials of Veterinary Hematology, Lea and Feber, Philadelphia, 417P.
- Lall, S.P, Oliver, G., Weerakoon, D.E.M., Hines, J.A., 1989. The effect of vitamin C deficiency and excess on immune response in Atlantic salmon Salmo salar L. In:

Proceedings of 3rd International Symposium on Feeding and Nutrition in Fish, Toba, Japan 28, August–1 September. pp. 427–441.

- Lanno, R.P., Slinger, S.J. and Hilton J.W., 1985. Effect of ascorbic acid on dietary copper toxicity in rainbow trout (*Salmo gairdneri*, Richardson). *Aquaculture*, 49, 269-287.
- Li, M.H., Wise, D.J. and Robinson, E.H., 1998. Effect of dietary vitamin C on weht gain, tissue ascorbate concentration, stress response, and disease resistance of channel catfish *Ictalurus punctatus. Journal of World Aquaculture Society*, 29, 1-8.
- Lim, C. and Lovell, R.T., 1978. Pathology of vitamin C deficiency syndrome in channel catfish. *Journal of Nutrition*, 108, 1137-1141.
- Lin, M.F. and Shiau, S.Y., 2005. Dietary L-ascorbic acid affects growth, nonspecific immune responses and disease resistance in juvenile grouper, *Epinephelus malabaricus*. *Aquaculture*, 244, 215-21.
- National Research Council (NRC), 2011. Nutrient requirements of fish and shrimp. Washington, DC, USA: National Academies Press, pp. 207-209.
- Ortuno, J., Esteban, M.A. and Meseguer, J., 1999. Effect of hh dietary intake of vitamin C on nonspecific immune response of gilthead seabream (*Sparus aurata* L). Fish and Shellfish Immunology, 9, 429-443.
- Pan, L., Xie, S., Zhu, X., Lei, W., Han, D. and Yang, Y., 2009. The

effect of different dietary iron levels on growth and hepatic iron concentration in juvenile gibel carp (*Carassius auratus gibelio*) Journal of Applied Ichthyology, 25, 428-431.

- Puangkaew, J., Kiron, V., Somamoto, T., Okamoto, N., Satoh, S., Takeuchi, T. and Watanabe T.,
 2004. Nonspecific immune response of rainbow trout (*Onchorhynchus mykiss*, Walbaum) in relation to different status of vitamin E and hhly unsaturated fatty acids. *Fish and Shellfish Immunology*, 16, 25–39.
- Rehulka, J., 2000. Influence of astaxanthin on growth rate condition and some blood indices of rainbow trout *Oncorhynchus mykiss*. *Aquaculture*, 190, 27-47.
- Roberts, M.L., Davies, S.J. and Pulsford, A.L., 1995. The influence of ascorbic acid (vitamin C) on nonspecific immunity in the turbot (*Scophthalmus maximus* L.). *Fish and Shellfish Immunology*, 5, 27-38.
- Sakamoto, S. and Yone, Y., 1978a. Iron deficiency symptoms of carp. Bulletin Japanese Society Science of Fisheries, 44, 1157-1160.
- Sakamoto, S. and Yone, Y., 1978b. Requirement of red sea bream for dietary iron - II. Bulletin Japanese Society Science of Fisheries, 44, 223–225.
- Sankaran K. and Gurnani, S., 1972. On the variation in the catalytic activity of lysozyme in fishes Indian. *Journal of Biochemistry and Biophysics*, 9, 162-165.
- Sahoo, P.K. and S.C. Mukherjee., 2003. Immunomodulation by dietary

897 Adel and Khara, The effects of different dietary vitamin C and iron levels on the growth,...

vitamin C in healthy and aflatoxin B1-induced immunocompromised rohu (*Labeo rohita*). Comparative Immunology, Microbiology & Infectious Diseases, 26, 65–76.

- Shiau, S.Y. and C.Y. Hsu, 2002. Vitamin E sparing effect by dietary vitamin C in juvenile hybrid tilapia *Oreochromis niloticus* x *O. Aureus. Aquaculture*, 210, 335-342.
- Shiau, S.Y. and Su, L.W., 2002. Ferric citrate is half as effective as ferrous sulfate in meeting the iron requirement of juvenile tilapia, Oreochromis niloticus × O. aureus. Journal of Nutrition, 133, 483-488.
- Verhlac, V., Obach, A., Gabaudan, J., Schuep, W. and Hole, R., 1998. Immunomodulation by dietary vitamin C and glucan in rainbow trout (Oncorhynchus mykiss). Fish and Shellfish Immunology 8, 409– 424.
- Zhou, Q.C., Wang, L.G., Wang, H.L., Xie, F.J., Wang, T., 2012. Effect of dietary vitamin C on the growth performance and innate immunity of juvenile cobia (*Rachycentron canadum*). Fish and Shellfish Immunology, 32, 969–975.