

## Effects of microelements (Fe, Cu, Zn) on growth and pigment contents of *Arthrospira (Spirulina) platensis*

Akbarnezhad M.<sup>1</sup>; Shamsaie Mehrgan M.<sup>1\*</sup>; Kamali A.<sup>1</sup>; Javaheri Baboli M.<sup>2</sup>

Received: July 2016

Accepted: November 2017

### Abstract

A laboratory experiment was conducted to assess the bioaccumulation of microelements Fe<sup>2+</sup>, Cu<sup>2+</sup>, Zn<sup>2+</sup> and their effects on the growth and pigment contents of *Spirulina platensis* in Zarrouk's media. The mentioned metals concentrations enhanced separately to tenfold of the Zarrouk's content. The results indicated no differences in the dry weights for the different medias ( $p>0.05$ ). A rapid increase in optical density values has been observed in tenfold of Cu<sup>2+</sup> concentration giving the maximum optical density values ( $0.498 \pm 0.234 \text{ mg L}^{-1}$ ). In contrast, a gradual increase rate in the optical density has been observed at all concentration of treatments. Enrichment factor (EF) improved with increasing the metal concentration of treatments. The maximum EF value has been observed at 13.54 in Zn<sup>2+</sup> concentrations. At all concentrations, the maximum production of chlorophyll A ( $1.213 \pm 0.514 \text{ mg L}^{-1}$ ) occurred in 7<sup>th</sup> day of incubation. The highest total carotenoid concentration was recorded in *S. platensis* treated with Cu<sup>2+</sup> ( $0.0042 \pm 0.0004 \text{ mg L}^{-1}$ ) after 14 days. The phycobiliproteins decreased in all of the treatments. Results suggested that *S. platensis* would be important part of in functional food developments.

**Keywords:** Growth, Bioaccumulation, Microelements, Pigments, *Spirulina platensis*.

1-Department of Fisheries, Science and Research Branch, Islamic Azad University, Tehran, Iran.

2-Department of Fisheries Science, Ahwaz Branch, Islamic Azad University, Ahwaz, Iran.

\*Corresponding author's Email: drshamsaie@gmail.com; m.shamsaie@srbiau.ac.ir

## Introduction

*Arthrospira (Spirulina) platensis* is a photosynthetic filamentous, helical shaped, multicellular, and green-blue microalga that grows vigorously in strong sunshine under high temperatures and alkaline conditions (Sanchez *et al.*, 2003; Habib *et al.*, 2008). *Spirulina* sp. has a high content of protein (up to 70%), along with high amounts of essential fatty acids, essential amino acids, minerals (such as iron, copper, zinc), vitamins (especially B12), antioxidant pigments (i.e. phycobiliproteins, carotenoids, and chlorophyllp-A) and polysaccharides (Belay *et al.*, 1993; Vonshak, 1997). The commercial production of *Spirulina* sp. has gained worldwide attention as a human food supplements, animal feed (terrestrial, fresh water and marine) and pharmaceuticals. In aquaculture, *Spirulina* sp. is using as an additive to improve growth, coloration and probiotic agent (Ramakrishnan *et al.*, 2008; Ghaeni *et al.*, 2011; Ansarifard *et al.*, 2018). The growth of microalgae and the composition of the biomass produced depend on many factors such as nutrient availability, temperature, and light (Sabzi *et al.*, 2018; Hadizadeh *et al.*, 2019).

Trace metals such as iron (Fe), manganese (Mn), cobalt (Co), zinc (Zn), copper (Cu) and nickel (Ni) are essential components which required by microalgae for various metabolic functions and growth (Bruland *et al.*, 1991; Rastar *et al.*, 2018). It has been reported that iron promotes the growth of cyanobacteria and increase photosynthesis and nitrogen fixation

(Rueter and Petersen, 1987). Microalgae utilize zinc as an enzyme cofactor; and, the higher concentrations are toxic to most aquatic life (Omar, 2002). Szabolcs *et al.* (2013) investigated on four microelements such as Fe<sup>+2</sup>, Cu<sup>+2</sup>, Zn<sup>+2</sup> and Mo<sup>+2</sup> on *S. platensis* and *Chlorella vulgaris* and proved to enhanced bioaccumulation ability and biomass. Arunakumara *et al.* (2008) results showed that bioaccumulation of Pb<sup>2+</sup> at low concentrations (5 µg ml<sup>-1</sup>), cause stimulate its growth slightly, although the chlorophyll α and β carotene were decreased. Balaji *et al.* (2014) investigated the effects of zinc and nickel on growth of different *Spirulina* sp. strains which significantly affected by the concentration of selected metals in the culture medium. Maximum growth has exhibited by *S. maxima* at 0.01mM zinc (6.9 mg L<sup>-1</sup>) and 0.01mM nickel (17 mg L<sup>-1</sup>). Deniz *et al.* (2011) reported that growth and chlorophyll A contents in *Spirulina* sp. decreased at most exposure levels to copper and sodium chloride, with increase in carotenoid pigment. Okmen *et al.*, (2011) reported that lower zinc concentration (2.5 mg L<sup>-1</sup>) were stimulated the biomass, chlorophyll A, total carbohydrate and protein content in *Anabaena* sp. GO1 and *Gloeotheca* GO9 cyanobacteria. Zinc trace element in the 10 mg L<sup>-1</sup> and more concentrations were decreased the biomass and other parameters except for *Anabaena* sp. GO2 species. Most of the previous studies have focused on the remediation of heavy metals by microalgae species to eliminate toxic

elements (Romera *et al.*, 2007). The objective of this study was to examine the accumulation of selected important microelements by *S. platensis* and its effects on growth and biopigments accumulation.

### Materials and methods

#### *Microorganism and applied chemicals*

*S. platensis* was obtained from the Research laboratory, Agriculture and Natural Resources of Ahwaz Islamic Azad University, Ahwaz, Iran. All the applied reagents and chemicals have been obtained from either Merck and/or Sigma-Aldrich companies.

#### *Culture medium*

The composition of the growth media in the case of *spirulina platensis* was in accordance with Zarrouk's medium

(Zarrouk, 1996). Three metals ( $\text{Fe}^{+2}$ ,  $\text{Cu}^{+2}$  and  $\text{Zn}^{+2}$ ) were chosen for experiments with concentration enhanced separately to tenfold of the Zarrouk's media (control) (OECD, 2011). Four different solutions have been prepared with control and three others with diverse metal concentrations. The metal content of the control was in accordance with the Zarrouk guideline, as  $[\text{FeSO}_4.7\text{H}_2\text{O}]$ :  $0.01 \text{ g L}^{-1}$ ,  $[\text{ZnSO}_4.4\text{H}_2\text{O}]$ :  $0.222 \text{ g L}^{-1}$ ,  $[\text{CuSO}_4.5\text{H}_2\text{O}]$ :  $0.079 \text{ g L}^{-1}$ . The metal solutions have prepared from  $\text{FeSO}_4.7\text{H}_2\text{O}$ ,  $\text{ZnSO}_4.4\text{H}_2\text{O}$ ,  $\text{CuSO}_4.5\text{H}_2\text{O}$  (Table 1). All experiments have been performed in triplicates ( $n=3$ ).

**Table 1: Applied metal concentrations in the growth media.**

Growth media	$\text{FeSO}_4.7\text{H}_2\text{O}$	$\text{Fe}^{2+}$	$\text{ZnSO}_4.4\text{H}_2\text{O}$	$\text{Zn}^{2+}$	$\text{CuSO}_4.5\text{H}_2\text{O}$	$\text{Cu}^{2+}$
control*	$0.01 \text{ g L}^{-1}$	$0.002 \text{ g L}^{-1}$	$0.222 \text{ g L}^{-1}$	$0.050 \text{ mg L}^{-1}$	$0.079 \text{ g L}^{-1}$	$0.020 \text{ mg L}^{-1}$
Enhanced media(Fe 10fold)	0.1	0.02	0.222	0.050	0.079	0.020
Enhanced media(Zn 10fold)	0.01	0.002	2.22	0.50	0.079	0.020
Enhanced media(Cu 10fold)	0.01	0.002	0.222	0.050	0.79	0.20

\*Control is accordance with Zarrouk media.

#### *Culture conditions and growth*

The alga growing apparatus consists of a horizontal glass surface where on the Erlenmeyer flasks had been placed. Erlenmeyer flasks of 1000 ml capacity have prepared containing 100 ml *S. platensis* (10%) with initial optical density 0.019 (Biomass concentration of  $0.002 \text{ g L}^{-1}$  dry weight) and 200 ml Zarrouk media (Zarrouk, 1966) at temperature  $32 \text{ }^\circ\text{C}$ , pH 8.7, salinity 20 ppt with an illumination of 2500 lux

light intensity, with a light/dark cycle of 12/12 h. Fresh air was pumped into the solution through plastic tubes in order to avoid the generation of alga film layer on the wall of the flasks for a period of 14 days.

#### *Growth of microalgae*

Biomass concentration was determined every day by measuring the optical density at 560 nm to produce a standard curve. Standard curve has been

subsequently used to calculate the biomass of individual samples based on their optical density (Gupta *et al.*, 2006; Rastar *et al.*, 2018).

#### *Determination of dry weight*

The dry weight of biomass has been determined by filtration of sample (15 ml) through dried Whatman filter (pore size 0.42  $\mu\text{m}$ ) with carefully up to 0.0001 g level. The sample is being filtered and dried at 80°C for 4 h and cooled in desiccators after it has been washed up twice with distilled water (Olguin *et al.*, 2001; Sabzi *et al.*, 2018).

#### *Bioaccumulation analysis*

To estimate the iron, copper and zinc metals, the samples have been separated from the solution by centrifugation MICRO 22R model manufactured by Hettich of Germany. One gram of the wet sample more accurately weigh scales Sartryvs 124S model. They were transferred to crucible and then in electric furnaces BATEC PC 21 model with ashing process has performed at 550 °C. The contents dissolved in 3 ml 1:1 nitric acid solution and then reach to 10 ml of distilled water in volumetric flask after cooling. The solutions of iron, copper and zinc ions have been prepared from standard stock solutions (1000 mg L<sup>-1</sup>) in the concentration range of 0.1 to 10 mg of iron, 0.1 to 15 mg per liter for copper and 0.1 to 1.5 mg per liter for zinc have analyzed by flame atomic absorption spectrometer PG-990 model. Results have been calculated from the values of three parallel measurements and were

expressed in mg kg<sup>-1</sup> dried alga (Forstner and Muller, 1974).

#### *EF calculation*

Data for the calculation EF gained from the results of measured metal content of 15 samples prepared as follows: Five samples were prepared in triplicates with four different the compositions of the metals (with a control sample).

Enrichment factor (EF) has calculated according to the following ratios (Szabolcs *et al.*, 2013):

$$EF = C_E / C_C$$

C<sub>E</sub> = Microelement concentration of dry alga grown in the media with enhanced metal content

C<sub>C</sub> = Microelement concentration of dry alga grown in the control media

#### *Biopigments estimation*

Chlorophyll content has determined by centrifugation for 10 minute at 4000 rpm. Chlorophyll A (Ch-A) was extracted by using 5 ml 90% acetone and placed the tube in dark for 24 hour. Samples have centrifuged after extraction at 5000 rpm for 15 minute and collect the supernatant. Read the absorbance at 630 nm, 645 nm, and 665nm against 90% acetone as blank by using UV/VIS spectrophotometer and concentration of Chl-a A was calculated using the below formula:

$$C = 11.6 A_{665} - 1.31 A_{645} - 0.14 A_{630}$$

The concentration of Chl-A in a given volume of culture can be determined by below formula:

$$\text{Chl-A} (\text{mg L}^{-1}) = \frac{C \times V_e}{V_c}$$

C = Value obtained from above equation

V<sub>e</sub> = Volume of extract (ml)

V<sub>c</sub> = Volume of culture (liters)

*Phycobiliproteins estimation*

Amount of 5 ml cyanobacterial cell suspension was taken and subjected to centrifugation at 4000 rpm for 10 minutes. Phycobiliproteins have extracted in 5 ml of phosphate at pH 6.7, 0.05 M by 3 times repeated freezing and thawing. Freeze thawed samples subjected to Centrifugation at 1000 rpm for 15 minutes. The final extract has been measured at 562 nm, 615 nm, and 652 nm against phosphate buffer as blank.

The concentration of phycocyanin (PC), allophycocyanin (APC), and phycoerythrin (PE) have calculated by using the formula (Bennett and Bogorad, 1971):

$$PC = \frac{A_{615} - 0.474(A_{652})}{5.34}$$

$$APC = \frac{A_{652} - 0.208(A_{615})}{5.09}$$

$$PE = \frac{A_{562} - 2.41(PC) - 0.849(APC)}{9.62}$$

The concentration of phycobiliprotein in a total volume of culture can be determined as follows:

$$\text{Phycobiliprotein (mg ml}^{-1}\text{)} = \frac{C \times V_e}{V_c}$$

C=Value of PC, APC and PE obtained from above equations

Ve=Volume of extract (ml)

Vc=Volume of culture (ml)

*Carotenoids estimation*

Harvested biomass has been homogenized in homogenizer with 5 ml, 90% acetone and centrifuged the sample at 5000 rpm for 15 minutes. The carotenoids in samples were determined spectrophotometrically at 450 nm by using the following calculation formula (Jensen, 1978):

$$C = \frac{A_{450} \times V \times f \times 10}{2500}$$

C=Total amount of Cart (mg ml<sup>-1</sup>)

V=Volume of extract (ml)

f=Dilution factor

*Statistical analysis*

Data analyzed statistically using one way analysis of variance (ANOVA) using SPSS version 18. Duncan's multiple range test was used to compare differences among treatment means at (p<0.05) level (Duncan, 1955).

**Results***Growth analysis*

The effect of different microelement contents in media on the growth of *S. platensis* was evaluated daily during 14 days by determination of dry weights (g L<sup>-1</sup>) (Fig. 1 and Table 2).




Figure 1: Changes in the optical density (O.D) of *Spirulina platensis* on Zarrouk's media as a control, and growth medias with enhanced trace element level.

Table 2: Dry weight and OD (Optical density) of *Spirulina platensis* cultivated in Zarrouk's medium as a control and growth medias with enhanced trace element level.

Parameter	Day	Control	Fe10fold	Cu10fold	Zn10fold
Dry weight (g L <sup>-1</sup> )	14	1.55±0.02 <sup>a</sup>	1.57±0.03 <sup>a</sup>	1.54±0.03 <sup>a</sup>	1.57±0.01 <sup>a</sup>
OD	1	0.196±0.009 <sup>a</sup>	0.026±0.005 <sup>a</sup>	0.018±0.005 <sup>a</sup>	0.045±0.025 <sup>a</sup>
OD	2	0.022±0.005 <sup>a</sup>	0.028±0.007 <sup>a</sup>	0.027±0.009 <sup>a</sup>	0.080±0.092 <sup>a</sup>
OD	3	0.036±0.011 <sup>ab</sup>	0.028±0.017 <sup>a</sup>	0.072±0.019 <sup>b</sup>	0.057±0.019 <sup>ab</sup>
OD	4	0.075±0.010 <sup>b</sup>	0.039±0.014 <sup>a</sup>	0.096±0.029 <sup>b</sup>	0.076±0.004 <sup>b</sup>
OD	5	0.094±0.011 <sup>b</sup>	0.049±0.013 <sup>a</sup>	0.121±0.032 <sup>b</sup>	0.093±0.010 <sup>b</sup>
OD	7	0.149±0.037 <sup>ab</sup>	0.080±0.019 <sup>b</sup>	0.182±0.053 <sup>b</sup>	0.093±0.036 <sup>a</sup>
OD	8	0.177±0.044 <sup>ab</sup>	0.096±0.026 <sup>a</sup>	0.236±0.075 <sup>b</sup>	0.111±0.057 <sup>a</sup>
OD	9	0.193±0.054 <sup>ab</sup>	0.113±0.045 <sup>a</sup>	0.273±0.104 <sup>b</sup>	0.110±0.063 <sup>a</sup>
OD	10	0.207±0.052 <sup>ab</sup>	0.150±0.043 <sup>a</sup>	0.327±0.113 <sup>b</sup>	0.100±0.058 <sup>a</sup>
OD	11	0.232±0.067 <sup>ab</sup>	0.143±0.083 <sup>a</sup>	0.366±0.132 <sup>b</sup>	0.107±0.071 <sup>a</sup>
OD	13	0.274±0.106 <sup>a</sup>	0.203±0.116 <sup>a</sup>	0.424±0.223 <sup>a</sup>	0.344±0.139 <sup>a</sup>
OD	14	0.296±0.115 <sup>ab</sup>	0.217±0.124 <sup>ab</sup>	0.498±0.234 <sup>b</sup>	0.132±0.089 <sup>a</sup>

Within rows, means followed by the different letters are significantly different at  $p \leq 0.05$  as determined by the Duncan's test.

Results indicated no significant differences in the dry weights for the different medium, although the values of dry weight in iron and zinc treatments were higher than the control treatment the (Table 2). Results showed that there are no significant differences in OD values between control and other treatments after 14<sup>th</sup> day of culture (Table 2). Similar to these results, the optical density increased by 13 days, but the growth of *S. platensis* started to decline after incubation in the medium

with ten times enhanced Zn<sup>+2</sup> concentration. The growth curves showed no lag phase for the Zarrouk, s and other medium. A rapid increase in OD values has observed in the media with ten times enhanced Cu<sup>+2</sup> concentration with maximum 0.498±0.234 mg L<sup>-1</sup> on the 14th day ( $p < 0.05$ ). In contrast, a gradual different rate of increase in the OD has measured at all concentration (Fig. 1).



*Bioaccumulation analysis*

The extent of bioaccumulation and EF of iron, copper and zinc by *Spirulina* sp. grown in media with diverse

microelement content were measured (Table 3).

**Table 3: Bioaccumulation and EF of iron, copper and zinc by *Spirulina* sp. grown in media with diverse microelement content after 14 days of incubation.**

Treatment	Bioaccumulation of ions by <i>Spirulina</i> ( mg kg <sup>-1</sup> )		EF
	Alga grown in normal medium	Alga grown in normal medium with 10 times enhanced microelement content	
Fe <sup>2+</sup>	668±25.53 <sup>a</sup>	4465±39.68 <sup>b</sup>	6.68
Cu <sup>2+</sup>	81.66±5.50 <sup>a</sup>	394±7.54 <sup>b</sup>	4.82
Zn <sup>2+</sup>	52±4.00 <sup>a</sup>	704.33±8.14 <sup>b</sup>	13.54


Within rows, means followed by the different letters are significantly different at  $p \leq 0.05$  as determined by the Duncan's test.

*Pigment contents*

*Chlorophyll-A and Carotenoids*

The maximum production of chlorophyll occurred in 7<sup>th</sup> days of incubation at all concentration of treatments with maximum chlorophyll-A values (1.21±0.30 mg L<sup>-1</sup>) in ten

times enhanced Cu concentration. Although the concentration of chlorophyll accumulation in *S. platensis* grown in different media with maximum values of chlorophyll contents in copper treatment by 14<sup>th</sup> days of culture (Fig. 2).



**Figure 2: Changes in the chlorophyll A content of *Spirulina platensis* on Zarrouk's media (control) and growth medias with enhanced trace element level ( $p \leq 0.05$ ).**

Fig. 3, shows that the highest total carotenoid concentration was recorded

in *S. platensis* treated with Cu<sup>2+</sup> (0.0042±0/00 mg L<sup>-1</sup>) after 14 day of

inoculation. There were no significant differences in concentration of

carotenoids between control and other treatments in all experimental days.





Figure 3: Changes in the carotenoid content of *Spirulina platensis* on Zarrouk's media (control) and growth medias with enhanced trace element level ( $p \leq 0.05$ ).

*Phycobiliproteins*


Fig. 4 (a, b and c), shows the concentration phycobiliproteins accumulation in Zarrouk's (control) and growth medias with enhanced trace element level. There was a significant

difference in phycobiliproteins content for the different medias as compared to control ( $p < 0.05$ ). It has observed that an increase of ions concentration caused reduction of phycobiliproteins of *S. platensis*.



Downloaded from jifro.ir at 8:50 +0430 on Saturday April 4th 2020





**Figure 4: (a,b,c). Changes in phycoerythrin, allophycocyanin and phycocyanin content of *Spirulina platensis* on Zarrouk's (control) and growth media with enhanced trace element level ( $p \leq 0.05$ ).**

### Discussion

Nutrition concentration is one of the key factors that controls growth and dry weight of microalgae (Vonshak and Richmond, 1988; Faintuch *et al.*, 1991). The results in adverse with previous published data have showed that lower biomass of *Spirulina* sp. was measured in case of the ten times enhanced Zn<sup>+2</sup> concentration (Szabolcs *et al.*, 2013). Biomass growth of *Spirulina* sp. is not so adversely affected by the increasing iron concentration of the media, even though slight biomass enhancement has

observed in the previous cases. This conclusion is in agreement with Dou *et al.* (2013) who has found that dry weight of microalgae decreased firstly and then up gradually with the increase of concentration of Fe<sup>3+</sup>, Zn<sup>2+</sup>, and EDTA. The increase of Zn<sup>2+</sup> concentration would high the efficiency of photosynthesis in microalgae (Yamasaki *et al.*, 2012).

Results of this experiment showed that media culture supplemented with zinc has declined after 13 days of incubation. This conclusion is in

agreement with Gok and Esra (2009) who suggested,  $Zn^{2+}$  is not a constituent of the enzyme but necessary for its synthesis. Bascik Remisiewicz *et al.* (2009) reported that, Zn promote growth rate, since it is a main metabolic requirement for microalgae where it acts as an important enzyme cofactor. Omar (2002) reported that, very low zinc concentration improved growth of *Scenedesmus obliquus* and *S. quadricauda* but low zinc concentrations (i.e. 1.5, 4.5 and 8.0  $\mu\text{g L}^{-1}$ ) inhibit growth of *S. quadricauda*. However, higher concentrations support toxicity (20.6-37.7% growth inhibition) as well as longer exposure period.

Copper is a micronutrient required by microalgae growth and plays an important role as an enzymatic cofactor and electron carrier in the photosynthetic and respiratory processes (Andrade *et al.*, 2004). This experiment in confirmed with Estevez *et al.* (2001) and Abd El Baky *et al.* (2012) showed that green microalgae with addition of iron concentration cause increases biomass, with lower considered limiting for algal growth. Our experiment with Soeprbowati and Hariyati (2014) findings showed that the growth of *Spirulina* at 1 mg  $\text{L}^{-1}$  concentration increased until 11 day. In adverse with Dou *et al.* (2013) results showed that *Spirulina* sp. treated with  $\text{Cu}^{2+}$  causes the decrease of this ion with little effects on the growth density. Rueter and Petersen (1987) and Kilulya *et al.* (2015) reported that iron promotes the growth of cyanobacteria in natural waters. Results of this experiment have been showed that photosynthesis of

microalgae depended on  $\text{Fe}^{3+}$ , which is an important part of nitrate and nitrite reductase. The possibility that excess iron ion could be responsible for the decrease in alga growth by inducing oxidative stress (Wells *et al.*, 1994; Wilhelm and Trick, 1994; Boyer and Brand, 1998; Davey and Geider, 2001; Estevez *et al.*, 2001).

Regarding to reduction OD in zinc treatment after 13<sup>th</sup> day of culture, its uptake in different growth stage is changed. The maximum biomass values occurred in experiment with ten times enhanced  $\text{Cu}^{+2}$  concentration with the largest slope value being  $0.498 \pm 0.234$  ( $p \leq 0.05$ ) which is in agreement with views of Mihova and Godjevargova (2000) and Pavasant *et al.* (2006) who reported low concentration of Cu and Zn even stimulate the growth and the activity of the metabolic processes. Sunda and Guillard (1976) reported that copper toxicity generally due to the presence of free copper ions in the water. At the same time, the result reflected that, Cu ion supports growth only within the lowest concentration (5  $\mu\text{g Cu L}^{-1}$ ) as shown in this study. Wong *et al.* (1979) and Mosleh and Mofeed (2014) explain that, presence of  $\text{Cu}^{+2}$  in the growth media by low concentration could enhance the peroxidase activity, which involved in IAA (Indole Acetic Acid) degradation, a hormone widely known by its ability for stimulating growth.

As agreed by this research, Szabolcs *et al.* (2013) reported that amount of iron, copper and zinc uptake and EF by *S. platensis* were higher than the control treatments. Similarly, Mane and Bhosle

(2012) reported the highest percent bioaccumulation by *Spirulina* sp. for Fe (98.93%), Cu (81.2%) and Zn (79%) respectively at 5 mg L<sup>-1</sup> initial metal concentration.

In similar tests with Szabolcs *et al.* (2013), reports, the difference uptake in iron, copper, zinc ions uptake and EF by the algae *Spirulina* sp. might be related to absorption capacity. The capability of metal uptake depends on several factors like the growing condition such as temperature and pH circumstances and the level of available nutrients and microelements, metal concentration, the amount of the alga in the solution (biomass) and the absorption capacity of microalgae (Lovley, 2000).

Concerning the influence of medium type on carotenoids concentrations, it has found that similar trend of total chlorophyll content profile. The highest total carotenoid concentration has been recorded in experiment with ten times enhanced Cu10 concentration media (0.0042±0.0004 mg L<sup>-1</sup>) after 14 days of inoculation. This may indicate a strong relation between both chlorophyll and carotenoids contents. Such correlation could be attributed to that the carotenoids protect chlorophyll molecules against photo destruction and oxidation by molecular oxygen (Krinsky, 1979). Similarly, Vonshak (1997) reported that there was a positive correlation between chlorophyll and carotenoids content of *S. platensis*. Collen *et al.* (2003) and Pinto *et al.* (2011) observed similar results in *Gracilaria tenuistipitata*

exposed to copper and cadmium, with the increase of lutein and b-carotene.

Our results demonstrated that phycobiliprotein levels, including APC, PC, and PE, decreased in *S. platensis* after iron, zinc and copper treatments. These molecules absorb solar energy, transferring it to the reaction center of photosystem II, where chlorophyll A is excited by the flow of electrons (Gantt, 1981). According to Xia *et al.* (2004), a high concentration of copper altered phycobilisome structure, and these changes resulted in a decline of absorbed light energy, thus inhibiting photosynthesis. We found a decrease in phycobiliprotein levels, similar to the findings of Xia *et al.* (2004), who studied the red macroalgae *Gracilaria lemaneiformis* cultivated with copper during 4 four days. This indicates that iron, zinc and copper strongly inhibited the accumulation of phycobiliproteins. Similarly, Gouveia *et al.*, (2013) reported the amounts of phycobiliproteins decreased in *Gracilaria domingensis* treated with lead and copper. In addition, it could be due to its peripheral position in phycobilisomes on the thylakoid membrane (Gantt, 1981) and attributable to its sensitivity to metals (Kiran and Thanasekaran, 2011).

In the results from the measurement, growth of *S. platensis* has the highest uptake ability and high metals accumulation in their cells. By conducting this research, it was determined that *S. platensis* is the suitable species for enrichment. More studies in this field with regard to its high usage need, by its industrial

culture in Iran it could be necessary in different industries especially in food industry.

### Acknowledgements

The authors thank from Islamic Azad University, Branch of Ahwaz, Iran for supporting of this research.

### References

- Abd El Baky, H.H., El-Baroty, G.S., Bouaid, A., Martinez, M. and Aracil, J., 2012.** Enhancement of lipid accumulation in *Scenedesmus obliquus* by optimizing CO<sub>2</sub> and Fe<sup>3+</sup> levels for biodiesel production. *Bioresource Technology*, 119, 429–432.
- Andrade, L.R., Farina, M. and Filho, A.M.G., 2004.** Effects of copper on *Enteromorpha flexuosa* (Chlorophyta) *in vitro*. *Ecotoxicology and Environmental Safety Journal*, 58, 117- 125.
- Ansarifard, F., Rajabi Islami, H., Shamsaie Mehrjan, M. and Soltani, M., 2018.** Effects of *Arthrospira platensis* on growth, skin color and digestive enzymes of Koi, *Cyprinus carpio*. *Iranian Journal of Fisheries Sciences*, 17(2), 381-393.
- Arunakumara, K., Zhang, X. and Song, X., 2008.** Bioaccumulation of Pb<sup>2+</sup> and its effects on growth, morphology and pigment contents of *Spirulina (Arthrospira) platensis*. *Journal of Ocean University of China (Oceanic and Coastal Sea Research)*, 7(4), 397-403.
- Balaji, S., Kalaivani, T. and Rajasekaran, C., 2014.** Biosorption of zinc and nickel and its effect on growth of different *Spirulina* strains. *Journal of Clean Soil Air Water*, 42(4), 507-512.
- Bascik-Remisiewicz, A., Tomaszewska, E., Labuda, K. and Tukaj, Z., 2009.** The effect of Zn and Mn on the toxicity of Cd to the green microalga *Desmodesmus armatus* cultured at ambient and elevated (2%) CO<sub>2</sub> concentrations. *Polish Journal of Environmental Studies*, 18(5), 775– 780.
- Belay, A., Ota, Y., Miyakawa, K. and Shimamatsu, H., 1993.** Current knowledge on potential health benefits of *Spirulina*. *Journal of Applied Phycology*, 5, 235–241.
- Bennett, A. and Bogorad, L., 1971.** Properties of subunits and aggregates of bluegreen algal biliproteins. *Biochemistry*, 10, 3625-3634.
- Boyer, G.L. and Brand, L.E., 1998.** “Trace elements and harmful algal blooms,” in *Physiological Ecology of Harmful Algal Blooms*, D.M. Anderson, A.D. Cembella, and G.M. Hallegraeff, Eds., Springer, Berlin, Germany. pp. 489–508
- Bruland, K.W., Donat, J.R. and Hutchins, D.A., 1991.** Interactive influences of bioactive trace metals on biological production in oceanic waters. *Limnology and Oceanography*, 36, 1555–1577.
- Collen, J., Pinto, E., Pedersen, M. and Colepicolo, P., 2003.** Induction of oxidative stress in the red macroalgae *Gracilaria tenuistipitata* by pollutant metals. *Archives of Environmental Contamination and Toxicology*, 45, 337–342.

- Davey, M. and Geider, R.J., 2001.** Impact of iron limitation on the photosynthetic apparatus of the diatom *Chaetoceros muelleri* (Bacillariophyceae). *Journal of Phycology*, 37, 987–1000.
- Deniz, F., Saygideger, S.D. and Karaman, S., 2011.** Response to copper and sodium chloride excess in *Spirulina* sp. (Cyanobacteria). *Bulletin of Environmental Contamination and Toxicology*, 87, 11–15.
- Dou, X., Xiang-Hong, L., Mei-Zhen, L., Li-Sheng, Y., Rong, X. and Jian-Bing, J., 2013.** The effects of trace elements on the lipid productivity and fatty acid composition of *Nannochloropsis oculata*. *Journal of Renewable Energy*, 2013, 1-6.
- Duncan, D.B., 1955.** Multiple-range and multiple F tests. *Biometrics*, 11, 1–42.
- Estevez, M.S., Malanga, G. and Puntarulo, S., 2001.** Iron-dependent oxidative stress in *Chlorella vulgaris*. *Plant Science*, 161, 9–17.
- Faintuch, B.L., Sato, S. and Aquarone, E., 1991.** Influence of the nutritional sources on the growth rate of cyanobacteria. *Archives of Biology and Technology*, 34, 13–30.
- Forstner, U. and Muller, G., 1974.** Heavy metal in river and sea. Springer-Verlag. pp. 11-45.
- Gantt, E., 1981.** Phycobilisomes. *Annual Review of Plant Physiology*, 32, 327–347.
- Ghaeni, M., Matinfar, A., Soltani, M., Rabbani, M. and Vosoughi, A., 2011.** Comparative effects of pure spirulina powder and other diets on larval growth and survival of green tiger shrimp, *Peneaus semisulcatus*. *Iranian Journal of Fisheries Sciences*, 10(2), 208-217.
- Gok, Z. and Esra, Y., 2009.** Heavy metal uptake kinetics of microalgae in aquatic media. Proceedings of the 4th Iasme / Wseas Int. Conference on Water Resources, Hydraulics and Hydrology (WHH'09). pp. 175-179.
- Gouveia, C., Kreusch, M., Schmidt, E.C., Felix, M.R., De, L., Osorio, L.K.P., Pereira, D.T., Santos, R.D., Ouriques, L.C., Martins, R.De.P., Latini, A., Ramlov, F., Carvalho, T.J.G. and Chow F., 2013.** The effects of lead and copper on the cellular architecture and metabolism of the red alga *Gracilaria Domingensis*. *Microscopy and Microanalysis*, 19, 513-524.
- Gupta, V.K., Arshi, R., Saini, V.K. and Neeraj, J., 2006.** Biosorption of copper (II) from aqueous solutions by *Spirogyra* species. *Journal of Colloid and Interface Science Elsevier*, 296(1), 1-382.
- Habib, M.A.B., Parvin, M., Huntington, T.C. and Hasan, M.R., 2008.** A review on culture, production and use of *Spirulina* as food for humans and feeds for domestic animals and fish. FAO Fisheries and Aquaculture Circular, No. (1034). FAO, Rome. 33 P.
- Hadizadeh, Z., Shamsaie Mehrgan, M. and Hosseini Shekarabi, S.P., 2019.** The potential use of stickwater from a kilka fishmeal plant in *Dunaliella salina* cultivation. *Environmental Science and*



- Pollution Research*, pp.1-11. DOI: 10.1007/s11356-019-06926-w
- Jensen, A., 1978.** Chlorophylls and carotenoids. In: Hellebust JA, Craigie JS (eds) Handbook of phycological methods, physiological and biochemical methods. Cambridge University Press, Cambridge. pp. 59-70.
- Kilulya, F.K., Mamba B.B. and Msagati, T.A., 2015.** Extraction procedures and GCxGC-TOFMS determination of fatty acids (FAs) in cyanobacteria cultures and the effect of growth media iron concentration variation on cellular FAs composition. *Journal of Analytical and Bioanalytical Techniques*, 6(2), 1-44.
- Kiran, B. and Thanasekaran, K., 2011.** Metal tolerance of an indigenous cyanobacterial strain, *Lyngbya putealis*. *International Biodeterioration and Biodegradation*, 65, 1128-1132.
- Krinsky, I., 1979.** Carotenoid protection against oxidation. *Pure and Applied Chemistry*, 51, 649-660.
- Lovley, DR., 2000.** Environmental microbe-metal interactions. ASM Press, Washington, D.C. pp. 3-30.
- Mane, P.C. and Bhosle, A.B., 2012.** Bioremoval of some metals by living algae *Spirogyra* sp. and *Spirullina* sp. from aqueous solution. *International Journal of Environmental Research*, 6(2), 571-576.
- Mihova, St. and Godjevargova, T., 2000.** Biosorption of heavy metals from aqueous solutions. *Journal International Research Publication*, 1, 34-42.
- Mosleh, Y.Y.I. and Mofeed, J., 2014.** Bio-chemical biomarkers in algae *Scenedesmus obliquus* exposed to heavy metals Cd, Cu and Zn. *Life Science Journal*, 11(10), 994-1004.
- OECD, 2011.** Guidelines for testing of chemicals proposal for updating guideline 201. Freshwater Alga and Cyanobacteria, Growth Inhibition Test.
- Okmen, G., Bozanta, B., Aysel, U. and Ceyhan, N., 2011.** Zinc effect on chlorophyll a, total carbohydrate, total protein contents and biomass of Cyanobacterial species. *Journal of Applied Behavioral Science (JABS)*, 5(2), 67-73.
- Olguin, E.J., Galicia, S., Angulo-Guerrero, O. and Hernandez, E., 2001.** The effect of low light flux and nitrogen deficiency on the chemical composition of *Spirulina* sp. (*Arthrospira*) grown on digested pig waste. *Bioresource Technology*, 77, 19-24.
- Omar, H.H., 2002.** Bioremoval of zinc ions by *Scenedesmus obliquus* and *Scenedesmus quadricauda* and its effect on growth and metabolism. *International Biodeterioration and Biodegradation*, 50, 95-100.
- Pavasant, P., Apiratikul, R., Sungkhum, V., Suthiparinyanont, P., Wattanachira, S. and Marhaba, T.F., 2006.** Biosorption of  $\text{Cu}^{2+}$ ,  $\text{Cd}^{2+}$ ,  $\text{Pb}^{2+}$  and  $\text{Zn}^{2+}$  using dried marine green macroalga *Caulerpa lentillifera*. *Bioresource Technology*, 97, 2321-2329.



- Pinto, E., Carvalho, A.P., Cardozo, K.H.M., Malcata, F.X., Anjos, F.M.D. and Colepicolo, P., 2011. Effects of heavy metals and light levels on the biosynthesis of carotenoids and fatty acids in the macroalgae *Gracilaria tenuistipitata* (var. *Liui Zhang and Xia*). *Brazilian Journal of Pharmacognosy*, 21, 349–354.
- Ramakrishnan, C.M., Haniffa, M.A., Manohar, M., Dhanaraj, M., Arokiaraj, A.J., Seetharaman S. and Arunsingh, S.V., 2008. Effects of probiotics and *Spirulina* on survival and growth of juvenile common carp (*Cyprinus carpio*). *Israeli Journal of Aquaculture-Bamidgeh*, 60(2), 128-133.
- Rastar, M., Hosseini Shekarabi, S.P., Shamsaie Mehrgan, M. and Sabz, S., 2018. Effects of iron and zinc concentrations on growth performance and biochemical composition of *Haematococcus pluvialis*: A comparison between nanoparticles and their corresponding metals bulks. *Journal of Algal Biomass Utilization*, 9(2), 59-67.
- Romera, E., Gonzales, F., Ballester, A., Blazquez, M.L. and Munoz, J.A., 2007. Comparative study of biosorption of heavy metals using different types of algae. *Bioresource Technology*, 98, 3344-3353.
- Rueter, J.G. and Petersen, R.R., 1987. Micronutrient effects on cyanobacterial growth and physiology. *New Zealand Journal of Marine and Freshwater Research*, 21, 435-445.
- Sabzi, S., Shamsaie Mergan, M., Rajabi Islami, H. and Hosseini Shekarabi, S.P., 2018. Changes in biochemical composition and fatty acid accumulation of *Nannochloropsis oculata* in response to different iron concentrations. *Journal of Biofuels*, DOI: 10.1080/17597269.2018.1489672
- Sanchez, M., Bernal-Castillo, J., Rozo, C. and Rodriguez, I., 2003. *Spirulina* (Arthrospira): An edible microorganism: A review. *Universitas Scientiarum Journal of the Faculty of Sciences Pontificia Universidad Javeriana*, 8(1), 7-24.
- Soeprbowati, T.R. and Hariyati, R., 2014. Phycoremediation of  $Pb^{+2}$ ,  $Cd^{+2}$ ,  $Cu^{+2}$ , and  $Cr^{+3}$  by *Spirulina platensis* (Gomont) Geitler. *American Journal of BioScience*, 2(4), 165-170.
- Sunda, W.G. and Guillard, R.R., 1976. Relationship between cupric ion activity and the toxicity of copper to phytoplankton. *Journal of Marine and Freshwater Research*, 34, 511-529.
- Szabolcs, M., Kiss, A., Virag, D. and Forgo, P., 2013. Comparative Studies on accumulation of selected microelements by *Spirulina Platensis* and *Chlorella Vulgaris* with the prospects of functional food development. *Journal of Chemical Engineering and Process Technology*, 4(7), 1-6.
- Vonshak, A. and Richmond, A., 1988. Mass production of blue-green alga *Spirulina* – an overview. *Biomass*, 15, 233–247.

- Vonshak, A., 1997.** *Spirulina platensis* (Arthrospira), physiology, cell biology and biotechnology. Taylor and Francis Ltd, London. pp. 43-65.
- Wells, M.L., Price, N.M. and Bruland, K.W., 1994.** Iron limitation and the *Cyanobacterium Synechococcus* in equatorial Pacific waters. *Limnology and Oceanography Journal*, 39, 1481–1486.
- Wilhelm, S.W. and Trick, C.G., 1994.** Iron-limited growth of cyanobacteria: multiple siderophore production is a common response. *Limnology and Oceanography Journal*, 39, 1979–1984.
- Wong, P.T.S., Burnison, G. and Chau, Y.K., 1979.** Cadmium toxicity to freshwater algae. *Bulletin of Environmental Contamination and Toxicology*, 23, 487-490.
- Xia, J.R., Li, Y.J., Lu, J. and Chen, B., 2004.** Effects of copper and cadmium on growth, photosynthesis, and pigment content in *Gracilaria lemaneiformis*. *Bulletin of Environmental Contamination and Toxicology*, 73, 979–986.
- Yamasaki, Y., Yokose, T. and Nishikawa, T., 2012.** Effects of alginate oligosaccharide mixtures on the growth and fatty acid composition of the green algae *Chlamydomonas reinhardtii*. *Journal of Bioscience and Bioengineering*, 113(1), 112–116.
- Zarrouk, C., 1966.** Contribution to the study of a cyanobacterium: Influence of various physical and chemical factors on growth and photosynthesis of *Spirulina maxima* (Setchell and Gardner) Geitler. PhD thesis, University of Paris, France. pp. 4-5.