Research Paper: Immediate Effects of Lumbosacral Belt on Postural Control During Challenging Tasks in Patients With Chronic Low Back Pain



Masoud Ghofrani¹, Gholamreza Olyaei^{2*}, Saeed Talebian², Hossein Bagheri², Kazem Malmir³, Shiva Mousavi⁴

1. PhD Candidate, Department of Physical Therapy, School of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran.

2. Full Professor, Department of Physical Therapy, School of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran.

3. Assistant Professor, Department of Physical Therapy, School of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran.

4. Educational Instructor, Department of Physical Therapy, School of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran.



doj*: ???

Citation: Ghofrani M, Olyaei Gh, Talebian S, Bagheri H, Malmir K, Mousavi Sh. Immediate Effects of Lumbosacral Belt on Postural Control During Challenging Tasks in Patients With Chronic Low Back Pain. Journal of Modern Rehabilitation. 2017; 11(2):119-126.

Article info:

Received: 12 Oct. 2016 Accepted: 23 Jan. 2017

ABSTRACT

Introduction: Chronic Low Back Pain (CLBP) may be associated with impaired proprioception which can result in joint instability and balance deficit. However, wearing a lumbosacral belt may be helpful in this situation. This study aimed to identify postural control impairments in patients with CLBP and determine the effect of lumbosacral support on postural control strategies in these patients.

Materials and Methods: A total of 16 patients with CLBP and 16 healthy subjects participated in this study. Center of Pressure (COP) data were recorded for 30 seconds while wearing or not wearing a belt in four standing positions; double leg with open eyes, double leg with closed eyes, single leg with open eyes, single leg with closed eyes. Postural control was assessed using range, area and total mean velocity for each experimental condition.

Results: Patients with CLBP showed significantly larger mean COP range and mean area compared to the healthy controls in single leg stance with closed eyes (P<0.05). Lumbosacral belt reduced significantly mean COP range, mean COP area and mean total velocity during challenging conditions (P<0.05).

Keywords:

Chronic Low Back Pain, Postural control. Lumbosacral belt

Conclusion: Individuals with CLBP had poorer postural control compared to the healthy controls as determined by changes in COP. Lumbosacral belt can improve postural control during challenging tasks.

1. Introduction

ow back pain (LBP) is one of the most common health problems. The prevalence rate of this disorder is 15% among adults and ranges from 60% to 80% over the course of an entire life time [1, 2]. Postural control system operates on the basis of integrated information from three independent sensory sources; somatosensory, vestibular, and visual [3]. Thus, derangement in any of these sensory systems will influence the overall output of

.....

* Corresponding Author:

Gholamreza Olyaei, PhD Address: Department of Physical Therapy, School of Rehabilitation, Tehran University of Medical Sciences, Tehran, Iran. Tel: +98 (21) 77528469 E-mail: olyaeigh@sina.tums.ac.ir

the postural control system. Several studies have shown impaired proprioception in these patients [4, 5].

Control of posture, therefore, may be affected in patients with Chronic Low Back Pain (CLBP) [6, 7]. In this regard, wearing a lumbar belt may be beneficial. Lumbar belt may increase compression forces on the skin, thereby providing extra sensory afferents to the central nervous system by the cutaneous mechanoreceptors, which in turn would improve lumbar proprioception [8-10]. This situation may compensate for proprioceptive impairment in CLBP patients and restore postural balance for them. Assessment of human postural or balance control is of high interest to researchers and clinicians. In this regard, many measures have been proposed from force platform signals to quantifying postural steadiness. However, Center of Pressure (COP), defined as the point of application of the ground reaction forces under the feet, is found the most commonly used variable [11].

Postural control can be assessed using COP variables such as COP range in the Anterior-Posterior (AP) or Medial-Lateral (ML) direction, sway area and mean velocity. To the authors' knowledge, no article has yet investigated the effect of lumbosacral belt on balance parameters in CLBP patients. Therefore, the main purpose of the present investigation was to determine whether a commonly used lumbosacral belt can improve COP range in the AP or ML direction, sway area, and mean velocity in CLBP patients. The next goal was to identify postural control impairments in patients with CLBP and compare them during four balance tasks related to different activities of daily living.

2. Materials and Methods

Participants

Sixteen patients with CLBP participated in this study. Samples were recruited in an unpredictable way through advertising in physiotherapy departments of Rehabilitation Schools in Tehran.

Inclusion criteria were as follows: having localized back pain, feeling pain intensity less than 4 based on visual analog scale, lasting pain more than 6 months and radiating no further than the buttock, and lacking previous history of sciatica or other radicular involvement. The patients had neither history of vestibular and neurological disease nor of hip, knee, ankle or foot problems. Sixteen matched healthy subjects as control group were also participated in the study. They had neither low back pain 6 month prior to testing, nor evidence of vestibular, postural or musculo-skeletal impairments. The demographic characteristics of the participants are presented in Table 1. Each participant signed an informed consent form approved by the Institutional Review Board of the Tehran University of Medical Sciences (IR.TUMS. REC.13942145).

Instruments

A force platform (Bertec Corporation, Columbus, OH, USA) with sampling frequency of 500 Hz was used to record the COP data. Data were stored on a Pentium-based PC and then fed to MATLAB software for calculation of COP parameters. AP and ML displacement of COP were measured along the y-axis and x-axis, respectively. An extensible lumbosacral belt with posterior metal bar (Sacro Lumbar Support 2164 Oppo USA www.reidko) was used to assess the effects of lumbosacral belt on postural control (Figure 1).

Testing procedures

After familiarization with the tests, all participants completed balance tests during four conditions barefoot on a force platform. The arms were parallel to the trunk. A physiotherapist blinded to experimental objectives performed all testing procedures. The balance conditions were

Table 1. Mean(SD) of the demographic characteristics of the participants (n=16 in each group)

Voriakla	Group				
variable	CLBP, Mean (SD)	Control, Mean (SD)			
Age (y)	24.8(3.8)	23.2(2.2)			
Mass (kg)	64.8(12.0)	68.7(10.9)			
Height (m)	1.66(7.6)	1.7(9.1)			
BMI (kg/m ²)	23.3(2.9)	23.5(2.9)			
BMI: Body Mass Index		JMR			



Figure 1: Lumbosacral belt

JMR

performed randomly with and without lumbosacral belt as follows: 1. Two-legged stance with eyes open; 2. Twolegged stance with eyes closed; 3. One-legged stance with eyes open; and 4. One-legged stance with eyes closed.

The position of the feet was standardized using a tape marker on the force platform. Three trials, with a rest period of approximately 2 minutes, were performed for each condition, and each trial lasted for 30 seconds.

Data analyses

The raw data were filtered with a sixth order Butterworth, zero-phase low-pass filter at 10 Hz and converted into COP data using MATLAB-based routines (Mathworks, Natick, MA, USA). Parameters calculated from COP data were as follows: AP displacement, ML displacement, area and total mean velocity. The formulae used to calculate each parameter are presented in Table 2 [12].

Statistical analyses

SPSS (version 20.0 for Windows, Inc., Chicago, IL) was used to analyze all the data, with significance level set at P \leq 0.05. After passing normality test of Kolmogorov-

Table 2. Formulae used to	calculate C	OP parameters
---------------------------	-------------	---------------

Smirnov, a 2(LBP vs. control)×4(conditions)×2(with belt vs. without belt) repeated-measures multivariate analysis of variance (MANOVA) of 2 groups was used to assess postural control during different conditions between groups. Independent t test was used to analyze balance control between groups. Paired t test and effect size were further used to determine lumbosacral belt effectiveness. Cohen's d was used to describe the effect size: 0.2 is considered a "low" effect size, 0.5 an "average" effect size and 0.8 a "strong" one [13].

3. Results

Table 3 presents comparisons between groups in four balance conditions. Participants with CLBP showed significantly larger mean COP AP displacement and area in tasks 1 and 4, larger mean COP ML displacement in task 4 and higher mean of COP total mean velocity in task 1 compared to those in controls (P<0.05). The CLBP group reported 6.21 cm, 5.65 cm and 15.10 cm² for the means of the AP, ML displacement and area, respectively, compared to the healthy controls with 5.03 cm, 4.20 cm and 7.86 cm² of increased COP sway in task 4 (with strong effect size d=0.68 for AP displacement, d=1.19 for ML displacement and d=1.24 for area) suggesting poor postural control balance in CLBP group in challenging tasks.

Table 4 presents the effect of lumbosacral belt in four balance tasks. Conditions with lumbosacral belt presented significantly smaller COP displacement in AP direction in task 3, significantly smaller COP displacements, area and lower COP mean velocity in task 4 compared to conditions without lumbosacral belt (P<0.05). For the mean of the variables, the condition with lumbosacral belt reported AP displacement 3.05 cm in task 3, AP displacement 4.97 cm, ML displacement 4.18 cm, area 9.20 cm² and mean

Parameter	Formula
Anterior-posterior displacement (cm)	/ y _{max} -y _{min} /
Medial-lateral displacement (cm)	x _{max} -x _{min}
Area (cm²)	$A=2\pi F_{0.05[2, n-2]} (\sigma_{x}^{2} \sigma_{y}^{2} - \sigma_{xy}^{2})$ Where $\sigma_{xy} = \sum \frac{(x_{i}^{-}x)(y_{i}^{-}y)}{n}$
Mean velocity (cm/s)	$\bar{\upsilon} = 1/T \sum_{1}^{T} \left[(X_{t+1} - X_t)^2 + (y_{t+1} + y_t)^2 \right]$

JMR

	CLBP	Independent T-Test				MANOVA			
Variables		Balance Tasks				Group	Task		
		1	2	3	4	P-Value	P-Value	interaction	
Range AP (cm)	No	1.98(0.81)	1.90(0.87)	3.23(0.94)	5.03(1.75)	0.003*	0.000*	0.040*	
	Yes	2.56(1.26)	1.75(0.79)	3.59(0.90)	6.21(2.32)				
	P between	0.038*	0.480	0.134	0.030*				
	Effect size (d)	0.71	_	_	0.68				
Rang ML (cm)	No	1.23(1.28)	1.08(0.90)	2.42(0.80)	4.20(1.21)	0.010*	0.000*	0.055*	
	Yes	1.49(0.84)	1.14(0.97)	2.71(0.64)	5.65(3.53)				
	P between	0.370	0.780	0.132	0.038*				
	Effect size (d)	_	-	_	1.19				
	No	1.11(0.93)	1.02(1.20)	3.05(1.97)	7.86(5.82)	0.008*	0.000*	0.009*	
Area (cm ²)	Yes	2.02(2.05)	1.01(1.30)	4.00(2.16)	15.10(17.05)				
	P between	0.030*	0.970	0.080	0.032*				
	Effect size (d)	0.97	-		1.24				
Mean velocity (cm/s)	No	0.55(0.17)	0.67(0.59)	1.51(0.60)	3.02(0.67)	0.277			
	Yes	0.65(0.20)	0.58(0.22)	1.58(0.35)	3.32(0.88)				
	P between	0.038*	0.422	0.580	0.316		0.000*	0.455	
	Effect size (d)	0.58	_	_	_				

Table 3. Mean(SD) of balance parameters of individuals with and without CLBP during four balance tasks (MANOVA, independent t test and effect size)

JMR

(1) Two-legged stance with eyes open; (2) Two-legged stance with eyes closed; (3) One-leg stance with eyes open; and (4) one-leg stance with eyes closed

Range AP: COP displacement in anterior-posterior direction; Range ML: COP displacement in medial-lateral direction. *Significant differences between groups (P<0.05; with vs. without CLBP)

velocity 2.88 cm/s in task 4 compared to conditions without lumbosacral belt with AP displacement 3.78 cm in task 3, AP displacement 6.26 cm, ML displacement 5.65 cm, area 13.76 cm² and mean velocity 3.37 in task 4 of decreased COP sway in task 3 and 4 (with medium effect size d=0.70 for AP displacement in task 3 and d=0.55 for displacements in directions, mean velocity and d=0.47 for area in task 4) suggesting better postural control in condition with lumbosacral belt in challenging tasks.

4. Discussion

Our study compared individuals with and without CLBP under four balance tasks. Our results supported

poorer postural control in CLBP group. Also more challenging balance tasks like "one-legged stance with closed eyes" can detect more tactfully CLBP effects on balance. In addition, the effect of lumbosacral belt on postural control was assessed under four balance tasks. It was found that wearing lumbosacral belt could improve postural control, especially in challenging conditions.

Between group comparisons

The present study showed significant differences between groups, in that CLBP group reported poorer postural control compared to the healthy controls. Apparently, the differences between CLBP and healthy

	Belt	Paired T-Test				MANOVA		
Variables		Balance Task				Belt P-	Task P-	
		1	2	3	4	Value	Value	Interaction
Range AP (cm)	No	2.20(1.41)	1.82(0.72)	3.78(0.90)	6.26(2.36)	0.005*	0.000*	0.007*
	Yes	2.35(0.24)	1.83(0.93)	3.05(0.82)	4.97(1.65)			
	P between	0.595	0.936	0.001*	0.005*			
	Effect size (d)	_	_	0.70	0.55			
Rang ML (cm)	No	1.32(1.38)	1.01(0.83)	2.67(0.62)	5.67(3.37)	0.050*	0.000*	0.009*
	Yes	1.43(0.71)	1.22(1.02)	2.47(0.83)	4.18(1.59)			
	P between	0.66	0.362	0.206	0.005*			
	Effect size (d)	_	_	_	0.55			
	No	1.30(1.38)	0.91(1.16)	3.80(2.35)	13.76(13.81)	0.195	0.000*	0.124
Area (cm²)	Yes	1.83(2.06)	1.11(1.33)	3.26(1.82)	9.20(12.25)			
	P between	0.205	0.505	0.346	0.013*			
	Effect size (d)	_	_	_	0.47			
Mean velocity (cm/s)	No	0.56(0.18)	0.62(0.57)	1.64(0.42)	3.37(0.71)	0.023* 0.0	0.000*	0.011*
	Yes	0.65(0.21)	0.63(0.27)	1.45(0.54)	2.88(0.78)			
	P between	0.089	0.971	0.101	0.005*		0.000*	
	Effect size (d)	-	-	-	0.55			

Table 4. Mean (standard deviation) of balance parameters of two conditions without belt and with belt during four balance tasks in CLBP (MANOVA, paired t test and effect size)

JMR

(1) Two-legged stance with eyes open; (2) Two-legged stance with eyes closed; (3) One-leg stance with eyes open; and (4) one-leg stance with eyes closed

Range AP: COP displacement in anterior-posterior direction; Range ML: COP displacement in medial-lateral direction *Significant differences between conditions (P<0.05; with vs. without Lumbosacral belt)

individuals were task dependent and these differences are seen in more challenging activities. Likewise, Da Silva et al. [14] compared the balance of individuals with and without CLBP in five tasks. They concluded individuals with CLBP would present poorer postural control determined by center of pressure measurements compared to the healthy controls, mainly during more challenging balance tasks like semi-tandem and one-legged stance conditions.

In a systematic review of the literature, Ruhe et al. [14] determined differences in COP between LBP patients and control group. The patients with LBP showed a greater postural instability than the control group. This difference was noticed in 88% of the studies. Only two studies reported significantly lower COP excursions in patients suffering from LBP [15, 16]. Compared to the control group, participants with LBP exhibited a greater sway area [17, 18], increased COP mean displacement [19-23], and finally higher COP sway velocity [24-28]. Mazaheri et al. [29] reviewed postural control studies which manipulating sensory systems (visual, vestibular, proprioception). The study showed controversial results. Participants with CLBP sometimes had less, sometimes more, and sometimes the same COP sway as participants without LBP in double leg standing.

Effect of wearing a lumbosacral belt

The present study showed a significant reduction in the COP measurements after wearing lumbosacral belt during challenging tasks. Previous studies showed that lumbosacral orthoses for patients with LBP can decrease the pain and help improve balance ability by stabilizing the lumbar region [30, 31]. Redford et al. [32] reported that lumbosacral orthoses would limit movement of the trunk and decrease the load on the lumbar region by transmitting forces applied to intervertebral disks to soft tissues surrounding the abdomen which ultimately result in pain relief. The result of Vogt et al. [33] study revealed that lumbosacral orthoses might increase joint position sense as a result of increasing afferent proprioceptive inputs through mechanoreceptors of the skin, facilitate voluntary extension of the spine and improve postural control.

In another study, Sinaki et al. [34] reported that wearing kypho-orthosis could heighten location awareness of the vertebral joint or proprioception and improve balance and walking quality in over 60 years old subjects with risk of falls. Balancing in a postural task is a dynamic response, consisting of the body (including the spine) and the Central Nervous System (CNS) as the controller. Apparently the CNS tunes stiffness to optimize system's performance [35, 36]. It is possible that the CNS senses the added information from the lumbosacral orthoses and modulates postural balance easier. This scenario is consistent with our results, where a decrease in COP measures occurred by lumbosacral orthoses in challenging tasks.

As a conclusion, this study and literature demonstrate that wearing lumbosacral may improve postural control by manipulating sensory inputs from lumbosacral region. These results will help prepare baseline data providing related information with proper mediation about wearing lumbosacral orthoses rightly. A limitation of this study was the duration that the subjects wore the orthoses which was just a few seconds. Further studies are needed to evaluate utility of orthoses prescribed as a therapy method for patients with LBP in longer periods. There are different types of orthoses including soft and hard ones. It is suggested to investigate application of different lumbosacral orthoses in patients with CLBP and their effects on postural control and low back pain.

Acknowledgements

We appreciate the Postgraduate Studies and Research Program, Tehran University of Medical Sciences, for providing the generous grant to carry out this research. We also acknowledge the generous assistance of the staff and students of the Rehabilitation school, Tehran University of Medical Sciences.

Conflict of Interest

The authors declared no conflicts of interest.

References

- [1] Panjabi MM. Clinical spinal instability and low back pain. Journal of Electromyography and Kinesiology. 2003; 13(4):371–9. doi: 10.1016/s1050-6411(03)00044-0
- [2] Walker BF. The Prevalence of Low Back Pain: A Systematic Review of the Literature from 1966 to 1998. Journal of Spinal Disorders. 2000; 13(3):205–17. doi: 10.1097/00002517-200006000-00003
- [3] Massion J. Movement, posture and equilibrium: Interaction and coordination. Progress in Neurobiology. 1992; 38(1):35– 56. doi: 10.1016/0301-0082(92)90034-c
- [4] Newcomer KL, Laskowski ER, Yu B, Johnson JC, An K-N. Differences in Repositioning Error Among Patients With Low Back Pain Compared With Control Subjects. Spine. 2000; 25(19):2488–93. doi: 10.1097/00007632-200010010-00011
- [5] Boucher JA, Abboud J, Nougarou F, Normand MC, Descarreaux M. The Effects of Vibration and Muscle Fatigue on Trunk Sensorimotor Control in Low Back Pain Patients. Hug F, editor. Plos One. 2015; 10(8):e0135838. doi: 10.1371/journal.pone.0135838
- [6] Brumagne S, Janssens L, Knapen S, Claeys K, Suuden-Johanson E. Persons with recurrent low back pain exhibit a rigid postural control strategy. European Spine Journal. 2008; 17(9):1177–84. doi: 10.1007/s00586-008-0709-7
- [7] da Silva RA, Vieira ER, Fernandes KBP, Andraus RA, Oliveira MR, Sturion LA, Calderon MG. People with chronic low back pain have poorer balance than controls in challenging tasks. Disability and Rehabilitation 2017; doi: 10.1080/09638288.2017.1294627.
- [8] Kang JI, Kwon HM, Jeong DK, Choi H, Moon YJ, Park JS. The effects on postural control and low back pain according to the types of orthoses in chronic low back pain patients. Journal of Physical Therapy Science. 2016; 28(11):3074–7. doi: 10.1589/jpts.28.3074
- [9] Newcomer K, Laskowski ER, Yu B, Johnson JC, An K-N. The effects of a lumbar support on repositioning error in subjects with low back pain. Archives of Physical Medicine and Rehabilitation. 2001; 82(7):906–10. doi: 10.1053/apmr.2001.23281
- [10] Sinaki M, Brey RH, Hughes CA, Larson DR, Kaufman KR. Significant reduction in risk of falls and back pain in osteoporotic-kyphotic women through a Spinal Proprioceptive Extension Exercise Dynamic (SPEED) program. InMayo Clinic Proceedings. 2005; 80(7):849-55. doi: 10.4065/80.7.849

- [11] Winter DA. Biomechanics and motor control of human movement. 4th edition. Hoboken, New Jersey: John Wiley & Sons; 2009.
- [12] Salavati M, Hadian MR, Mazaheri M, Negahban H, Ebrahimi I, Talebian S, et al. Test-retest reliabty of center of pressure measures of postural stability during quiet standing in a group with musculoskeletal disorders consisting of low back pain, anterior cruciate ligament injury and functional ankle instability. Gait & Posture. 2009; 29(3):460–4. doi: 10.1016/j. gaitpost.2008.11.016
- [13] Lakens D. Calculating and reporting effect sizes to facilitate cumulative science: a practical primer for t-tests and ANO-VAs. Frontiers in Psychology. Frontiers Media SA; 2013; 4. doi: 10.3389/fpsyg.2013.00863
- [14] da Silva RA, Vieira ER, Fernandes KBP, Andraus RA, Oliveira MR, Sturion LA, et al. People with chronic low back pain have poorer balance than controls in challenging tasks. Disability and Rehabilitation. 2017; 1–7. doi: 10.1080/09638288.2017.1294627
- [15] Mok NW, Brauer SG, Hodges PW. Hip Strategy for Balance Control in Quiet Standing Is Reduced in People With Low Back Pain. Spine. 2004; 29(6):E107–E112. doi: 10.1097/01. brs.0000115134.97854.c9
- [16] Salavati M, Mazaheri M, Negahban H, Ebrahimi I, Jafari AH, Kazemnejad A, et al. Effect of Dual-Tasking on Postural Control in Subjects With Nonspecific Low Back Pain. Spine. 2009; 34(13):1415–21. doi: 10.1097/brs.0b013e3181a3a917
- [17] Lafond D, Champagne A, Descarreaux M, Dubois J-D, Prado JM, Duarte M. Postural control during prolonged standing in persons with chronic low back pain. Gait & Posture. 2009; 29(3):421–7. doi: 10.1016/j.gaitpost.2008.10.064
- [18] Harringe ML, Halvorsen K, Renström P, Werner S. Postural control measured as the center of pressure excursion in young female gymnasts with low back pain or lower extremity injury. Gait & Posture. 2008; 28(1):38-45. doi: 10.1016/j. gaitpost.2007.09.011
- [19] Hamaoui A, Do MC, Poupard L, Bouisset S. Does respiration perturb body balance more in chronic low back pain subjects than in healthy subjects? Clinical Biomechanics. 2002; 17(7):548–50. doi: 10.1016/s0268-0033(02)00042-6
- [20] Grimstone SK, Hodges PW. Impaired postural compensation for respiration in people with recurrent low back pain. Experimental Brain Research. 2003; 151(2):218–24. doi: 10.1007/s00221-003-1433-5
- [21] Hamaoui A, Do M., Bouisset S. Postural sway increase in low back pain subjects is not related to reduced spine range of motion. Neuroscience Letters. 2004; 357(2):135–8. doi: 10.1016/j.neulet.2003.12.047
- [22] Smith M, Coppieters MW, Hodges PW. Effect of experimentally induced low back pain on postural sway with breathing. Experimental Brain Research. 2005; 166:109–117.
- [23] Popa T, Bonifazi M, Della Volpe R, Rossi A, Mazzocchio R. Adaptive changes in postural strategy selection in chronic low back pain. Experimental Brain Research. 2006; 177(3):411–8. doi: 10.1007/s00221-006-0683-4
- [24] Della Volpe R, Popa T, Ginanneschi F, Spidalieri R, Mazzocchio R, Rossi A. Changes in coordination of postural control during dynamic stance in chronic low back pain pa-

tients. Gait & Posture. 2006; 24(3):349–55. doi: 10.1016/j.gait-post.2005.10.009

- [25] Mann L, Kleinpaul JF, Pereira Moro AR, Mota CB, Carpes FP. Effect of low back pain on postural stability in younger women: Influence of visual deprivation. Journal of Bodywork and Movement Therapies. 2010; 14(4):361–6. doi: 10.1016/j. jbmt.2009.06.007
- [26] Luoto S, Taimela S, Hurri H, Aalto H, Pyykkö I, Alaranta H. Psychomotor Speed and Postural Control in Chronic Low Back Pain Patients. Spine. 1996; 21(22):2621–7. doi: 10.1097/00007632-199611150-00012
- [27] Lafond D, Champagne A, Descarreaux M, Dubois J-D, Prado JM, Duarte M. Postural control during prolonged standing in persons with chronic low back pain. Gait & Posture. 2009; 29(3):421–7. doi: 10.1016/j.gaitpost.2008.10.064
- [28] Harringe ML, Halvorsen K, Renström P, Werner S. Postural control measured as the center of pressure excursion in young female gymnasts with low back pain or lower extremity injury. Gait & Posture. 2008; 28(1):38–45. doi: 10.1016/j. gaitpost.2007.09.011
- [29] Mazaheri M, Coenen P, Parnianpour M, Kiers H, van Dieën JH. Low back pain and postural sway during quiet standing with and without sensory manipulation: A systematic review. Gait & Posture. 2013; 37(1):12–22. doi: 10.1016/j. gaitpost.2012.06.013
- [30] Reyna JR, Leggett SH, Kenney K, Holmes B, Mooney V. Lumbar Muscle. Spine. 1995; 20(1):68–73. doi: 10.1097/00007632-199501000-00013
- [31] Kim MH. A biomechanical effectiveness of corset and back brace for low back pain syndrome. Physical Therapy Korea. 1996; 28(11): 3074–3077. doi: 10.1589/jpts.28.3074
- [32] Redford JB, Basmajian JV, Trautman P. Orthotics: Clinical practice and rehabilitation technology. New York: Churchill Livingstone; 1995.
- [33] Vogt L, Pfeifer K, Portscher M, Banzer W. Lumbar corsets: their effect on three-dimensional kinematics of the pelvis. Journal of Rehabilitation Research & Development. 2000; 37(5):495-9. PMID: 11322147
- [34] Sinaki M, Brey RH, Hughes CA, Larson DR, Kaufman KR. Significant Reduction in Risk of Falls and Back Pain in Osteoporotic-Kyphotic Women Through a Spinal Proprioceptive Extension Exercise Dynamic (SPEED) Program. Mayo Clinic Proceedings. 2005; 80(7):849–55. doi: 10.4065/80.7.849
- [35] Reeves NP, Everding VQ, Cholewicki J, Morrisette DC. The effects of trunk stiffness on postural control during unstable seated balance. Experimental Brain Research. 2006; 174(4):694–700. doi: 10.1007/s00221-006-0516-5
- [36] Selen LPJ, Beek PJ, van Dieën JH. Impedance is modulated to meet accuracy demands during goal-directed arm movements. Experimental Brain Research. 2005; 172(1):129–38. doi: 10.1007/s00221-005-0320-7