

Effect of Milk Thistle (*Silybum marianum* L.) on Biochemical Parameters and Immunity of Broiler Chicks Fed Aflatoxin B1 after Three Weeks

Maliheh Amiri Dumari¹, Hadi Sarir^{*1}, Omid Fani Makki¹, Nazar Afzali¹

Received: 20.02.2014

Accepted: 30.03.2014

ABSTRACT

Background: This study was conducted to determine the efficacy of milk thistle seeds (MTSs) in counteracting the toxic effects of aflatoxin B1 (AFB₁) in a contaminated diet fed to broilers.

Methods: Two dietary inclusion rates of AFB₁ (0, 0.250 and 500 ppb) and MTS (0, 0.5 and 1%) were tested in a 3×3 factorial manner. The effect of nine experimental treatments was assessed using 216 one-d-old Ross 308 male broiler chicks in a randomized complete design with 4 replicates of 6 birds each from one to 21 days of age. The effects of dietary AFB₁ and MTS on serum biochemistry factors, antibody titer against Newcastle disease (ND) and influenza disease (ID) in broilers were evaluated at the end of this period.

Results: Statistical analysis of the main effects of diets indicated no significant changes in uric acid, cholesterol, triglycerides, low density lipoprotein (LDL), ID, and phosphorus compared to the control (P>0.01). Also, addition of 500 ppb of dietary AFB₁ into the diet was associated with significant decreases in serum glucose, calcium, high density lipoprotein (HDL), and ND compared to the control group (P<0.01). The contaminated diet significantly increased the activities of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) (P<0.05).

Conclusion: Milk thistle showed protective effects and resulted in some serum enzyme activities and serum biochemical changes associated with aflatoxin toxicity.

Keywords: Aflatoxin, Antibody Titer, Broiler, Lipid Profiles, Liver Enzymes, Milk Thistle.

IJT 2014; 1098-1103

INTRODUCTION

Mycotoxins are a group of structurally diverse secondary fungal metabolites that occur worldwide as contaminants of grain [1]. Various mycotoxins, like aflatoxin (AF), ochratoxin A (OA), zearalenone, T-2 toxin, vomitoxin, and fumoninsin, naturally contaminate foods and feed of poultry [1, 2]. Aflatoxins are a group of heterocyclic toxic metabolites of toxigenic fungi *Aspergillus flavus* and *A. parasiticus* [2]. Aflatoxicosis in poultry is a chronic illness characterized by high mortality, anorexia, decreased growth rates, negative feed conversions, fatty liver, decreased egg production, poor pigmentation, and increased susceptibility to some diseases [3]. In order to eliminate or reduce mycotoxin in animal diets, different physical, chemical, and biological methods are used [4]. The mechanisms of action of AFB₁ involve their metabolism to reactive intermediates, which bind to macromolecules with consequent disruption of transcriptional and

translational processes [5]. Milk thistle seed (MTS), a medicinal herb, has been extensively used in folk medicine for treating liver diseases [6]. Certain active ingredients found in the seed of this plant possess numerous medicinal properties. Earlier in 1960, a German scientist isolated a flavonoid 'silymarin' from milk thistle [6]. Chemically, it is composed of four flavonoids- silybin, isosilybin, silydianin, and silychristin [7]. Silybin is the major component constituting 50 to 70% of silymarin and exhibit greater biological activities [8]. Chakarverty and Parsad reported that milk thistle has hepatoprotective and hepatorestorative functions and protects liver and kidney from both exo- and endo-toxins [7]. It reduces liver enzyme production and shows anti-inflammatory and T-cell modulating effects [9]. Birds affected by AFB₁ can effectively be recovered if treated with milk thistle [10, 11]. Increase in AST concentration after challenge with aflatoxin has been reported [12, 13]. In broilers, higher levels of silymarin afford partial protection against a

1. Department of Animal Science, Birjand University, Birjand, Iran.

*Corresponding Author: E-mail: hsarir@birjand.ac.ir

35-day challenge with 0.8 ppm (body weight) aflatoxin [10]. Basaga *et al.* (1997) reported that milk thistle supports the immune system through its powerful antioxidant and free radical scavenging action. It has the ability to preserve the supply of glutathione and has direct effects on immune cells [14]. The main objective of this study was to investigate the effects of MTS on serum parameters, liver enzymes, lipid profiles, and immune performance of broiler chicks contaminated with AFB₁ against Newcastle and influenza diseases (ND and ID).

MATERIALS AND METHODS

Contaminating Poultry Feed

A known AFB₁ producing strain of *A. flavus* (PTCC NO: IR 111) was maintained on Potato Dextrose Agar (PDA) at optimum conditions in the Animal Science Institute, Birjand University, Iran. This culture was, then, introduced to commercial feed placed in a humid (15%) and hotter (27°C) regions to fully propagate the toxin in feed. The AFB₁ was semi-quantified according to Tapia [15] using thin layer chromatography (TLC). Commercial feed was contaminated weekly to maintain the toxin within the required limits. The AFB₁ in the rice powder consisted of 60 ppm AFB₁.

Bird Husbandry and Experimental Protocol

A total of 216 one-d-old Ross 308 male broiler chicks were randomly allocated to nine groups in a 3×3 factorial experiment. During this study, the birds were submitted to conventional broiler chicken management and housed in floor pens in an environmentally controlled broiler house with litter floors. The chicks were divided into nine treatment groups, with four replicates per treatment and 12 chicks per replicate: T1) basal diet (BD) free of toxin, T2) BD contaminated with 250 ppb AFB₁, T3) BD contaminated with 500 ppb AFB₁, T4) BD supplemented with 0.5% MTS, T5) BD supplemented with 0.5 % MTS plus 250 ppb AFB₁, T6) BD supplemented with 0.5% MTS plus 500 ppb AFB₁, T7) BD supplemented with 1% MTS, T8) BD supplemented with 1.0 % MTS plus 250 ppb AFB₁, T9) BD supplemented with 0.1 % MTS plus 500 ppb AFB₁. MTS was collected from the outskirts of Zabol district in Sistan and Blochestan province, Iran (Figure 1). Chickens consumed the diets and water *ad*

libitum. They received a commercial diet formulated to meet or exceed the nutritional requirements of broilers (for 21 day of age) as recommended by the National Research Council (NRC) (Table 1).

Table 1. Composition of the starter, grower and finisher diets fed to broilers (as fed).

Feed Stuffs	Starter Period (1-14 day)	Grower Period (15-21 day)
Maize	5443	50.42
Soybean meal(44%CP)	35	30.29
Wheat	-	10
Fish meal (60 % CP)	3.07	2.04
Soybean Fat	3.29	3.57
Dicalcium phosphate	1.73	1.47
Oyster shell	1.16	1.04
Mineral Premix	0.5	0.5
Vitamin premix	0.5	0.5
Salt	0.2	0.2
DL-methionine	0.35	0.28
L-lysine	0.24	0.19
Analyzed values		
ME (Kcal Per kg)	2980	3050
CP (Percent)	22	20
Lys (Percent)	1.43	1.24
Met + Cys (Percent)	1.07	0.95
Thr (Percent)	0.84	0.74
Ca (Percent)	1.05	0.90
P (Percent)	0.52	0.45



Figure 1. Milk thistle seed (MTS): Collected from the outskirts of Zabol district in Sistan balochestan province, Iran.

Serum Biochemical Parameters

Blood was collected in nonheparinized tubes from two broilers of each treatment by puncturing the brachial vein on the last day of

the period. Serum was separated after 8 to 10 hours through the standard procedures and was stored at -20°C for subsequent analysis. The individual and combined effects of dietary AFB₁ and MTS on serum biochemistry factors, lipid profile, and liver enzymes of broilers were evaluated. The individual serum samples were analyzed for uric acid, creatinine, cholesterol, triglycerides, glucose, LDL, HDL, iron, phosphorus, and calcium, and the activities of aspartate aminotransferase (AST) and alanine aminotransferase (ALT) were measured by the ultraviolet spectrometry method (Anonymous, 1984) - using an automatic analyzer (Boehringer Mannheim Hitachi 704 automatic analyzer, Japan). Then, the serum samples were assayed for Newcastle and Influenza vaccine titers using ELISA technique on the last day of the period (day 21).

Statistical Analysis

The data were subjected to ANOVA as a completely randomized design in the factorial procedure (Macros software, 2010). Differences were considered significant at $P < 0.05$. Also, the Tukey-Kramer test was adapted to compare mean values. All of the procedures used in testing the chickens in this experiment were

approved by the Department of Poultry Science at University of Birjand, Iran.

RESULTS

Data presented in Tables 2 and 3 demonstrate that AFB₁ and MTS had no significant effect on uric acid, iron, phosphorus, cholesterol, triglycerides, LDL, and influenza vaccine titer compared to the uncontaminated diet. In addition, MTS alone did not have any significant effects on uric acid, cholesterol, triglycerides, LDL, phosphorus, and influenza vaccine titer compared to the contaminated diet alone. As shown in Tables 2 and 3, addition of 500 ppb of AFB₁ into the diet resulted in a significant decrease in glucose (228.6 mole/L), HDL (65.1 mmole/L), calcium (7.82 mmole/L), and Newcastle vaccine titer (7.42) compared to the control ($P < 0.01$). Moreover, thistle-supplemented diet showed a significant increase in glucose level compared to the control groups ($P < 0.01$). Feeding diets contaminated with AF resulted in a significant increase in creatinine (1.99 $\mu\text{mole/L}$), AST (221.5 U/L), and ALT (31.49 U/L) ($P < 0.05$) compared to the control group. Moreover, the mean total of AST was significantly lower in the birds fed with MTS diets compared to the values exhibited by the chicks in the control group ($P < 0.05$).

Table 2. Effect of aflatoxin B₁ (AFB₁) and Milk thistle seeds (MTS) on serum biochemistry and some liver enzymes of broiler chickens at the end of period (21 days).

Groups		Glucose	Creatinine	Uric acid	Iron	Phosphorus	Calcium	AST	ALT
AFB ₁	MTS	(mole/L)	($\mu\text{mole/L}$)	(mmole/L)	(mole/L)	(mmole/L)	(mmole/L)	(U/L)	(U/L)
(ppb)	(%)								
0	-	242.1 ^a	0.56 ^b	5.24	396.9	7.14	9.31 ^a	198.5 ^b	21.12 ^b
250	-	245.3 ^a	1.61 ^a	6.05	359.8	6.41	8.93 ^{ab}	206.9 ^b	29.48 ^a
500	-	228.6 ^b	1.99 ^a	6.61	355.1	6.62	7.82 ^b	221.5 ^a	31.49 ^a
$\pm\text{SEM}$		2.89	0.13	0.27	2.99	0.15	0.19	2.7	1.64
-	0	218.8 ^b	1.54	6.49	248.1	6.31	8.22	220.8 ^a	28.91
-	0.5	246.3 ^a	1.35	5.58	254.1	6.84	9.12	204.1 ^b	27.75
-	1.0	251.1 ^a	1.27	5.84	289.4	6.92	8.91	202.1 ^b	25.41
$\pm\text{SEM}$		24.18	0.021	0.28	28.41	0.12	0.16	17.54	3.11
Probabilities									
AFB ₁		0.01	0.05	NS	NS	NS	0.01	0.05	0.05
MTS		0.01	NS	NS	NS	NS	NS	0.05	NS
AFB ₁ ×MTS		NS	NS	NS	NS	NS	NS	NS	NS

(a,b) Main effect means within a column lacking a common superscript differ significantly ($P < 0.05$) & ($P < 0.01$).

[Main effect values are mean \pm SEM from 6 observations each].

Ns: Not significant.

Table 3. Effect of aflatoxin B1 (AFB1) and Milk thistle seeds (SMS) on Lipid profiles and antibody titers at the end of period (21 days).

Groups		Lipid profiles				Antibody titers (Log2)	
AFB1 (ppb)	MTS (%)	Cholesterol (mole/L)	Triglycerides (mole/L)	LDL (mmole/L)	HDL (mmole/L)	Newcastle Disease	Influenza Disease
0	-	154.5	133.8	39.5	88.5 ^a	8.46 ^a	4.49
250	-	153.5	133.8	38.8	84.5 ^a	8.06 ^{ab}	4.34
500	-	156.1	144.9	29.9	65.1 ^b	7.24 ^b	4.29
±SEM		4.21	3.02	2.42	2.46		
-	0	157.2	143.7	36.9	74.6	8.36	3.66
-	0.5	152.7	135.3	41.4	81.4	8.46	3.69
-	1.0	145.1	133.5	30.3	82.1	8.49	3.89
±SEM		4.22	2.42	1.37	2.13	0.41	0.35
Probabilities							
AFB1		NS	NS	NS	0.01	0.01	NS
SMS		NS	NS	NS	NS	NS	NS
AFB1×SMS		NS	NS	NS	NS	NS	NS

-(a,b) Main effect means within a column lacking a common superscript differ significantly ($P<0.05$) & ($P<0.01$). [Main effect values are mean \pm SEM from 6 observations each].

Ns: Not significant.

DISCUSSION

AFB₁ is known to be hepatotoxin and causes genetic damage [16, 17]. In the present study, significant increases in the levels of AST and ALT were observed upon treatment with AFB₁ during 21 days. This increase in the values of serum AST and ALT might be due to hepatotoxic effects of AFB₁ which is in agreement with the findings of Fani Makki *et al.* [18]. Several studies have reported similar hepatotoxic property for AFB₁, which is in agreement with the findings of the present study [9-11]. Bares *et al.* (2008) reported the hepatoprotective property of silybin, which is the active ingredient of milk thistle [16]. In addition, the findings of the present study are in close agreement with those of Schrieber *et al.* (2008) who reported reduction of ALT and AST by milk thistle in human hepatitis C patients [17]. The findings of the present study are supported by others who reported reduction of AST by supplementation of feed with medicinal herbs as well [10, 13, 18].

Serum total glucose, HDL, and AST levels were significantly different between groups contaminated with different levels of AFB₁. Creatinine and ALT experienced significant increases and, in fact, were the values which were significantly affected in groups contaminated by AFB₁ alone, but not influenced by milk thistle consumption alone, whereas

calcium and HDL decreased after aflatoxin administration. However, treatments did not influence the serum concentrations of uric acid, cholesterol, triglycerides, LDL, phosphorus, and iron. Previous studies performed with high levels of AFB₁ (0 and 1.0 ppm) showed significant decreases in serum total glucose, protein, albumin, total cholesterol, and uric acid levels [19]. However, 1.0 ppm of AFB₁ did not affect uric acid, γ -GT, and phosphorus levels [19]. Rastogi *et al.* (2000) reported that silymarin reversed biochemical changes in liver and serum in AFB₁ intoxicated rats, indicating that it has a hepatoprotective action in preventing AFB₁ induced injury [20]. Lutensko *et al.* (2008) reported that silymarin phytosome caused no difference in ALT and AST serum activity, while total protein, calcium, phosphorus, albumin, and globulin levels increased compared with the control ($P<0.05$) [21].

Serum antibody titer against Newcastle and influenza diseases did not significantly change in treatments containing thistle seed, whereas lowest titers were recorded in groups AFB₁. It is clear from our observations and those of other researchers that aflatoxin causes severe immunosuppression that might be due to reduction in phagocytic activity of blood monocytes and depressed complement activity, hence, depressed opsonization and phagocytic activity [22]. Moreover, there are some reports on the immune-stimulatory effects of milk thistle

[23]. The results of the present study are supported by Tedesco *et al.* (2004) who reported the negative role of contaminated feed in the presence of immunosuppressant aflatoxin in diet [10]. In conclusion, the results of the present study showed that increases in the amount of liver enzymes occurred as a result of AFB₁. The fact that milk thistle supported liver function in conjunction with acute aflatoxin challenge is encouraging and warrants further research.

ACKNOWLEDGEMENTS

This study was conducted at University of Birjand and was supported by the Office of Graduate Study as a thesis project.

REFERENCES

- Hussein HS, Brasel JM. Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicology*. 2001;167(2):101-34.
- Arafa A, Bloomer R, Wilson H, Simpson C, Harms R. Susceptibility of various poultry species to dietary aflatoxin. *British Poultry Science*. 1981;22(5):431-6.
- Yunus AW, Razzazi-Fazeli E, Bohm J. Aflatoxin B1 in affecting broiler's performance, immunity, and gastrointestinal tract: A review of history and contemporary issues. *Toxins*. 2011;3(6):566-90.
- Huwig A, Freimund S, Käppeli O, Dutler H. Mycotoxin detoxication of animal feed by different adsorbents. *Toxicology letters*. 2001;122(2):179-88.
- Do JH, Choi D-K. Aflatoxins: detection, toxicity, and biosynthesis. *Biotechnology and Bioprocess Engineering*. 2007;12(6):585-93.
- Imanek V, Kottova N, Bartek J, Psotova J, Kosina P, Balejova LE, et al. Extract from *Silybum marianum* as a nutraceutical: a double-blind placebo-controlled study in healthy young men. *Czech J Food Sci* Vol. 2000;19(3):106-10.
- Chakarverty A, Parsad J. Study on the effect of Milk Thistle extract on the performance of broiler chicks. *Ind Poult Advis*. 1991;24(9):37-8.
- Abenavoli L, Capasso R, Milic N, Capasso F. Milk thistle in liver diseases: past, present, future. *Phytotherapy Research*. 2010;24(10):1423-32.
- Muhammad D, Chand N, Khan S, Sultan A, Mushtaq M. Hepatoprotective role of milk thistle (*Silybum marianum*) in meat type chicken fed aflatoxin B1 contaminated feed. *Pakistan Veterinary Journal*. 2012;32(3):443-6.
- Tedesco D, Steidler S, Galletti S, Tameni M, Sonzogni O, Ravarotto L. Efficacy of silymarin-phospholipid complex in reducing the toxicity of aflatoxin B1 in broiler chicks. *Poultry science*. 2004;83(11):1839-43.
- Kalorey D, Kurkure N, Ramgaonkar J, Sakhare P, Warke S, Nigot N. Effect of polyherbal feed supplement "Growell" during induced aflatoxicosis, ochratoxicosis and combined mycotoxicoses in broilers. *Asian-Australasian J Anim Sci*. 2005;18:375-83.
- Valdivia A, Martinez A, Damian F, Quezada T, Ortiz R, Martinez C, et al. Efficacy of N-acetylcysteine to reduce the effects of aflatoxin B1 intoxication in broiler chickens. *Poultry science*. 2001;80(6):727-34.
- Amiridumari H, Sarir H, Afzali N, FaniMakki O. Effects of milk thistle seed against aflatoxin B1 in broiler model. *Journal of research in medical sciences: the official journal of Isfahan University of Medical Sciences*. 2013;18(9):786-90.
- Basaga H, Poli G, Tekkaya C, Aras I. Free radical scavenging and antioxidative properties of 'silibin' complexes on microsomal lipid peroxidation. *Cell biochemistry and function*. 1997;15(1):27-33.
- Tapia M. A quantitative thin layer chromatography method for the analysis of aflatoxins, ochratoxin A, zearalenone, T-2 toxin and sterigmatocystin in foodstuffs. *Revista Argentina de microbiologia*. 1984;17(4):183-6.
- Bares JM, Berger J, Nelson JE, Messner DJ, Schildt S, Standish LJ, et al. Silybin treatment is associated with reduction in serum ferritin in patients with chronic hepatitis C. *Journal of clinical gastroenterology*. 2008;42(8):937-44.
- Schrieber SJ, Wen Z, Vourvahis M, Smith PC, Fried MW, Kashuba AD, et al. The pharmacokinetics of silymarin is altered in patients with hepatitis C virus and nonalcoholic fatty liver disease and correlates with plasma caspase-3/7 activity. *Drug Metabolism and Disposition*. 2008;36(9):1909-16.
- Fani Makki O, Omid A, Afzali N, Sarir H, Frouzanmehr M. Efficacy of *Silybum Marianum* Seeds in Ameliorating the Toxic Effects of Aflatoxin B1 in Broilers. *Iranian Journal of toxicology*. 2014;8(24):977-82.
- Gowda N, Ledoux D, Rottinghaus G, Bermudez A, Chen Y. Efficacy of turmeric (*Curcuma longa*), containing a known level of curcumin, and a hydrated sodium calcium aluminosilicate to ameliorate the adverse effects of aflatoxin in

- broiler chicks. Poultry science. 2008;87(6):1125-30.
20. Rastogi R, Srivastava AK, Srivastava M, Rastogi AK. Hepatocurative Effect of Picroliv and Silymarin Against Aflatoxin B1 Induced Hepatotoxicity in Rats. *Planta medica*. 2000;66(08):709-13.
21. Lutsenko S, Kashnikova T, Khmyrov A, Ledeshkova O, Fel'dman N, Luzhnov N, et al. Study of the effect of a liposomal form of silymarin on biochemical indices of the blood serum and productivity of broiler chicks. *Russian Agricultural Sciences*. 2008;34(6):415-7.
22. Wilasrusmeebce C, Kitturac S, Shahb G, Siddiquib J, Bruchbe D, Wilasrusmeeb S, et al. Immunostimulatory effect of Silybum marianum (milk thistle) extract. *Signature*. 2002;8(11):439-43.
23. Thyagarajan S, Jayaram S, Gopalakrishnan V, Hari R, Jeyakumar P, Sripathi M. Herbal medicines for liver diseases in India. *Journal of Gastroenterology and Hepatology*. 2002;17(s3):S370-S6.

Archive of SID