

## Serogroups, subtypes and virulence factors of shiga toxin-producing *Escherichia coli* isolated from human, calves and goats in Kerman, Iran

Rohollah Taghadosi<sup>1</sup>, Mohammad Reza Shakibaie<sup>1</sup>, Hesam Alizade<sup>2</sup>, Hossein Hosseini-Nave<sup>1</sup>, Asma Askari<sup>3</sup>, Reza Ghanbarpour<sup>3</sup>

<sup>1</sup>Department of Microbiology and Virology, Kerman University of Medical Sciences, Kerman, Iran

<sup>2</sup>Infectious and Tropical Disease Research Center, Hormozgan Health Institute, Hormozgan University of Medical Sciences, Bandar Abbas, Iran

<sup>3</sup>Department of Pathobiology, Faculty of Veterinary Medicine, Shahid Bahonar University of Kerman, Kerman, Iran

### ABSTRACT

**Aim:** The present study was conducted to detect the occurrence, serogroups, virulence genes and phylogenetic relationship of shiga toxin-producing *Escherichia coli* (STEC) in human, calves and goat in Kerman (southeast of Iran).

**Background:** STEC have emerged as the important foodborne zoonotic pathogens causing human gastrointestinal disease and confirming the risk to public health.

**Methods:** A total of 671 fecal samples were collected from diarrheic patients (n=395) and healthy calves (n=156) and goats (n=120) and screened for the presence of *stx* gene. Furthermore, the prevalence of *stx1* and *stx2* variants, serotypes (O157, O145, O103, O26, O111, O91, O128, and O45), phylogenetic groups and the presence of *ehxA*, *eae*, *hlyA*, *iha* and *saa* virulence genes were studied.

**Results:** Prevalence of STEC in human diarrheic isolates was 1.3% (5 isolates), in calves was 26.3% (41 isolates) and in goats was 27.5% (33 isolates). *stx1* gene was the most prevalent variant and detected in 75 isolates. Furthermore, *stx1c* was the most predominant *stx* subtype, found in 56 isolates. The *ehxA* identified in 36 (45.6%) isolates, followed by *iha* 5 (6.3%), *eaeA* 4 (5.1%), *hlyA* 2 (2.5%) and *saa* 2 (2.5%). Most of the isolates belonged to phylogroup B1. Only two O26 and one O91 isolates were detected in our study.

**Conclusion:** Our results show that STEC strains were widespread among healthy domestic animals in the southeast of Iran

**Keywords:** Shiga toxin-producing *E. coli*, serogroup, virulence factors.

(Please cite as: Taghadosi R, Shakibaie MR, Alizade H, Hosseini-Nave H, Askari A, Ghanbarpour R. Serogroups, subtypes and virulence factors of shiga toxin-producing *Escherichia coli* isolated from human, calves and goats in Kerman, Iran. *Gastroenterol Hepatol Bed Bench* 2018;11(1):60-67).

### Introduction

Shiga toxin-producing by *Escherichia coli* (STEC) is an important enteric pathogen, has been reported in several outbreaks with clinical manifestations including mild diarrhea, hemorrhagic colitis (HC) and hemolytic uremic syndrome (HUS) (1, 2). The disease in human is primarily a food-borne infection. Although STEC strains have been isolated from other animals such as goats, sheep, swine, wild animals and humans, cattle are the major source of food contamination (3). The

ability of STEC strains to cause human disease is mainly due to the production of shiga-like toxins (stx) which are classified into two closely related subgroups, *stx1* and *stx2* (encoded by the *stx1* and *stx2* genes). *Stx1* is a homologous group with only three variants (*stx1a*, *stx1c*, and *stx1d*), while *stx2* is a more heterogeneous group and is comprised of several subtypes (*stx2a*, *stx2b*, *stx2c*, *stx2d*, *stx2e*, *stx2f* and *stx2g*) (4, 5). STEC strains producing *stx2a*, *stx2c*, or *stx2d* subtypes are more associated with HC and HUS in humans. In contrast, *stx2b*, *stx2e*, *stx2f* and *stx2g* are related to animal infections (6). Additional factors that contribute to virulence have also been described, including intimin (encoded by the *eae* gene), involved

Received: 21 November 2017 Accepted: 27 January 2018

**Reprint or Correspondence:** Reza Ghanbarpour, PhD, Department of Pathobiology, Faculty of Veterinary Medicine, Shahid Bahonar University of Kerman, Kerman, Iran.

**E-mail:** ghanbar@uk.ac.ir

in the attachment of *E. coli* to the enterocyte, plasmid-encoded enterohemolysin (encoded by *ehxA* gene) which acts as a pore-forming cytolysin, alpha-hemolysin (encoded by the *hlyA* gene), IrgA homologue adhesin (*iha*) which is a STEC adherence-conferring molecule and Saa which is an autoagglutinating adhesin produced by LEE-negative STEC (3, 7-9). Epidemiologic investigations demonstrated that O157 is the main cause of HC and HUS in human; however, additional serogroups that have been reported in human clinical cases are O26, O45, O91, O103, O111, O128 and O145, and others in recent years (10, 11).

*E. coli* can also be assigned to one of the four major phylogenetic groups (A, B1, B2 and D) based on the presence or absence of *chuA*, *yjaA* and TspE4.C2 (12). Bearing in mind the importance of *E. coli* as food-borne pathogens, as vehicle of human disease, the objectives of this study were to investigate the distribution of subtypes, serotypes, virulence factors and phylogenetic groups among STEC strains from healthy domestic animals (calves and goats) and patients with diarrhea in Kerman, southeast of Iran.

## Methods

### Specimen collection and microbiological processing

In a prospective study, from October 2014 to November 2015, a total of 671 fecal samples were collected from diarrheic patients (n=395) and fecal healthy calves (n=156) and goats (n=120). The human samples were related to both male (n=215) and female (n=180). Their age ranged from <5 years old (n=107), 5 to 15 years old (n=146), 15 to 40 years old (n=75) and 40 to 90 years old (n=67). The human isolates obtained from the rectal swab of the patient with diarrhea referred to Afzalipour and Payambar-Azam hospitals. All animal samples were collected by veterinarians from School of Veterinary Medicine, Shahid Bahonar University, Kerman, Iran. All samples were placed into Amies medium (Becton Dickinson, BBL, and USA) and were sent out to the laboratory in ice-cooled containers. The samples were taken to the microbiology laboratory, Kerman University of Medical Sciences and identified as *E. coli* by biochemical characteristics and conventional diagnostic tests (13). All strains were stored at  $-70^{\circ}\text{C}$  in Trypticase Soy broth (Difco

Laboratories, Detroit, Mich.) containing 30% glycerol for further study.

### Detection of STEC strains

For the detection of *stx* gene, DNA template was obtained by boiling method (14). Presence of *stx* gene in the selected *E. coli* colonies was verified by PCR method (15). In addition, *stx*-positive isolates were examined for the presence of *stx*<sub>1</sub> and *stx*<sub>2</sub> genes by using duplex-PCR (16). A positive control for PCR was *E. coli* Sakai (*stx*<sub>1</sub>+/*stx*<sub>2</sub>+/*eaeA*+). The *E. coli* strain MG1655 was used as a negative control for virulence genes. Details of the primers and the length of the expected amplification product are listed in Table 1.

### Identification of subtype genes

We used PCR method for determination of *stx*<sub>1</sub> and *stx*<sub>2</sub> subtypes. PCR for detection of *stx*<sub>1a</sub>, *stx*<sub>1c</sub>, *stx*<sub>1d</sub>, *stx*<sub>2a</sub>, *stx*<sub>2b</sub>, *stx*<sub>2c</sub>, *stx*<sub>2d</sub>, *stx*<sub>2e</sub>, *stx*<sub>2f</sub> and *stx*<sub>2g</sub> subtypes was carried out by methods described previously (17-19) (Table 1).

### Identification of serogroup genes

Furthermore, PCR assay was used for the identification of O157, O145, O103, O26, O111, O91, O128 and O45 as described by Hemmatinezhad *et al.* (20) (Table 1).

### Identification of virulence genes

The presence of following virulence genes *ehxA*, *eae*, *hlyA*, *iha* and *saa* were detected by PCR assay (21-24) (Table 1).

### Determination of STEC strains phylogenetic groups

Strains assigned to one of the four main phylogenetic group of *E. coli* (A, B1, B2 and D) by using a PCR-based method as described previously (12). The genomic DNA of bacterial strains amplified by triplex-PCR using primers targeted at three markers, *chuA*, *yjaA* and TspE4.C2.

### Statistical analysis

SPSS version 15.0 software for Windows (SPSS Inc., Chicago) was used for statistical analysis. *P* values of less than 0.05 were considered to be significant.

## Results

Among 671 *E. coli* isolates isolated from healthy farm calves, goats and patients with sign of diarrhea, 79 strains were positive for the presence of *stx* gene and identified as STEC. Among STEC strains 41 strains were positive in calves, 33 strains in goats and 5 strains

62 Molecular characterization of STEC

Table 1. Oligonucleotide Primers Used in this Study.

| Target gene  | Primer sequence (5'-3')                                 | Size (bp) | Annealing temp (°C) | References |
|--------------|---|-----------|---------------------|------------|
| <i>stx</i>   | GAGCGAAATAATTTATATGTG<br>TGATGATGGCAATTCAGTAT           | 518       | 55                  | 15         |
| <i>stx1</i>  | TAAATCGCCATTCGTTGACTAC<br>AGAACGCCCACTGAGATCATC         | 180       | 58                  | 16         |
| <i>stx2</i>  | GGCACTGTCTGAAACTGCTCC<br>TCGCCAGTTATCTGACATTCTG         | 255       | 60                  | 16         |
| <i>stx1a</i> | CACGTTACAGCGTGTGCA<br>CGCCCACTGAGATCATCC                | 219       | 57                  | 18         |
| <i>stx1c</i> | TTTTACATGTTACCTTTCTC<br>CATAGAAGGAAACTCATTAGG           | 498       | 54                  | 17         |
| <i>stx1d</i> | CTTTTCAGTTAATGCGATTGCT<br>AACCCCATGATATCGACTGC          | 192       | 57                  | 18         |
| <i>stx2a</i> | AGATATCGACCCCTCTTGAA<br>GTCAACCTTCACTGTAATG             | 969       | 55                  | 18         |
| <i>stx2b</i> | AAATATGAAGAAGATATTTGTAGCGGC<br>CAGCAAATCCTGAACCTGACG    | 251       | 60                  | 19         |
| <i>stx2c</i> | GCGGTTTTATTTGCATTAGT<br>AGTACTCTTTCCGGCCACT             | 124       | 55                  | 17         |
| <i>stx2d</i> | GGTAAAATTGAGTTCTCTAAGTAT<br>CAGCAAATCCTGAACCTGACG       | 175       | 58                  | 17         |
| <i>stx2e</i> | ATGAAGAAGATGTTTATAGCG<br>TCAGTTAAACTTCACTGGGC           | 267       | 56                  | 17         |
| <i>stx2f</i> | TGGGCGTCATTCCTGTTG<br>TAATGGCCGCCCTGTCTCC               | 424       | 60                  | 19         |
| <i>stx2g</i> | CACGGGTAGTTATATTTCTGTGGATATC<br>GATGGCAATTCAGAATAACCGCT | 573       | 62                  | 19         |
| <i>chuA</i>  | GACGAACCAACGGTCAGGAT<br>TGCCGCCAGTACCAAAGACA            | 279       | 59                  | 12         |
| <i>YjaA</i>  | TGAAGTGTACAGGAGACGCTG<br>ATGGAGAATGCGTTCCTCAAC          | 211       | 59                  | 12         |
| TspE4.C2     | CTGGCG AAAGACTGTATCAT<br>CGCGCCAACAAAGTATTA CG          | 152       | 59                  | 12         |
| <i>ehxA</i>  | GGTGCAGCAGAAAAAGTTGTAG<br>TCTCGCCTGATAGTGTTTGGTA        | 1551      | 61                  | 24         |
| <i>hlyA</i>  | AACAAGGATAAGCACTGTTCTGGCT<br>ACCATATAAGCGGTCATTCCCGTCA  | 1177      | 61                  | 21         |
| <i>saa</i>   | CGTGATGAACAGGCTATTGC<br>ATGGACATGCCTGTGGCAAC            | 119       | 57                  | 23         |
| <i>iha</i>   | CTGGCGGAGGCTCTGAGATCA<br>TCCTTAAGCTCCCGCGGCTGA          | 827       | 57                  | 23         |
| <i>eaeA</i>  | CAGGTCGTCGTCTGCTAAA<br>TCAGCGTGGTTGGATCAACCT            | 1087      | 65                  | 22         |
| O157         | CGGACATCCATGTGATATGG<br>TTGCCTATGTACAGCTAATCC           | 259       | 58                  | 20         |
| O145         | CCATCAACAGATTTAGGAGTG<br>TTTCTACCGCGAATCTATC            | 609       | 58                  | 20         |
| O111         | TAGAGAAATTATCAAGTTAGTTCC<br>ATAGTTATGAACATCTTGTTAGC     | 406       | 58                  | 20         |
| O91          | GCTGACCTTCATGATCTGTTGA<br>TAATTTAACCCGTAGAATCGCTGC      | 291       | 58                  | 20         |
| O128         | GCTTTCTGCCGATATTGGC<br>CCGACGGACTGATGCCGGTGATT          | 289       | 58                  | 20         |
| O45          | CCGGGTTTTCGATTTGTGAAGTTG<br>CACAACAGCCACTACTAGGCAGAA    | 527       | 58                  | 20         |
| O103         | TGGAGCGTTAACTGGACCT<br>GCTCCCGAGCACGTATAAG              | 321       | 58                  | 20         |
| O26          | CAGAATGGTTATGCTACTGT<br>CTTACATTTGTTTTCCGGCATC          | 423       | 58                  | 20         |

**Table 2.** Summary of *stx* subtyping in 79 non O157-STECC strains isolated from calves, goats and human fecal samples.

| No (%) of strains | source |        |       | <i>stx</i> subtype       |                          |                          |                          |                          |                          |
|-------------------|--------|--------|-------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|--------------------------|
|                   | humans | calves | goats | <i>stx</i> <sub>1a</sub> | <i>stx</i> <sub>1c</sub> | <i>stx</i> <sub>2a</sub> | <i>stx</i> <sub>2b</sub> | <i>stx</i> <sub>2c</sub> | <i>stx</i> <sub>2d</sub> |
| 35 (44.3)         | 2      | 20     | 13    | -                        | +                        | -                        | -                        | -                        | -                        |
| 16 (20.3)         | 1      | 6      | 9     | +                        | -                        | -                        | -                        | -                        | -                        |
| 8 (10.1)          | 0      | 4      | 4     | -                        | +                        | -                        | +                        | +                        | +                        |
| 5 (6.3)           | 0      | 5      | 0     | -                        | +                        | -                        | +                        | +                        | -                        |
| 5 (6.3)           | 0      | 2      | 3     | +                        | +                        | -                        | -                        | -                        | -                        |
| 2 (2.5)           | 2      | 0      | 0     | -                        | -                        | -                        | +                        | -                        | -                        |
| 2 (2.5)           | 0      | 2      | 0     | +                        | +                        | -                        | +                        | +                        | +                        |
| 1 (1.3)           | 0      | 0      | 1     | +                        | -                        | -                        | -                        | +                        | -                        |
| 1 (1.3)           | 0      | 1      | 0     | -                        | +                        | -                        | +                        | -                        | -                        |
| 1 (1.3)           | 0      | 0      | 1     | +                        | -                        | +                        | -                        | -                        | -                        |
| 1 (1.3)           | 0      | 0      | 1     | -                        | -                        | -                        | +                        | +                        | -                        |
| 1 (1.3)           | 0      | 1      | 0     | -                        | -                        | -                        | +                        | +                        | +                        |
| 1 (1.3)           | 0      | 0      | 1     | -                        | -                        | -                        | -                        | +                        | -                        |
| Total= 79         | 5      | 41     | 33    | 25                       | 56                       | 1                        | 20                       | 19                       | 11                       |

No = Number of non O157-STECC strains, % = percent of non O157-STECC strains; - = negative, + = positive.

**Table 3.** Distribution of STECC in phylogenetic groups from calves, goats and human fecal samples

| Phylogenetic group | No. (%) of STECC strains isolated from |           |           | Total     |
|--------------------|--|-----------|-----------|-----------|
|                    | humans                                 | calves    | goats     |           |
| A                  | 0 (0)                                  | 6 (14.6)  | 6 (18.2)  | 12 (15.2) |
| B1                 | 0 (0)                                  | 35 (85.4) | 27 (81.8) | 62 (78.5) |
| B2                 | 0 (0)                                  | 0 (0)     | 0 (0)     | 0 (0)     |
| D                  | 5 (100)                                | 0 (0)     | 0 (0)     | 5 (6.3)   |

NO = Number of STECC strains isolated from calves, goats and human fecal samples, % = percent of STECC strains isolated from calves, goats and human fecal samples

in human samples. Our results showed that 54 (68.4%) of the strains carried *stx*<sub>1</sub> only, 4 (5.1%) contained *stx*<sub>2</sub> only, and 21 (26.6%) possessed both *stx*<sub>1</sub> and *stx*<sub>2</sub>.

Two *stx*<sub>1</sub> subtypes (*stx*<sub>1a</sub> and *stx*<sub>1c</sub>) and four *stx*<sub>2</sub> subtypes (*stx*<sub>2a</sub>, *stx*<sub>2b</sub>, *stx*<sub>2c</sub> and *stx*<sub>2d</sub>) were detected with a total of 13 different *stx*<sub>1</sub> and *stx*<sub>2</sub> subtypes combinations as shown in Table 2. Among the subtypes, *stx*<sub>1c</sub> was detected in 56 strains, followed by *stx*<sub>1a</sub> (25 strains), *stx*<sub>2b</sub> (20 strains), *stx*<sub>2c</sub> (19 strains) and *stx*<sub>2a</sub> (1 strain). In addition, *stx*<sub>2d</sub> (11 strains) was detected only in combination with other *stx* genes (Table 2). There was no correlation between *stx* subtypes and animal sources ( $P \leq 0.05$ ).

The STECC strains were further tested for five putative virulence factors, including *eaeA*, *hlyA*, *iha*, *ehxA* and *saa*. Out of 79 strains, 49 (62%) carried at least one virulence gene tested. The *ehxA* was detected in 36 (45.6%), *eaeA* in 4 (5.1%), *iha* in 5 (6.3%), *hlyA* in 2 (2.5%) and *saa* in 2 (2.5%) of the isolates.

Phylogroup B1 was the most prevalent (62/79; 78.5%) among the STECC strains, followed by phylogroups A (12/79; 15.2%) and D (5/79; 6.3%). As shown in Table 3, all isolates of human origin belonged

to the D Phylogroup. In this study, phylogenetic group B2 was not detected in STECC strains.

Serogroup analysis showed that none of the isolates belonged to O45, O103, O111, O128, O145 and O157 serogroups, while O26 and O91 were detected in two (clave and goat) and one (clave) isolate, respectively.

### Discussion

STECC can be found in various food sources, transmission of this pathotype from undercooked or unpasteurized animal products to human is problematic (1, 10). It is estimated that STECC to cause more than 265,000 illnesses each year in the USA, with more than 3,600 hospitalizations and 30 deaths (25). In the present study, STECC strains were isolated just from 1.2% of patients with diarrhea which was consistent with previous studies (26-28). According to a survey, high variability of genes-encoding *stx* was detected in the *E. coli* isolates in HIV and thalassemia patients in Kerman, south-east of Iran. Among *E. coli* isolates from faecal samples, 30.8% isolates were positive for *stx* genes (34). However, 26.8% of *E. coli* isolated from goats and calves carried at least one of the *stx* genes.

#### 64 Molecular characterization of STEC

This frequency was, lower compared to reports from Spain and Brazil (37% and 44%) (29, 30), but higher from results reported in Iran (8.5%) (31). Another study from West Azerbaijan province in Iran revealed that 21.92% of the *E. coli* isolates recovered from fecal healthy calves harbored *stx* genes (32). These variations may likely be due to geographical and climatic conditions and differences in the natural intestinal flora present in animal's gastrointestinal tract (33).

In STEC strains characterized in this study, *stx*<sub>1</sub> was the most common *stx* gene identified, a result which is similar to previous reports (31, 34). In contrast, some studies have detected *stx*<sub>2</sub> as a dominant *stx* gene in fecal samples of animals (35, 36). Although, this variant mainly found in strains isolated from healthy human carriers and most likely does not cause severe diseases in human (36).

In the present study, *stx*<sub>1c</sub> was the predominant variant among the STEC strains isolated. *Stx*<sub>1c</sub> Subtype has also frequently been reported in previous studies (5, 37). However, *stx*<sub>1c</sub>-encoding strains are associated with asymptomatic human carriage or mild illness (38). *Stx*<sub>2c</sub> and *stx*<sub>2d</sub> are associated with HUS. However, they are less toxic on Vero cells compared to *stx*<sub>2a</sub>. STEC strains with *stx*<sub>2a</sub> are associated with several clinical symptoms, such as HUS and HC (39). Stephan and Hoelzle suggested that *stx*<sub>2b</sub> was not associated with severe human diseases, because most strains carrying *stx*<sub>2b</sub> were isolated from healthy human carriers (40). In the present study, two strains carrying only *stx*<sub>2b</sub> were isolated from human and it was possible that these two STEC strains were not the main causative agent of diarrhea. In this study, two strains isolated from calves carried 5 subtypes of *stx*<sub>1a</sub>, *stx*<sub>1c</sub>, *stx*<sub>2b</sub>, *stx*<sub>2c</sub>, *stx*<sub>2d</sub> simultaneously. The combination of five *stx* genes in one isolate had not been previously reported. In the study of Bertin *et al.* strains with a combination of *stx*<sub>1</sub> and/or *stx*<sub>2</sub> subtypes were found to be more toxic toward Vero cells than other strains (41). In our study, other *stx*<sub>2</sub> subtypes such as *stx*<sub>2e</sub>, *stx*<sub>2f</sub> and *stx*<sub>2g</sub> were not found. These subtypes are related to animal infections (42).

In addition, we studied the distribution of eight important serotypes in the above isolates which associated more frequently with HUS and HC. None of the isolates belonged to O45, O103, O111, O128, O145 and O157 serogroups, while O26 and O91 were

detected in two and one isolates respectively. This finding is in agreement with the failure to find these serotypes in yaks and cattle (8, 43). It seems that in some regions, ruminants are not important reservoirs for the outbreak isolates. Although, human infections with *stx*-producing *E. coli* O26 is uncommon and has resulted in less severe illness, but is a major cause of HUS in Europe continent (44).

In this study, only four strains contained *eaeA* gene; however, none of the isolates carried the *stx*<sub>2</sub> subtypes. The low frequency of the *eaeA* gene found in the present study may be related to the low frequency of certain serogroups, as it has been reported that the presence of the *eaeA* gene is associated with specific O serogroups of STEC, such as the O157, O145, O103, O26, and O111 (33). Since the majority of the STEC strains lacked *eaeA* gene, we investigated other factors associated with adherence including *iha* and *saa*. These two virulence factors have been reported to be highly important for pathogenicity of *eae*-negative STEC strains (8). Only 2.5% and 6.35% of strains were positive for *saa* and *iha* respectively. It is possible that other virulence factors, that were not investigated in the present study like *lpfa* and *paa* play important role in the adherence of STEC strains. Also, we detected *ehxA* and *hlyA* genes in 45.6% and 2.5% of strains respectively. Overall, the frequency of virulence factors in STEC isolates was lower than that observed in other studies (8, 45). Carriage of *stx* gene positive *E. coli* isolates in the gastrointestinal tract of healthy ruminants proposes that these are transient commensal bacteria in these animals and the virulence genes of these isolates were either not or very poorly expressed (32).

Investigation on STEC phylogroups indicated that majority of commensal and diarrhogenic strains are belonged to group B1 and A, while extra intestinal *E. coli* strains belong mainly to group B2 and D (46). In this study, phylogenetic group B2 were not detected in STEC isolates, which was consistent to previous study (46). However, like in many studies, phylogenetic group B1 was predominant among isolates from animals (47, 48). All of the human strains belonged to phylogenetic group D2, while it was not found in strains isolated from animals.

In conclusion, although STEC strains were widespread among healthy domestic animals in the southeast of Iran, prevalence of STEC in patient with diarrhea was

low and most of the STEC strains did not belong to O serogroups that are commonly associated with severe disease in humans. Furthermore, these strains were mainly belonged to phylogenetic group B1. These facts together with the high prevalence of *stx*<sub>1c</sub>, *stx*<sub>2b</sub>, *stx*<sub>2c</sub> subtypes and low prevalence of *stx*<sub>2a</sub>, suggest that most of STEC in Iranian calves and goats may not pose a serious public health concern.

### Acknowledgment

The authors acknowledge Kerman University of Medical Sciences for the financial support.

### Conflict of interests

The authors declare that they have no conflict of interest.

### References

- Doregirae F, Alebouyeh M, Nayeri Fasaee B, Charkhkar S, Tajedin E, Zali MR. Isolation of atypical enteropathogenic and shiga toxin encoding *Escherichia coli* strains from poultry in Tehran, Iran. *Gastroenterol Hepatol Bed Bench* 2016;9:53-7.
- Kargar M, Homayoon M. Prevalence of shiga toxins (*stx1*, *stx2*), *eaeA* and *hly* genes of *Escherichia coli* O157: H7 strains among children with acute gastroenteritis in southern of Iran. *Asian Pac J Trop Med* 2015;8:24-8.
- Baliere C, Rince A, Delannoy S, Fach P, Gourmelon M. Molecular profiling of shiga toxin-producing *Escherichia coli* and enteropathogenic *E. coli* strains isolated from French coastal environments. *Appl Environ Microbiol* 2016;82:3913-27.
- Jajarmi M, Imani Fooladi AA, Askari Badouei M, Ahmadi A. Virulence genes, shiga toxin subtypes, major O-serogroups, and phylogenetic background of shiga toxin-producing *Escherichia coli* strains isolated from cattle in Iran. *Microb Pathog* 2017;109:274-9.
- Vu-Khac H, Cornick NA. Prevalence and genetic profiles of shiga toxin-producing *Escherichia coli* strains isolated from buffaloes, cattle, and goats in central Vietnam. *Vet Microbiol* 2008;126:356-3.
- Fuller CA, Pellino CA, Flagler MJ, Strasser JE, Weiss AA. Shiga toxin subtypes display dramatic differences in potency. *Infect Immun* 2011;79:1329-37.
- Beutin L, Aleksic S, Zimmermann S, Gleier K. Virulence factors and phenotypical traits of verotoxigenic strains of *Escherichia coli* isolated from human patients in Germany. *Med Microbiol Immunol* 1994;183:13-21.
- Bai X, Zhao A, Lan R, Xin Y, Xie H, Meng Q, et al. Shiga toxin-producing *Escherichia coli* in Yaks (*Bos grunniens*) from the Qinghai-Tibetan Plateau, China. *PLoS One* 2013;8:e65537.
- Caine L, Nwodo U, Okoh A, Ndip R, Green E. Occurrence of virulence genes associated with diarrheagenic *Escherichia coli* isolated from raw cow's milk from two commercial dairy farms in the eastern cape province, South Africa. *Int J Environ Res Public Health* 2014;11:11950-63.
- Wang F, Yang Q, Kase J, Meng J, Clotilde LM, Lin A, et al. Current trends in detecting non-O157 shiga toxin-producing *Escherichia coli* in food. *Foodborne Pathog Dis* 2013;10:665-77.
- Taghadosi R, Shakibaie MR, Ghanbarpour R, Hosseini-Nave H. Role of antigen-43 on biofilm formation and horizontal antibiotic resistance gene transfer in non-O157 shiga toxin producing *Escherichia coli* strains. *Iran J Microbiol* 2017;9:89-96.
- Clermont O, Bonacorsi S, Bingen E. Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl Environ Microbiol* 2000;66:4555-8.
- Patricia M. Bailey & Scott's Diagnostic Microbiology. St. Louis, Missouri: Elsevier, 2014.
- Hosseini Nave H, Mansouri S, Emaneini M, Moradi M. Distribution of genes encoding virulence factors and molecular analysis of *Shigella* spp. isolated from patients with diarrhea in Kerman, Iran. *Microb Pathog* 2016;92:68-71.
- Nakhjavani FA, Emaneini M, Hosseini H, Iman-Eini H, Aligholi M, Jabalameli F, et al. Molecular analysis of typical and atypical enteropathogenic *Escherichia coli* (EPEC) isolated from children with diarrhoea. *J Med Microbiol* 2013;62:191-5.
- Paton A, Paton J. Detection and characterization of shiga toxigenic *Escherichia coli* by using multiplex enterohemorrhagic *E. coli* hlyA, rfb O111, and detection and characterization of shiga toxigenic *Escherichia coli* by using multiplex PCR assays for *stx1*, *stx2*, *eae*. *J Clin Microbiol* 1998;36:598-602.
- Kumar A, Taneja N, Kumar Y, Sharma M. Detection of Shiga toxin variants among shiga toxin-forming *Escherichia coli* isolates from animal stool, meat and human stool samples in India. *J Appl Microbiol* 2012;113:1208-16.
- He X, Quiñones B, McMahon S, Mandrell RE. A single-step purification and molecular characterization

## 66 Molecular characterization of STEC

of functional Shiga toxin 2 variants from pathogenic *Escherichia coli*. *Toxins* 2012;4:487-504.

19. Scheutz F, Teel LD, Beutin L, Piérard D, Buvens G, Karch H, et al. Multicenter evaluation of a sequence-based protocol for subtyping shiga toxins and standardizing Stx nomenclature. *J Clin Microbiol* 2012;50:2951-63.

20. Hemmatinezhad B, Khamesipour F, Mohammadi M, Safarpour Dehkordi F, Mashak Z. Microbiological investigation of O-serogroups, virulence factors and antimicrobial resistance properties of shiga toxin-producing *Escherichia coli* isolated from Ostrich, Turkey and Quail Meats. *J Food Saf* 2015;35:491-500.

21. Yamamoto S, Terai A, Yuri K, Kurazono H, Takeda Y, Yoshida O. Detection of urovirulence factors in *Escherichia coli* by multiplex polymerase chain reaction. *FEMS Immunol Med Microbiol* 1995;12:85-90.

22. Gannon VP, D'souza, S, Graham T, King RK, Rahn K, Read S. Use of the flagellar H7 gene as a target in multiplex PCR assays and improved specificity in identification of enterohemorrhagic *Escherichia coli* strains. *J Clin Microbiol* 1997;35:656-62.

23. Chapman TA, Wu XY, Barchia I, Bettelheim KA, Driesen S, Trott D, et al. Comparison of virulence gene profiles of *Escherichia coli* strains isolated from healthy and diarrheic swine. *Appl Environ Microbiol* 2006;72:4782-95.

24. Meng Q, Bai X, Zhao A, Lan R, Du H, Wang T, et al. Characterization of shiga toxin-producing *Escherichia coli* isolated from healthy pigs in China. *BMC Microbiol* 2014;14:5.

25. Scallan E, Hoekstra RM, Angulo FJ, Tauxe RV, Widdowson MA, Roy SL, et al. Foodborne illness acquired in the United States-major pathogens. *Emerg Infect Dis* 2011;17:7-15.

26. Alizade H, Ghanbarpour R, Aflatoonian MR. Molecular study on diarrheagenic *Escherichia coli* pathotypes isolated from under 5 years old children in southeast of Iran. *Asian Pac J Trop Dis* 2014;4:S813-7.

27. Alizade H, Ghanbarpour R, Nekoubin M. Phylogenetic of shiga toxin-producing *Escherichia coli* and atypical enteropathogenic *Escherichia coli* strains isolated from human and cattle in Kerman, Iran. *Int J Entric Pathog* 2014;1:e15195.

28. Nguyen BM, Phung DC, Nakasone N, Toma C, Higa N, Iyoda S, et al. Shiga-toxin producing *Escherichia coli* in Vietnam. *Trop Med Health* 2004;32:339-41.

29. Blanco M, Blanco JE, Blanco J, Mora A, Prado C, Alonso MP, et al. Distribution and characterization of faecal verotoxin-producing *Escherichia coli* (VTEC) isolated from healthy cattle. *Vet. Microbiol* 1997;54:309-19.

30. Pereira MA, Brod CS, Rodrigues DP, Carvalhal JB, Aleixo JAG. Shiga toxin-producing *Escherichia coli* (STEC) isolated from healthy dairy cattle in southern Brazil. *Vet Microbiol* 2003;93:179-83.

31. Sepehriseresht S, Zahraei Salehi T, Sattari M, Tadjbakhsh H, Aslani M. Detection of shiga toxigenic *Escherichia coli* from fecal samples of calves and cattle by molecular and serological methods. *Comp Clin Path* 2009;18:53-7.

32. Dastmalchi Saei H, Ayremlou N. Characterization of shiga toxin-producing *Escherichia coli* (STEC) in feces of healthy and diarrheic calves in Urmia region, Iran. *Iran J Microbiol* 2012;4:63-9.

33. Yaghobzadeh N, Ownagh A, Mardani K, Khalili M. Prevalence, molecular characterization and serology of shiga toxin-producing *Escherichia coli* isolated from buffaloes in west Azerbaijan, Iran. *Int J Vet Res* 2011;5:113-7.

34. Alizade H, Sharifi H, Naderi Z, Ghanbarpour R, Bamorovat M, Aflatoonian MR. High frequency of diarrheagenic *Escherichia coli* in HIV-infected patients and patients with thalassemia in Kerman, Iran. *J Int Assoc Provid AIDS Care* 2017;16:353-8.

35. Oporto B, Esteban JI, Aduriz G, Juste RA, Hurtado A. *Escherichia coli* O157:H7 and non-O157 shiga toxin-producing *E. coli* in healthy cattle, sheep and swine herds in Northern Spain. *Zoonoses Public Health* 2008;55:73-81.

36. Hofer E, Cernela N, Stephan R. Shiga toxin subtypes associated with shiga toxin-producing *Escherichia coli* strains isolated from red deer, roe deer, chamois, and ibex. *Foodborne Pathog Dis* 2012;9:792-5.

37. Nguyen TD, Vo TT, Vu-Khac H. Virulence factors in *Escherichia coli* isolated from calves with diarrhea in Vietnam. *J Vet Sci* 2011;12:159-64.

38. Zhang W, Bielaszewska M, Kuczius T, Karch H. Identification, characterization, and distribution of a shiga toxin 1 gene variant (stx1c) in *Escherichia coli* strains isolated from humans. *J Clin Microbiol* 2002;40:1441-6.

39. Melton-Celsa AR. Shiga Toxin (Stx) Classification, Structure, and Function. *Microbiol Spectr* 2014;2:1-21.

40. Stephan R, Hoelzle LE. Characterization of shiga toxin type 2 variant B-subunit in *Escherichia coli*

strains from asymptomatic human carriers by PCR-RFLP. *Lett Appl Microbiol* 2000;31:139-42.

41. Bertin Y, Boukhors K, Pradel N, Livrelli V, Martin C. Stx2 subtyping of shiga toxin-producing *Escherichia coli* isolated from cattle in France: detection of a new stx2 subtype and correlation with additional virulence factors. *J Clin Microbiol* 2001;3:3060-5.

42. Feng PCH, Reddy S. Prevalence of shiga toxin subtypes and selected other virulence factors among Shiga-toxigenic *Escherichia coli* strains isolated from fresh produce. *Appl Environ Microbiol* 2013;79:6917-23.

43. Auvray F, Dilasser F, Bibbal D, Kérourédan M, Oswald E, Brugère H. French cattle is not a reservoir of the highly virulent enteroaggregative shiga toxin-producing *Escherichia coli* of serotype O104: H4. *Vet Microbiol* 2012;158:443-5.

44. Chase-Topping ME, Rosser T, Allison LJ, Courcier E, Evans J, McKendrick IJ, et al. Pathogenic Potential to Humans of Bovine *Escherichia coli* O26, Scotland. *Emerg Infect Dis* 2012;18:439-48.

45. Franz E, Van Hoek AHAM, Wuite M, Van Der Wal FJ, De Boer AG, Bouw EI, et al. Molecular hazard identification of non-O157 shiga toxin-producing *Escherichia coli* (STEC). *PLoS One* 2015;10:e0120353.

46. Carlos C, Pires MM, Stoppe NC, Hachich EM, Sato MIZ, Gomes TAT, et al. *Escherichia coli* phylogenetic group determination and its application in the identification of the major animal source of fecal contamination. *BMC Microbiol* 2010;10:161.

47. Ishii S, Meyer KP, Sadowsky MJ. Relationship between phylogenetic groups, genotypic clusters, and virulence gene profiles of *Escherichia coli* strains from diverse human and animal sources. *Appl Environ Microbiol* 2007;73:5703-10.

48. Mora A, López C, Dhahi G, López-Beceiro AM, Fidalgo LE, Díaz EA, et al. Seropathotypes, phylogroups, stx subtypes, and intimin types of wildlife-carried, shiga toxin-producing *Escherichia coli* strains with the same characteristics as human-pathogenic isolates. *Appl Environ Microbiol* 2012;78:2578-85.