Available online at http://ijim.srbiau.ac.ir

Int. J. Industrial Mathematics Vol. 3, No. 1 (2011) 25-33





# An Iterative Approach for Estimation of Efficiency by Weighted Distance

R. Saneifard  $\,^*$ 

Department of Mathematics, Oroumieh Branch, Islamic Azad University, Oroumieh, Iran. Received 26 April 2010; revised 11 December 2010; accepted 26 December 2010.

#### Abstract

In group decision analysis, numerous approaches have been suggested in an attempt to solve the problem of aggregation of individual fuzzy opinion to form a group consensus as the basis of a group decision. If the inputs and outputs are fuzzy quantity, the decision making units can not be easily evaluated and ranked using the obtained efficiency scores. In this article, a kind of modified idea based on interactive method is introduced for ranking of decision making units with fuzzy data by weighted distance.

Keywords: Fuzzy number; Efficiency; Data envelopment analysis; Weighted distance.

## 1 Introduction

More and more, modeling techniques, control problems and operation research algorithms have been designed to fuzzy data since the concept of fuzzy number and arithmetic operations with these numbers was introduced and investigated first by Zadeh. Data envelopment analysis was suggested by Charnes, Cooper and Rhodes [2], and built on the idea of Farrell [3] which is concerned with the estimation of technical efficiency and efficient frontiers. In some cases, we have to use imprecise input and output. To deal quantitatively with imprecision in decision progress, Bellman Zadeh [1] introduce the notion of fuzziness. Some researchers have proposed several fuzzy models to evaluate decision making units with fuzzy data and introduce a ranking approach with efficiency measure of the model [9]. In this paper, the researcher first introduces an approach with weighted distance for ranking of decision making units with crisp data. Second, this model is used for ranking of decision making units with fuzzy data.

This paper is organized as follows. In Section 2, the researcher recalls some fundamental

<sup>\*</sup>Email address: Saneifard@iaurmia.ac.ir, Tel:+989149737077

results on fuzzy numbers. The proposed model is introduced in Section 3. An approach for ranking by using weighted distance is introduced in section 4. Interactive method is introduced in Section 5.

#### 2 Preliminaries

The basic definition of a fuzzy number given in [5, 11, 12, 13, 14, 15] is as follows:

**Definition 2.1.** A fuzzy number is a mapping  $\mu : \Re \rightarrow [0,1]$  with the following properties:

- 1.  $\mu$  is an upper semi-continuous function on  $\Re$ ,
- 2.  $\mu(x) = 0$  outside of some interval  $[a_1, b_2] \subset \Re$ ,
- 3. There are real numbers  $a_2, b_1$  such as  $a_1 \leq a_2 \leq b_1 \leq b_2$  and
  - 3.1  $\mu(x)$  is a monotonic increasing function on  $[a_1, a_2]$ ,
  - 3.2  $\mu(x)$  is a monotonic decreasing function on  $[b_1, b_2]$ ,

3.3 
$$\mu(x) = 1$$
 for all x in  $[a_2, b_1]$ .

**Definition 2.2.** A fuzzy number is a fuzzy set A on the real line  $\Re$  such that

$$\mu(x) = \begin{cases} f_A(x) & if \quad x \in [a_1, a_2], \\ 1 & if \quad x \in [a_2, a_3], \\ g_A(x) & if \quad x \in [a_3, a_4], \\ 0 & otherwise . \end{cases}$$
(2.1)

Such that  $f_A(.)$  is increasing function on  $[a_1, a_2]$  and  $g_A(.)$  is decreasing function on  $[a_3, a_4]$ .

The  $\alpha$ -cut of a fuzzy number A is defined as  $[A]^{\alpha} = \{x \mid u(x) \geq \alpha\}$ . Since  $\mu(.)$  is upper semi-continuous then  $\alpha$ -cuts are closed and bounded intervals and we represent by  $[A]^{\alpha} = [f_A^{-1}(\alpha), g_A^{-1}(\alpha)].$ 

**Definition 2.3.** A fuzzy number A in parametric form is a pair  $(\underline{A}, \overline{A})$  of functions  $\underline{A}(\alpha)$ and  $\overline{A}(\alpha)$  that  $0 \le \alpha \le 1$ , which satisfies the following requirements:

- 1.  $\underline{A}(\alpha)$  is a bounded monotonic increasing left continuous function,
- 2.  $\overline{A}(\alpha)$  is a bounded monotonic decreasing left continuous function,
- 3.  $\underline{A}\alpha \leq \overline{A}(\alpha), 0 \leq \alpha \leq 1.$

**Definition 2.4.** The symmetric triangular fuzzy number  $A = (x_0, \sigma)$ , with defuzzifier  $x_0$ and fuzziness  $\sigma$  is a fuzzy set where the membership function is as

$$\mu(x) = \begin{cases} \frac{1}{\sigma}(x - x_0 + \sigma) & x_0 - \sigma \le x \le x_0, \\ \frac{1}{\sigma}(x_0 - x + \beta) & x_0 \le x \le x_0 + \sigma, \\ 0 & otherwise . \end{cases}$$

26

The parametric form of symmetric triangular fuzzy number is

$$\underline{A}(\alpha) = x_0 - \sigma + \sigma \alpha$$
,  $\overline{A}(\alpha) = x_0 + \sigma - \sigma \alpha$ .

**Definition 2.5.** For fuzzy set A Support function is defined as follows:

$$supp(A) = \overline{\{x|\mu(x) > 0\}},$$

where  $\overline{\{x|\mu(x)>0\}}$  is closure of set  $\{x|\mu(x)>0\}$ .

**Definition 2.6.** [16], A function  $f : [0, 1] \to [0, 1]$  symmetric around  $\frac{1}{2}$ , i.e.  $f(\frac{1}{2} - \alpha) = f(\frac{1}{2} + \alpha)$  for all  $\alpha \in [0, \frac{1}{2}]$ , which reaches its minimum in  $\frac{1}{2}$ , is called the bi-symmetrical weighted function. Moreover, the bi-symmetrical weighted function is called regular if

- (1)  $f(\frac{1}{2}) = 0$ ,
- (2) f(0) = f(1) = 1,
- (3)  $\int_0^1 f(\alpha) d\alpha = \frac{1}{2}$ .

**Definition 2.7.** [14], For two arbitrary fuzzy numbers A and B with  $\alpha$ -cuts  $[f_A^{-1}(\alpha), g_A^{-1}(\alpha)]$ and  $[f_B^{-1}(\alpha), g_B^{-1}(\alpha)]$  respectively, the quantity

$$d(A,B) = \left[\int_0^1 f(\alpha)(f_A^{-1}(\alpha) - f_B^{-1}(\alpha))^2 d\alpha + \int_0^1 f(\alpha)(g_A^{-1}(\alpha) - g_B^{-1}(\alpha))^2 d\alpha\right]^{\frac{1}{2}}$$
(2.2)

is the weighted distance between A and B.

**Definition 2.8.** [4, 16], Let A is an arbitrary fuzzy number, the expected interval and expected value of a fuzzy number A are noted by EI(A) and EV(A) respectively, and considered as follows, (with  $f(\alpha) = \alpha$ ),

$$EI(A) = [E_1^A, E_2^A] = [2 \int_0^1 \alpha f_A^{-1}(\alpha) d\alpha, 2 \int_0^1 \alpha g_A^{-1}(\alpha) d\alpha],$$
  

$$EV(A) = \frac{E_1^A + E_2^A}{2} = \int_0^1 \alpha f_A^{-1}(\alpha) d\alpha + \int_0^1 \alpha g_A^{-1}(\alpha) d\alpha$$
(2.3)

If  $A = (a_1, a_2, a_3, a_4)$  is a trapezoidal fuzzy number then:

$$EI(A) = \left[\frac{a_1 + 2a_2}{3}, \frac{2a_3 + a_4}{3}\right],\tag{2.4}$$

$$EV(A) = \frac{1}{3}(a_1 + 2a_2 + 2a_3 + a_4).$$
(2.5)

**Proposition 2.1.** If A and B are two fuzzy numbers and  $\lambda, \nu \in \mathbb{R}$  then:

$$EI(\lambda A + \nu B) = \lambda EI(A) + \nu EI(B),$$
  

$$EV(\lambda A + \nu B) = \lambda EV(A) + \nu EV(B),$$
(2.6)

www.SID.ir

The addition and scalar multiplication of fuzzy numbers are defined by the extension principle and can be equivalently represented as follows. For arbitrary fuzzy numbers  $A = (\underline{A}, \overline{A})$  and  $B = (\underline{B}, \overline{B})$ , this article defines addition (A + B) and multiplication by scalar k > 0 as

$$(\underline{A} + \underline{B})(\alpha) = \underline{A}(\alpha) + \underline{B}(\alpha) \quad , \quad (\overline{A} + \overline{B})(\alpha) = \overline{A}(\alpha) + \overline{B}(\alpha), \tag{2.7}$$

$$(\underline{kA})(\alpha) = \underline{kA}(\alpha) \quad , \quad (\overline{kA})(\alpha) = \underline{kA}(\alpha).$$
(2.8)

To emphasis the collection of all fuzzy numbers with addition and multiplication as defined by (2.2) and (2.3) denoted by F, which is a convex cone.

#### 3 The proposed model

In this section, the researcher introduce ranking model based on weighted distance in data envelopment analysis. This article assumes that the  $DMU_p$  is extreme efficient [12, 6]. By omitting  $(X_p, Y_p)$  from  $T_c$  (PPS of CCR model), the researcher defines the production possibility set  $T'_c$  as follows, [7]:

$$T'_{c} = \{ (X,Y) \mid X \ge \sum_{j=1, j \neq p}^{n} w_{j}X_{j} , Y \le \sum_{j=1, j \neq p}^{n} w_{j}Y_{j}, w_{j} \ge 0, j = 1, \cdots, n, j \neq p \}.$$
(3.9)

Where

$$T_c = \{ (X, Y) \mid X \ge \sum_{j=1}^n w_j X_j , Y \le \sum_{j=1, j \ne p}^n w_j Y_j, w_j \ge 0, j = 1, \cdots, n \}.$$
(3.10)

To obtain the ranking score of  $DMU_p$ , this article considers the following model:

$$\min \quad \Gamma_{c}^{p}(X,Y) = \sum_{i=1}^{m} f(\alpha)(x_{i} - x_{ip})^{2} + \sum_{r=1}^{s} f(\alpha)(y_{r} - y_{rp})^{2}$$

$$s.t \quad \sum_{j=1, j \neq p}^{n} w_{j}x_{ij} \leq x_{i}, \qquad i = 1 \cdots, m,$$

$$\sum_{j=1, j \neq p}^{n} w_{j}y_{rj} \geq y_{r}, \qquad i = 1 \cdots, s,$$

$$x_{i} \geq 0, \qquad i = 1 \cdots, m,$$

$$y_{r} \geq 0, \qquad r = 1 \cdots, s,$$

$$w_{j} \geq 0, \qquad j = 1 \cdots, n, j \neq p.$$

$$(3.11)$$

Where  $X = (x_1, \ldots, x_n)$ ,  $Y = (y_1, \ldots, y_n)$  and  $\lambda = (\lambda_1, \ldots, \lambda_n)$  are the variables of the model (3.11) and  $\Gamma_c^p(X, Y)$  is the weighted distance  $(X_p, Y_p)$  from (X, Y) by weighted distance, also  $f(\alpha)$  is regular weighted function. Quadratic programming represents a special class of nonlinear programming in which the objective function is quadratic and the constraints are linear. The KKT conditions of a quadratic programming problem reduce to a linear complementary problem. Thus the complementary pivoting algorithm can be used for solving a quadratic programming problem.

28

### 4 Comparison In Fuzzy DEA

In this section, the researcher supposes that inputs and outputs of DMUs are fuzzy numbers. Therefore,

$$\tilde{T}'_{c} = \{ (X,Y) \mid X \ge \sum_{j=1, j \neq p}^{n} w_{j} \tilde{X}_{j}, Y \le \sum_{j=1, j \neq p}^{n} w_{j} \tilde{Y}_{j}, w_{j} \ge 0, j = 1, \cdots, n, j \neq p \}$$
(4.12)

Weighted distance model with Eqs. (2.2) can be extended to the following model:

$$\min \quad \Gamma_{c}^{p}(X,Y) = \sum_{i=1}^{m} \left( \int_{0}^{1} f(\alpha) (f_{x_{i}}^{-1}(\alpha) - f_{x_{ip}}^{-1}(\alpha))^{2} d\alpha + \int_{0}^{1} f(\alpha) (g_{x_{i}}^{-1}(\alpha) - g_{x_{ip}}^{-1}(\alpha))^{2} d\alpha \right)$$
$$+ \sum_{r=1}^{s} \left( \int_{0}^{1} f(\alpha) (f_{y_{r}}^{-1}(\alpha) - f_{y_{rp}}^{-1}(\alpha))^{2} d\alpha + \int_{0}^{1} f(\alpha) (g_{y_{r}}^{-1}(\alpha) - g_{y_{rp}}^{-1}(\alpha))^{2} d\alpha \right)$$
$$s.t \qquad (X,Y) \in \tilde{T}_{c}'.$$
(4.13)

Where  $X = (x_1, \ldots, x_n)$ ,  $Y = (y_1, \ldots, y_n)$  and  $\lambda = (\lambda_1, \ldots, \lambda_n)$  are the variables of the model (4.13), that all components of vectors X and Y for all DMUs are non-negative and each DMU has at least one strictly positive input and output.

For solving the model (4.13), we have some following definitions:

**Definition 4.1.** [8], For any pair of fuzzy numbers A and B the degree in A is bigger than B has the following form:

$$\mu_M(A,B) = \begin{cases} 0 & if \quad E_2^A - E_1^B < 0, \\ \frac{E_2^A - E_1^B}{E_2^A - E_1^A + E_2^B - E_1^B} & if \quad 0 \in [E_1^A - E_2^B, E_2^A - E_1^B], \\ 1 & if \quad E_1^A - E_2^B > 0. \end{cases}$$
(4.14)

Where  $[E_1^A, E_2^A]$  and  $[E_1^B, E_2^B]$  are the expected intervals of A and B. When  $\mu_M(A, B) = \frac{1}{2}$ , we will say that A and B are different. When  $\mu_M(A, B) \ge \alpha$ , we will say that A is bigger than, or equal to B at least in degree  $\alpha$  and we will represent it by  $A \ge_{\alpha} B$ .

**Definition 4.2.** Given a production possibility  $(X, Y) \in \tilde{T}'_c$ , we will say that it is product in degree  $\alpha$  in  $\tilde{T}'_c$  if:

$$min \left\{ \begin{array}{ccc} \mu_{M}(x_{i}, \sum_{j=1, j\neq p}^{n} w_{j} \tilde{x}_{ij}) &, & \mu_{M}(\sum_{j=1, j\neq p}^{n} w_{j} \tilde{y}_{rj}, y_{r}) \\ \\ \mu_{M}(x_{i}, \tilde{x}_{ip}) &, & \mu_{M}(\tilde{y}_{rp}, y_{r}) \\ & i = 1, \dots, m & r = 1, \dots, s \end{array} \right\} = \alpha.$$
(4.15)

That is to say

$$x_{i} \geq_{\alpha} \sum_{j=1, j \neq p}^{n} w_{j} \tilde{x}_{ij} , \quad i = 1, \dots, m,$$
  
$$y_{r} \leq_{\alpha} \sum_{j=1, j \neq p}^{n} w_{j} \tilde{y}_{rj} , \quad r = 1, \dots, s.$$
 (4.16)

www.SID.ir

30

R. Saneifard / IJIM Vol. 3, No. 1 (2011) 25-33

There is:

$$x_{i} \geq \sum_{j=1, j \neq p}^{n} w_{j} (\alpha E_{2}^{x_{ij}} + (1 - \alpha) E_{1}^{x_{ij}}) , \quad i = 1, \dots, m,$$

$$y_{r} \leq \sum_{j=1, j \neq p}^{n} w_{j} (\alpha E_{1}^{y_{rj}} + (1 - \alpha) E_{2}^{y_{rj}}) , \quad r = 1, \dots, s.$$
(4.17)

(For more details see [12]).

**Definition 4.3.** A production possibility  $(X_o, Y_o)^{\alpha} \in \tilde{T}'_c$  is an  $\alpha$ -acceptable optimal solution of model (4.13) if it is an optimal solution of the following model:

$$min \quad \Gamma_{c}^{p}(X,Y)_{\alpha} = \sum_{i=1}^{m} \left( \int_{0}^{1} f(\alpha) (f_{x_{i}}^{-1}(\alpha) - f_{x_{ip}}^{-1}(\alpha))^{2} d\alpha + \int_{0}^{1} f(\alpha) (g_{x_{i}}^{-1}(\alpha) - g_{x_{ip}}^{-1}(\alpha))^{2} d\alpha \right)$$
$$+ \sum_{r=1}^{s} \left( \int_{0}^{1} f(\alpha) (f_{y_{r}}^{-1}(\alpha) - f_{y_{rp}}^{-1}(\alpha))^{2} d\alpha + \int_{0}^{1} f(\alpha) (g_{y_{r}}^{-1}(\alpha) - g_{y_{rp}}^{-1}(\alpha))^{2} d\alpha \right)$$
$$s.t \qquad (X,Y) \in \tilde{T}_{c}^{\prime \alpha}.$$
(4.18)

Where

$$\tilde{T}_{c}^{\prime\alpha} = \{(X,Y) \mid X \ge_{\alpha} \sum_{j=1, j \neq p}^{n} w_{j} \tilde{X}_{j}, Y \le_{\alpha} \sum_{j=1, j \neq p}^{n} w_{j} \tilde{Y}_{j}, w_{j} \ge 0, j = 1, \cdots, n\}$$
(4.19)

**Proposition 4.1.** If  $\alpha_1 < \alpha_2$  then  $\tilde{T}_c^{\prime \alpha_2} \subseteq \tilde{T}_c^{\prime \alpha_1}$ .

We write model (4.18) as follows:

$$\min \quad \Gamma_{c}^{p}(X,Y)_{\alpha} = \sum_{i=1}^{m} \left( \int_{0}^{1} f(\alpha) (f_{x_{i}}^{-1}(\alpha) - f_{x_{ip}}^{-1}(\alpha))^{2} d\alpha + \int_{0}^{1} f(\alpha) (g_{x_{i}}^{-1}(\alpha) - g_{x_{ip}}^{-1}(\alpha))^{2} d\alpha \right)$$

$$+ \sum_{r=1}^{s} \left( \int_{0}^{1} f(\alpha) (f_{y_{r}}^{-1}(\alpha) - f_{y_{rp}}^{-1}(\alpha))^{2} d\alpha + \int_{0}^{1} f(\alpha) (g_{y_{r}}^{-1}(\alpha) - g_{y_{rp}}^{-1}(\alpha))^{2} d\alpha \right)$$

$$s.t \qquad x_{i} \geq \sum_{j=1, j \neq p}^{n} w_{j} (\alpha E_{2}^{x_{ij}} + (1 - \alpha) E_{1}^{x_{ij}}) \quad i = 1, \dots, m,$$

$$y_{r} \leq \sum_{j=1, j \neq p}^{n} w_{j} (\alpha E_{1}^{y_{rj}} + (1 - \alpha) E_{2}^{y_{rj}}) \quad r = 1, \dots, s,$$

$$y_{r} \geq 0 \qquad r = 1, \dots, s,$$

$$w_{j} \geq 0 \qquad j = 1, \dots, n.$$

$$(4.20)$$

Model (4.20) is a crisp  $\alpha$ -parametric model. Therefore this article can solve it by the interactive method. Now this study is going to explain the interactive method.

Table 1

Fuzzy data of DMUs in fuzzy data in Example 4.1.

## Archive of SID

R. Saneifard / IJIM Vol. 3, No. 1 (2011) 25-33

DMUs	Input	$lpha ext{-cut}$	Output	$lpha ext{-cut}$
A	(11, 12, 14)	$[11+\alpha, 14-\alpha]$	(10, 10, 10)	[10, 10]
В	$(30,\!30,\!30)$	[30, 30]	(12, 13, 14, 16)	$[12+\alpha, 16-2\alpha]$
C	(40, 40, 40)	[40, 40]	(11, 11, 11)	[11, 11]
D	(45, 47, 52, 55)	$^{[45+2\alpha,55\text{-}3\alpha]}$	(12, 15, 19, 22)	[12+3lpha,22-3lpha]

Table 2

The optimal values of $\alpha$ -parametric model.							
α	$\Gamma_c^A$	$\Gamma_c^B$	$\Gamma_c^C$	$\Gamma_c^D$	$(Ranking)_{\alpha}$		
0.0	25.5	3.5	0	55.2	$(C \prec B \prec A \prec D)_{0.0}$		
0.1	27.0	3.5	0	55.2	$(C \prec B \prec A \prec D)_{0.1}$		
0.2	28.5	3.5	0	55.2	$(C \prec B \prec A \prec D)_{0.2}$		
0.3	30.0	3.5	0	55.2	$(C \prec B \prec A \prec D)_{0.3}$		
0.4	31.6	3.5	0	55.2	$(C \prec B \prec A \prec D)_{0.4}$		
0.5	33.3	3.5	0	55.2	$(C \prec B \prec A \prec D)_{0.5}$		
0.6	35.0	3.5	0	55.2	$(C \prec B \prec A \prec D)_{0.6}$		
0.7	36.7	3.5	0	55.2	$(C \prec B \prec A \prec D)_{0.7}$		
0.8	38.6	3.5	0	55.2	$(C \prec B \prec A \prec D)_{0.8}$		
0.9	40.4	3.5	0	55.2	$(C \prec B \prec A \prec D)_{0.9}$		
1.0	42.3	3.5	0	55.2	$(C \prec B \prec A \prec D)_{1.0}$		

#### 4.1 Interactive Method

Regarding to proposition (4.1), to obtain the nearest (X, Y) of  $\tilde{T}'_c$  implies a lesser degree of production possibility. Then the decision-maker runs in to two conflicting objectives: to find the nearest (X, Y) and to improve the degree of production possibility. Following Kaufman [11] the researcher considers 11 scales, which allow for different choice of decision-maker idea in (4.20) model.

1:  $\alpha = 0.0$  unacceptable solution 2:  $\alpha = 0.1$  Practically unacceptable solution 3:  $\alpha = 0.2$  Almost unacceptable solution 4:  $\alpha = 0.3$  Very unacceptable solution 5:  $\alpha = 0.4$  Quite unacceptable solution 6:  $\alpha = 0.5$  Neither acceptable nor unacceptable solution 7:  $\alpha = 0.6$  Quite acceptable solution 8:  $\alpha = 0.7$  Very acceptable solution 9:  $\alpha = 0.8$  Almost acceptable solution 10:  $\alpha = 0.9$  Practically acceptable solution 11:  $\alpha = 1.0$  Completely acceptable solution

We choice the  $\alpha_0$  is the minimum acceptable degree with decision-maker idea. Then, this article solving the (4.20)  $\alpha$ -parametric model for each  $\alpha_k$  that  $k = 1, \ldots, (10 - 10\alpha_0)$ . We obtain the  $\alpha_k$ -acceptable optimal fuzzy value of objective function of original model (4.13) with  $\alpha_k$ -acceptable solution of model (4.20) in model (4.13).

#### 4.2 Example

**Example 4.1.** We will consider a simple example was introduced in [10] with its data listed in Table 1. These DMUs are evaluated by proposed model in 4.13 with different  $\alpha_k$ . The  $\alpha$ -parametric model is as follows:

$$\min \int_{0}^{1} \alpha (x - 11 - \alpha) d\alpha + 2 \int_{0}^{1} \alpha (y - 10) d\alpha + \int_{0}^{1} \alpha (x - 14 + 2\alpha) d\alpha$$

$$s.t \qquad x \ge 30w_{B} + 40w_{C} + w_{D}(53.5 - 7.5\alpha)$$

$$y \le w_{B}(15 - 2.5\alpha) + 11w_{C} + w_{D}(20.5 - 7\alpha)$$

$$(4.21)$$

$$x \ge 0,$$

$$y \ge 0,$$

$$w_{B}, w_{C}, w_{D} \ge 0.$$

The  $\alpha$ -parametric model for B, C and D can be showed similarly. The results is shown in Table 2.

## 5 Conclusion

In the present article a modified approach based on weighted distance is introduced for ranking of decision making units with fuzzy data. The method is based on the interactive method.  $\alpha$ -acceptable optimal solution of proposed model for  $\alpha \geq \frac{1}{2}$  is an acceptable solution. For any decision making unit, the score of ranking is obtained by solving  $\alpha$ -parametric model (4.20).

#### References

- R. E. Bellman, L. A. Zadeh, Decision making in a fuzzy environment, Managemant Science 17 (1970) 141 - 164.
- [2] A. Charnes, W. W. Cooper, Rhodes, Measuring the efficiency of decision making units, Eur. J. Operation Research 2 (1978) 429 - 444.
- [3] M. J. Farrell, The measurment of productive efficiency, Journal of the Royal Statistical Society 120 (1957) 253 - 281.
- [4] S. Heilpern, The expected valued of a fuzzy number, Fuzzy Sets and Systems 47 (1996) 81 - 86.
- [5] S. Heilpern, The expected value of a fuzzy number, Fuzzy Sets and Systems 47 (1992) 81 - 86.
- [6] G. R. Jahanshahloo, F. Hosseinzadeh Lotfi, T. Allahviranloo, Ranking DMUs with fuzzy data in DEA, International Journal of Contempt Mathematics Sciences 2 (2007) 203 - 211.
- [7] G. R. Jahanshahloo, F. Hosseinzadeh Lotfi, N. Shoja, G. Tohidi, Ranking by using  $l_2$ -Norm in in DEA, Applied Mathematics and Computation 153 (2004) 215 224.
- [8] M. Jimenez, Ranking fuzzy numbers through the comparison of its expected intervals, International Journal of Uncertainty, Fuzziness and Knowledge - Based Systems 4 (2004) 379 - 388. Mathematics. (2) 163 - 173.
- [9] C. Kao, S. T. Liu, A mathematical programming approach to fuzzy efficiency ranking, International Journal Production Economics 86 (2003) 45 - 54.
- [10] C. Kao, S. T. Liu, A mathematical programming approach to fuzzy efficiency ranking, International Journal Production Economic 86 (2003) 45 - 54.
- [11] A. Kauffman, M.M. Gupta, Introduction to Fuzzy Arithmetic: Theory and Application, Van Nostrand Reinhold, New York, 1991.
- [12] R. Saneifard, T. Allahviranloo, F. Hosseinzadeh, N. Mikaeilvand, Euclidean ranking DMUs with fuzzy data in DEA, Applied Mathematical Sciences 60 (2007) 2989 -2998.
- [13] R. Saneifard, Ranking L-R fuzzy numbers with weighted averaging based on levels, International Journal of Industrial Mathematics 2 (2009)163 - 173.
- [14] R. Saneifard, A method for defuzzification by weighted distance, International Journal of Industrial Mathematics 3 (2009) 209 - 217.
- [15] R. Saneifard, Defuzzification method for solving fuzzy linear systems, International Journal of Industrial Mathematics 4 (2009) 321 - 331.
- [16] R. Saneifard, Defuzzification Through a Bi Symmetrical weighted Function, Aust. J. Basic appl. sci. 10 (2010) 4976 - 4984.