

Linear optimization of fuzzy relation inequalities with max-Lukasiewicz composition

E. Shivanian ^{*†}

Abstract

In this paper, we study the finitely many constraints of fuzzy relation inequalities problem and optimize the linear objective function on this region which is defined with fuzzy max-Lukasiewicz operator. In fact Lukasiewicz t-norm is one of the four basic t-norms. A new simplification technique is given to accelerate the resolution of the problem by removing the components having no effect on the solution process. Also, an algorithm and one numerical example are offered to abbreviate and illustrate the steps of the problem resolution process.

Keywords : Linear objective function optimization; Fuzzy relation equations; Fuzzy relation inequalities; max-Lukasiewicz composition.

1 Introduction

Fuzzy relation equations (FRE), fuzzy relation inequalities (FRI) and their connected problems have been investigated by many researchers in both theoretical and applied areas [4, 5, 8, 11, 13, 18, 32, 33, 35, 42]. Sanchez [34] started a development of the theory and applications of FRE treated as a formalized model for non-precise concepts. Generally, FRE and FRI has a number of properties that make it suitable for formulizing the uncertain information upon which many applied concepts are usually based. The application of (FRE) and (FRI) can be seen in many areas, for instance, fuzzy control, fuzzy decision making, system analysis, fuzzy modeling, fuzzy arithmetic, fuzzy symptom, diagnosis, and especially fuzzy medical diagnosis and so on (see [1, 2, 5, 7, 8, 9, 23, 27, 30, 31, 32, 41, 44]).

An interesting extensively investigated kind

of such these problems is the optimization of the objective functions on the region whose feasible solutions sets have been defined as FRE or FRI constraints [3, 10, 14, 16, 18, 21, 22, 25, 26, 37, 38, 39, 40]. Fang and Li solved the linear optimization problem with respect to the FRE constraints by considering the max-min composition [10]. The max-min composition is commonly used when a system requires conservative solutions in sense that the goodness of one value can not compensate the badness of another value [21]. Recent results in the literature, however, show that the min operator is not always the best choice for intersection operation. Instead, the max-product composition provided results better or equivalent to the max-min composition in some application [1].

The fundamental result for fuzzy relation equations with max-product composition goes back to Pedrycz [30]. Recent study in this regard can be found in Bourk and Fisher [3]. They extended the study of an inverse solution of a system of fuzzy relation equations with max-product

*Corresponding author. shivanian@sci.ikiu.ac.ir

†Department of Mathematics, Imam Khomeini International University, Qazvin, 34149-16818, Iran.

composition. They provided theoretical results for determining the complete solution sets as well as the conditions for the existence of resolutions. Their results showed that such complete solution sets can be characterized by one maximum solution and a number of minimal solutions. Furthermore, the monograph by Di Nola, Sessa, Pedrycz and Sanchez [8] contains a thorough discussion of this class of equations. Nonetheless, recent published literatures show that max-L composition in which L is Lukasiewicz t-norm is played important role in applications [6, 7, 24, 28, 29, 36].

In this paper, we consider the linear optimization problem of the fuzzy relation inequalities (FRI) with max-Lukasiewicz operator which, we show max-L for simplicity [20]. This problem can be formulated as following:

$$\begin{aligned} & \min c^t x \\ \text{s.t.} \quad & A \circ_L x \geq d^1 \\ & B \circ_L x \leq d^2 \\ & x \in [0, 1]^n \end{aligned} \tag{1.1}$$

where $A = (a_{ij})_{m \times n}$, $a_{ij} \in [0, 1]$, $B = (b_{ij})_{l \times n}$, $b_{ij} \in [0, 1]$, are fuzzy matrices, $d^1 = (d_i^1)_{m \times 1} \in [0, 1]^m$, $d^2 = (d_i^2)_{l \times 1} \in [0, 1]^l$ are fuzzy vectors and, $x = (x_j)_{n \times 1} \in [0, 1]^n$ is unknown fuzzy vector and, $c = (c_j)_{n \times 1} \in R^n$ is vector of cost coefficients, and “ \circ_L ” denotes the fuzzy max-L operator. Problem (1.1) can be rewritten as following problem:

$$\begin{aligned} & \min c^t x \\ \text{s.t.} \quad & a_i \circ_L x \geq d_i^1, \quad i \in I^1 = \{1, 2, \dots, m\}, \\ & b_i \circ_L x \leq d_i^2, \quad i \in I^2 = \{1, 2, \dots, l\}, \\ & 0 \leq x_j \leq 1, \quad j \in J = \{1, 2, \dots, n\}, \end{aligned} \tag{1.2}$$

where a_i and b_i are i 'th row of the matrices A and B , respectively and the constraints are expressed by the max-L operator definition as:

$$\begin{aligned} & \forall i \in I^1 : \\ & a_i \circ_L x = \max_{j \in J} \{ \max(a_{ij} + x_j - 1, 0) \} \geq d_i^1 \\ & \forall i \in I^2 : \\ & b_i \circ_L x = \max_{j \in J} \{ \max(a_{ij} + x_j - 1, 0) \} \leq d_i^2, \end{aligned} \tag{1.3}$$

In Section 2, the feasible solutions set of the problem (1.2) and its properties are studied also, necessary and sufficient conditions are given to

realize the feasibility of the problem (1.2). In Section 2, some simplification operations are presented to accelerate the resolution process. Also, in Section 4 an algorithm is introduced to solve the problem and one example is given to illustrate the algorithm. Finally, a conclusion is stated in Section 5.

2 The characteristics of the feasible solution set

Definition 2.1 Define $S(A, d^1)_i = \{x \in [0, 1]^n : a_i \circ_L x \geq d_i^1\}$ for each $i \in I^1$, $S(B, d^2)_i = \{x \in [0, 1]^n : b_i \circ_L x \leq d_i^2\}$ for each $i \in I^2$, and $S(A, B, d^1, d^2) = S(A, d^1) \cap S(B, d^2) = \{x \in [0, 1]^n : A \circ_L x \geq d^1, B \circ_L x \leq d^2\}$.

Lemma 2.1 (a) $S(A, d^1) \neq \emptyset$ if and only if for each $i \in I^1$ there exists some $j \in J$ such that $d_i^1 \leq a_{ij}$.

(b) If $S(A, d^1) \neq \emptyset$ then $\bar{1} = [1, 1, \dots, 1]_{1 \times n}^t$ is the single maximum solution of $S(A, d^1)$.

Proof.

(a) Suppose $x \in S(A, d^1)$. Thus, $x \in S(A, d^1)_i$, $\forall i \in I^1$ by Definition 2.1, and thus, by Relation (1.3), for each $i \in I^1$ there are some $j_i \in J$ such that $\max(a_{ij_i} + x_{j_i} - 1, 0) \geq d_i^1$. Since for each $i \in I^1$ and $j \in J$ we have $d_i^1 \geq 0$ and $x_j \leq 1$, then $d_i^1 \leq a_{ij_i} + x_{j_i} - 1 \leq a_{ij_i}$ therefore for each $i \in I^1$ there exists some $j \in J$ such that $d_i^1 \leq a_{ij}$. Conversely, suppose there exist some $j_i \in J$ such that $d_i^1 \leq a_{ij_i}$, $\forall i \in I^1$. Set $x = \bar{1} = [1, 1, \dots, 1]_{1 \times n}^t$. Since $x \in [0, 1]^n$ and $\max_{j \in J} \{ \max(a_{ij_i} + x_{j_i} - 1, 0) \} \geq$

$\max(a_{ij_i} + x_{j_i} - 1, 0) \geq a_{ij_i} + x_{j_i} - 1 = a_{ij_i} \geq d_i^1$, $\forall i \in I^1$ hence, by Relation (1.3), $x \in S(A, d^1)_i$, $\forall i \in I^1$ and as a result $x \in S(A, d^1)$.

(b) Proof of this part is easily attained from the part (a) and this fact that $x_j \leq 1, \forall j \in J$.

Lemma 2.2 (a) $S(B, d^2) \neq \emptyset$

(b) The single minimum solution of $S(B, d^2)$ is $\bar{0} = [0, 0, \dots, 0]_{1 \times n}^t$.

Proof.

Set $x = \bar{0} = [0, 0, \dots, 0]_{1 \times n}^t$. We select $i \in I^2$ arbitrary and constant hereafter. Since, $0 \leq b_{ij}, d_i^2 \leq 1$ we have $b_{ij} + x_j - 1 \leq d_i^2, \forall j \in J$ then $\max(b_{ij} + x_j - 1, 0) \leq d_i^2, \forall j \in J$ and hence $\max_{j \in J} \{ \max(b_{ij} + x_j - 1, 0) \} \leq d_i^2$ therefore $x \in S(B, d^2)$ and then part (a) and (b) are proved.

Theorem 2.1 (Necessary condition)

If $S(A, B, d^1, d^2) \neq \emptyset$ then, $\forall i \in I^1 \exists j \in J$ Such that $d_i^1 \leq a_{ij}$.

Proof.

This Theorem is clearly proved from Lemmas 2.1 and 2.2 and Definition 2.1.

Definition 2.2 Set $\bar{x} = (\bar{x}_j)_{n \times 1}$, where $\bar{x}_j = \min_{i \in I^2} \{ \min\{1 + d_i^2 - b_{ij}, 1\} \} = \min\{1, \min_{i \in I^2} \{1 + d_i^2 - b_{ij}\} \}$.

Lemma 2.3 \bar{x} is the single maximum solution of $S(B, d^2)$.

Proof.

Suppose $x \in S(B, d^2)$ then, $x \in S(B, d^2)_i, \forall i \in I^2$, and then, $\max_{j \in J} \{ \max(b_{ij} + x_j - 1, 0) \} \leq d_i^2, \forall i \in I^2$ by definition 2.1 and Relation 3, therefore $b_{ij} + x_j - 1 \leq d_i^2, \forall i \in I^2$ and $\forall j \in J$, and hence, for each $j \in J$, we have $x_j \leq 1 + d_i^2 - b_{ij}, \forall i \in I^2$, and then, $x_j \leq \min_{i \in I^2} \{1 + d_i^2 - b_{ij}\}, \forall j \in J$. By the way, since $x_j \leq 1, \forall j \in J$, therefore we have $x_j \leq \min_{i \in I^2} \{1 + d_i^2 - b_{ij}\} = \bar{x}_j$, and then $x \leq \bar{x}$, because of being arbitrary $j \in J$, and the proof is completed.

Theorem 2.2

$$S(B, d^2) = [\bar{0}, \bar{x}].$$

Proof.

It is clearly proved from part (b) of Lemma 2.2 and Lemma 2.3.

Definition 2.3 Let $J_i = \{j \in J : d_i^1 \leq a_{ij}\}, \forall i \in I^1$. For each $j \in J_i$, we define $i_{x(j)} = (i_{x(j)_k})_{n \times 1}$ such that

$$i_{x(j)_k} = \begin{cases} 1 + d_i^1 - a_{ij} & k = j \\ 0 & k \neq j \end{cases}$$

Lemma 2.4 Assume $i \in I^1$ is a fixed number.

(a) For each $j \in J_i$, the vectors $i_{x(j)}$ are the minimal solutions of $S(A, d^1)_i$.

(b) If $d_i^1 = 0$ then $\bar{0}$ is the single minimum solution of $S(A, d^1)_i$.

Proof.

(a) Suppose $j \in J_i$ and $i \in I^1$, since $i_{x(j)_j} = 1 + d_i^1 - a_{ij}$ then, $i_{x(j)} \in S(A, d^1)_i$ through relation 3. Now by contrary, let there exist $x \in S(A, d^1)_i$ such that $x < i_{x(j)}$, as a result $x_j < 1 + d_i^1 - a_{ij}$ and $x_k = 0$ for $k \in J - \{j\}$. After that $a_{ij} + x_j - 1 < d_i^1, \forall j \in J$ and followed that $x \notin S(A, d^1)_i$, by means of Relation 1.3, that is a contradiction.

(b) The proof of this part of lemma is clear because the one of the minimal solutions will be $i_{x(j)} = \bar{0}$.

Corollary 2.1 If $S(A, d^1)_i \neq \emptyset$, then $S(A, d^1)_i = \bigcup_{j \in J_i} [i_{x(j)}, \bar{1}]$, where $i \in I^1$.

Proof.

Take into account $S(A, d^1)_i \neq \emptyset$ means the vector $\bar{1}$ is the maximum solution and the vectors $i_{x(j)}, \forall j \in J_i$ are the minimal solutions in $S(A, d^1)_i$ as a result of Lemmas 2.1 and 2.4, respectively. Now, let $x \in \bigcup_{j \in J_i} [i_{x(j)}, \bar{1}]$, so, for

some $j \in J_i, x \in [i_{x(j)}, \bar{1}]$ and also $x \in [0, 1]^n$ and $x_j \geq i_{x(j)_j} = 1 + d_i^1 - a_{ij}$ via Definition 2.3, Hence, $x \in S(A, d^1)_i$ through Relation 1.3. Conversely, let $x \in S(A, d^1)_i$. Then there exists some $j' \in J$ such that $x_{j'} \geq 1 + d_i^1 - a_{ij'}$ as a result of Relation 1.3. Since, $x \in [0, 1]^n$ so, $1 + d_i^1 - a_{ij'} \leq 1$, then $d_i^1 \leq a_{ij'}$, and for that reason $j' \in J_i$. Therefore, $i_{x(j')} \leq x \leq \bar{1}$ that implies $x \in \bigcup_{j \in J_i} [i_{x(j)}, \bar{1}]$.

Definition 2.4 Let $e = (e(1), e(1.2), \dots, e(m)) \in J_1 \times J_2 \times \dots \times J_m$ such that $e(i) = j \in J_i$. We define $x(e)_j = \max_{i \in I_j^e} \{1 + d_i^1 - a_{ij}\}$ if $I_j^e \neq \emptyset$ and $x(e)_j = 0$ if $I_j^e = \emptyset$, where $I_j^e = \{i \in I^1 : e(i) = j\}$.

Lemma 2.5 Let $S(A, d^1) \neq \emptyset$, then $S(A, d^1) = \bigcup_{x(e) \in X(e)} [x(e), \bar{1}]$, where $X(e) = \{x(e) : e \in J_I\}$.

Proof.

If $S(A, d^1) \neq \emptyset$ then, $S(A, d^1)_i \neq \emptyset, \forall i \in I^1$. Hence, by Corollary 2.1 and Definitions 2.1 and 2.4, we have

$$\begin{aligned} S(A, d^1) &= \bigcap_{i \in I^1} S(A, d^1)_i = \\ &= \bigcap_{i \in I^1} \left[\bigcup_{j \in J_i} [i_{x(j)}, \bar{1}] \right] = \bigcap_{i \in I^1} \left[\bigcup_{e(i) \in J_i} [i_{x(e(i))}, \bar{1}] \right] \\ &= \bigcup_{e \in J_I} \left[\bigcap_{i \in I^1} [i_{x(e(i))}, \bar{1}] \right] = \bigcup_{e \in J_I} [x(e), \bar{1}] \\ &= \bigcup_{x(e) \in X(e)} [x(e), \bar{1}] \end{aligned} \tag{2.4}$$

From Lemma 2.5, it is obvious that $S(A, d^1) = \bigcup_{x(e) \in X_0(e)} [x(e), \bar{1}]$ and $X_0(e) = S_0(A, d^1)$, where $X_0(e)$ and $S_0(A, d^1)$ are the set of minimal solutions of $X(e)$ and $S(A, d^1)$, respectively.

Corollary 2.2 (a) If $d_i^1 = 0$ for $i \in I^1$, then we can remove the i 'th row of the matrix A .

(b) If $j \notin J_i, \forall i \in I^1$ then we can omit j 'th column of the matrix A for the purpose of finding $x(e)$.

Proof.

(a) It is proved from Definition 2.4 and the part (b) of the Lemma 2.4, because we will get minimal elements of $S(A, d^1)$

(b) It is proved only by using Definition 2.4.

It is recalled that in part (a), by Definition 2.4 and the part (b) of the Lemma 2.4, the i 'th row of the matrix A has no effect in the calculation of the vectors $x(e)$ belong to $X_0(e) = S_0(A, d^1)$, and also in part (b), before calculating the vectors $x(e), \forall e \in I_J$, we can remove j 'th column of the matrix A by the use of Definition 2.4 and set $x(e)_j = 0$.

Theorem 2.3 If $S(A, B, d^1, d^2) \neq \emptyset$, then $S(A, B, d^1, d^2) = \bigcup_{x(e) \in X_0(e)} [x(e), \bar{x}]$.

Proof.

It is obvious from Definition 2.1, Theorem 2.2 and Lemma 2.5.

Corollary 2.3 (Necessary and Sufficient Condition)

$S(A, B, d^1, d^2) \neq \emptyset$ if only if $\bar{x} \in S(A, d^1)$ or, equivalently, $S(A, B, d^1, d^2) \neq \emptyset$ if only if there exists some $e \in J_I$ such that $x(e) \leq \bar{x}$.

Proof:

It is clearly resulted from Theorem 2.2, Lemma 2.2 and Lemma 2.3.

3 Simplification operations and resolution algorithm

In order to solve the problem (1.2), it is initially converted into two follow sub-problems

$$\begin{aligned} & \min c^{+t}x \\ \text{s.t. } & A \circ_L x \geq d^1 \\ & B \circ_L x \leq d^2 \\ & x \in [0, 1]^n \end{aligned} \tag{4a}$$

$$\begin{aligned} & \min c^{-t}x \\ \text{s.t. } & A \circ_L x \geq d^1 \\ & B \circ_L x \leq d^2 \\ & x \in [0, 1]^n \end{aligned} \tag{4b}$$

where, $c_j^+ = \max(0, c_j)$ and $c_j^- = \min(0, c_j)$.

It is understandable that \bar{x} is an optimal solution of (4b). Also, (4a) achieves its optimal points at

some $x(e) \in X_0(e)$. Once $x(e_0)$ optimizes (4a), we set $x^* = (x_j^*)_{n \times 1}$ such that

$$x_j^* = \begin{cases} \bar{x}_j & , c_j \leq 0 \\ x(e_0)_j & , c_j > 0 \end{cases}$$

Now following lemma gives us an optimal point of the problem (1.2).

Lemma 3.1 x^* is an optimal solution of the problem (1.2).

Proof. See the Theorem 2.1 in [14].

In order to calculate x^* , it is enough to find \bar{x} and $x(e_0)$. Although \bar{x} is easily attained through Definition 2.2, but $x(e_0)$ is not so, because, $X_0(e)$ is attained by pairwise comparison of $X(e)$ members. For that reason, having complete set of $X_0(e)$ is time-consuming, especially, while $X(e)$ has several members. Therefore, simplification operations can hasten the resolution of the problem (4a). With the intention of simplification the vectors $e \in J_I$ is removed at what time $x(e)$ is not optimal of (4a). One of such these operations is given by Corollary 2.2. Other operations are attained by follow theorems.

Definition 3.1 Let $\bar{J}_i = \{j \in J_i : 1 + d_i^1 - a_{ij} \leq \bar{x}_j\}, \forall i \in I^1$ where \bar{x} comes from Definition 2.2.

Theorem 3.1 $S(A, B, d^1, d^2) \neq \emptyset$ if only if $\bar{J}_i \neq \emptyset, \forall i \in I^1$.

Proof.

Suppose $S(A, B, d^1, d^2) \neq \emptyset$. Therefore by Corollary 2.3, $\bar{x} \in S(A, B, d^1, d^2)$ and so we have $\bar{x} \in S(A, d^1)_i, \forall i \in I^1$. Thus, for each $i \in I^1$ there exists some $j \in J$ such that $\bar{x}_j \geq 1 + d_i^1 - a_{ij}$, as a result of Corollary 2.12.1, consequently $\bar{J}_i \neq \emptyset, \forall i \in I^1$. Conversely, suppose $\bar{J}_i \neq \emptyset, \forall i \in I^1$. It means that, $\forall i \in I^1$ there exists some $j \in J$ such that $\bar{x}_j \geq 1 + d_i^1 - a_{ij}$. Hence, $\bar{x} \in S(A, d^1)_i, \forall i \in I^1$ through Corollary 2.1, as a result $\bar{x} \in S(A, d^1)$. This fact go with Lemma 2.3 implies $\bar{x} \in S(A, B, d^1, d^2)$, therefore, $S(A, B, d^1, d^2) \neq \emptyset$.

Theorem 3.2 Let $S(A, B, d^1, d^2) \neq \emptyset$, then $S(A, B, d^1, d^2) = \bigcup_{x(e) \in \bar{X}(e)} [x(e), \bar{x}]$ where, $\bar{X}(e) = \{x(e) : e \in \bar{J}_I = \bar{J}_1 \times \bar{J}_2 \times \dots \times \bar{J}_m\}$.

Proof.

By considering Theorem 3.1, it is sufficient to show $x(e) \notin S(A, B, d^1, d^2)$ once $e \notin \bar{J}_I$. Suppose $e \notin \bar{J}_I$. Thus, there exist $i' \in I^1$ and

$j' \in J_{i'}$ such that $e(i') = j'$ and $1 + d_{i'}^1 - a_{i'j'} > \bar{x}_{j'}$, Then $i' \in I_{j'}^e$ and by means of Definition 2.4, $x(e)_{j'} = \max_{i \in I_{j'}^e} \{1 + d_i^1 - a_{ij'}\} \geq 1 + d_{i'}^1 - a_{i'j'} > \bar{x}_{j'}$. Therefore, $x(e) \leq \bar{x}$ will not be correct, and as a consequence of Theorem 3.13.1, we can obtain $x(e) \notin S(A, B, d^1, d^2)$.

It is noticeable that as a result of Definition 3.1, we have $\bar{J}_i \subseteq J_i, \forall i \in I^1$ that means $\bar{X}(e) \subseteq X(e)$. Also, by Theorem 3.2, $S_0(A, B, d^1, d^2) \subseteq \bar{X}(e)$ in which $S_0(A, B, d^1, d^2)$ is minimal elements of $S(A, B, d^1, d^2)$. Thus, the region of search can be reduced to find the set $S_0(A, B, d^1, d^2)$.

Definition 3.2 Let $J_i^* = \{j \in \bar{J}_i : c_j^- \neq 0\}, \forall i \in I^1$.

Theorem 3.3 Suppose $x(e_0)$ is an optimal solution in (4a) and $J_{i'}^* \neq \emptyset$ for some $i' \in I^1$, then there exist $x(e')$ such that $e'(i') \in J_{i'}^*$ and also, $x(e')$ is the optimal solution in (4a).

Proof.

Suppose $J_{i'}^* \neq \emptyset$ for some $i' \in I^1$ and $e_0(i') = j'$. Define $e' \in \bar{J}_I$ such that $e'(i') = k \in J_{i'}^*$ and $e'(i) = e_0(i)$ for each $i \in I^1$ and $i \neq i'$. By means of Definition 2.4, we have

$$\begin{aligned} x(e_0)_{j'} &= \max_{i \in I_{j'}^{e_0}} \{1 + d_i^1 - a_{ij'}\} \\ &\geq \max_{i \in I_{j'}^{e_0} \& i \neq i'} \{1 + d_i^1 - a_{ij'}\} = x(e')_{j'} \end{aligned}$$

and $x(e_0)_j = x(e')_j$ for each $j \in J$ and $j \neq j', k$. Therefore, with noting $c_k^+ = 0$ we have:

$$\begin{aligned} c^{+t}x(e_0) &= c_{j'+x(e_0)_{j'}} + \\ &\sum_{j \in J \& j \neq j'} c_{j+x(e_0)_j} \geq c_{j'+x(e')_{j'}} + \\ &\sum_{j \in J \& j \neq j'} c_{j+x(e')_j} = c^{+t}x(e') \end{aligned}$$

Therefore $x(e')$ is an optimal solution in (4a) then, proof is completed.

Corollary 3.1 If $J_i^* \neq \emptyset$ for some $i \in I^1$ then by omitting i 'th row we reach a reduced problem for which each optimal solution is an optimal solution for the previous (main) problem.

Proof.

It is resulted from Theorem 3.3 and also, notes that $c_j^+ = 0$ for each $j \in J_i^*$.

Definition 3.3 Let $j_1, j_2 \in J, c_{j_1} > 0$ and $c_{j_2} > 0$. We say j_2 dominates j_1 if only if

- (a) $j_1 \in \bar{J}_i$ implies $j_2 \in \bar{J}_i, \forall i \in I^1$.
- (b) For each $i \in I^1$ such that $j_1 \in \bar{J}_i, c_{j_1}(1 + d_i^1 - a_{ij_1}) \geq c_{j_2}(1 + d_i^1 - a_{ij_2})$.

Theorem 3.4 Suppose $x(e_0)$ is the optimal in (4a) and j_2 dominates j_1 for $j_1, j_2 \in J$, then, there exist $x(e')$ such that $I_{j_1}^{e'} = \emptyset$ and also, $x(e')$ is an optimal solution in (4a).

Proof. Define $e' = (e'(i))_{m \times 1}$ such that

$$e'(i) = \begin{cases} e_0(i) & i \notin I_{j_1}^{e_0} \\ j_2 & i \in I_{j_1}^{e_0} \end{cases}$$

It is obvious that $I_{j_1}^{e'} = \emptyset$ and so $x(e')_{j_1} = 0$. Also, $x(e_0)_j = x(e')_j$ for each $j \in J$ and $j \neq j_1, j_2$, $x(e')_{j_2} = 1 + d_{i_0}^1 - a_{i_0j_2}$. Now, if $i_0 \notin I_{j_1}^{e_0}$ then:

$$x(e_0)_{j_2} = x(e')_{j_2} = 1 + d_{i_0}^1 - a_{i_0j_2},$$

so we have

$$\begin{aligned} c^{+t}x(e_0) &= c_{j_1}^+x(e_0)_{j_1} + \sum_{j \in J \& j \neq j_1} c_{j+x(e_0)_j} \\ &\geq \sum_{j \in J \& j \neq j_1} c_{j+x(e')_j} = c^{+t}x(e') \end{aligned}$$

That proof is completed in this case. Otherwise, assume $i_0 \in I_{j_1}^{e_0}$. We show $c^{+t}x(e_0) \geq c^{+t}x(e')$. As a result of definition 2.4, let $x(e_0)_{j_2} = 1 + d_{i_0}^1 - a_{i_0j_2}$. Therefore, we have $c_{j_2}^+x(e_0)_{j_2} \geq 0$ by Definition 3.3. Consequently, since

$$c^{+t}x(e_0) = c_{j_1}^+x(e_0)_{j_1} + c_{j_2}^+x(e_0)_{j_2}$$

$$+ \sum_{j \neq j_1, j_2} c_j^+x(e_0)_j,$$

and

$$c^{+t}x(e') = c_{j_2}^+x(e')_{j_2} + \sum_{j \neq j_1, j_2} c_j^+x(e')_j$$

It is sufficient to show $c_{j_1}^+x(e_0)_{j_1} \geq c_{j_2}^+x(e')_{j_2}$. Now, by definition 2.4, set

$$x(e_0)_{j_1} = 1 + d_{i'}^1 - a_{i'j_1}$$

Since j_2 dominates j_1 , so we have

$$c_{j_1}^+(1 + d_{i'}^1 - a_{i'j_1}) \geq c_{j_2}^+(1 + d_{i_0}^1 - a_{i_0j_2})$$

That means $c_{j_1}^+x(e_0)_{j_1} \geq c_{j_2}^+x(e')_{j_2}$ once $i_0 = i'$. Otherwise, suppose $i_0 \neq i'$. Since $i_0 \in I_{j_1}^{e_0}$ and j_2 dominates j_1 , thus

$$c_{j_1}^+(1 + d_{i_0}^1 - a_{i_0j_1}) \geq c_{j_2}^+(1 + d_{i_0}^1 - a_{i_0j_2})$$

Also, through definition 2.4 we have $x(e_0)_{j_1} = \max_{i \in I_{j_1}^{e_0}} \{1 + d_i^1 - a_{ij_1}\} = 1 + d_{i'}^1 - a_{i'j_1}$ that implies

$$1 + d_{i'}^1 - a_{i'j_1} \geq 1 + d_i^1 - a_{ij_1}, \forall i \in I_{j_1}^{e_0}$$

Therefore

$$\begin{aligned} c_{j_1}^+(1 + d_{i'}^1 - a_{i'j_1}) &\geq c_{j_1}^+(1 + d_{i_0}^1 - a_{i_0j_1}) \\ &\geq c_{j_2}^+(1 + d_{i_0}^1 - a_{i_0j_2}) \end{aligned}$$

That results $c_{j_1}^+x(e_0)_{j_1} \geq c_{j_2}^+x(e')_{j_2}$, hence $c^{+t}x(e_0) \geq c^{+t}x(e')$, therefore proof is completed.

Corollary 3.2 If j_2 dominates j_1 for $j_1, j_2 \in J$, then, by omitting j_1 'th column we reach a reduced problem for which each optimal solution is an optimal solution for the previous (main) problem.

4 An algorithm for finding an optimal solution and example

Definition 4.1 Consider the Problem (1.1). We call $\bar{A} = (\bar{a}_{ij})_{m \times n}$ and $\bar{B} = (\bar{b}_{ij})_{l \times n}$ the characteristic matrices of the matrix A and matrix B , respectively, where $\bar{a}_{ij} = 1 + d_i^1 - a_{ij}$ for each $i \in I^1$ and $j \in J$, also $\bar{b}_{ij} = 1 + d_i^2 - b_{ij}$ for each $i \in I^2$ and $j \in J$.

Algorithm: Given problem (1.2),

1. Find the matrices \bar{A} and \bar{B} by Definition 2.4.
2. If there exists $i \in I^1$ such that $a_{ij} < d_i^1$, $\forall j \in J$ then stop. Problem 1.2 is infeasible (see Theorem 2.1).
3. Calculate \bar{x} from \bar{B} by Definition 2.2.
4. If there exists $i \in I^1$ such that $d_i^1 = 0$ then remove i 'th row of the matrix \bar{A} (see the part (a) of the Corollary 2.2).
5. If $\bar{a}_{ij} > \bar{x}_j$ then, set $\bar{a}_{ij} = 0 \forall i \in I^1$ and $\forall j \in J$.
6. If there exists $i \in I^1$ such that $\bar{a}_{ij} = 0, \forall j \in J$ then stop. Problem (1.2) is infeasible (see Theorem 3.1 and 3.2)
7. If there exists $j' \in J$ such that, $\bar{a}_{ij'} = 0, \forall i \in I^1$ then remove j' th column of the matrix \bar{A} (see Theorem 3.2) and set $x(e_0)_{j'} = 0$.

8. For each $i \in I^1$, if $J_i^* \neq \emptyset$ then remove i 'th row of the matrix \bar{A} (see Corollary 3.1).
9. Remove each column $j \in J$ from \bar{A} such that $c_j < 0$ and set $x(e_0)_j = 0$.
10. If j_2 dominates j_1 then remove the column j_1 from \bar{A} , $\forall j_1, j_2 \in J$ (see Corollary 3.2) and set $x(e_0)_{j_1} = 0$.
11. Let $J_i^{new} = \{j \in \bar{J}_i : \bar{a}_{ij} \neq 0\}$ and $J_I^{new} = J_1^{new} \times J_2^{new} \times \dots \times J_m^{new}$. Find the vectors $x(e)$, $\forall e \in J_I^{new}$ by Definition 2.4 from \bar{A} , and $x(e_0)$ by pairwise comparison between the vectors $x(e)$.
12. Find x^* via Lemma 3.1.

5 Numerical example

Consider the problem in below:

$$\begin{aligned} \min \quad & 2x_1 - x_2 - 3x_3 + 2.5x_4 - x_5 \\ & + 6x_6 - 3x_7 + 2x_8 + x_9 + 5x_{10} \end{aligned}$$

$$\begin{bmatrix} 1 & 0.16 & 0.37 & 0.95 & 0.17 & 0.07 \\ 0.08 & 0.51 & 0.26 & 0.1 & 0.3 & 0.4 \\ 0.99 & 0.59 & 0.28 & 0.34 & 0.34 & 0.74 \\ 0.83 & 0.75 & 0.25 & 0.35 & 0.2 & 0.5 \\ 0.73 & 0.84 & 0.94 & 0.44 & 0.54 & 0.84 \\ 0.37 & 0.7 & 0.55 & 0.4 & 0.2 & 0.2 \end{bmatrix}$$

$$\begin{bmatrix} 0.77 & 0.14 & 0.8 & 0.6 \\ 0.3 & 0.35 & 0.9 & 1 \\ 0.19 & 0.21 & 0.7 & 0.65 \\ 0.2 & 0.2 & 0.95 & 0.85 \\ 0.99 & 0.44 & 0.1 & 0.5 \\ 0.73 & 0.24 & 0.98 & 0.9 \end{bmatrix} \circ_L \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \\ x_7 \\ x_8 \\ x_9 \\ x_{10} \end{bmatrix}$$

$$\geq \begin{bmatrix} 0 \\ 0.5 \\ 0.57 \\ 0.6 \\ 0.72 \\ 0.6 \end{bmatrix}$$

$$\begin{bmatrix} 0 & 0.94 & 0.69 & 0.49 & 0.5 & 0.51 \\ 0.02 & 1 & 0.82 & 0.59 & 0.89 & 0.76 \\ 0.1 & 0.73 & 0.4 & 0.37 & 0.45 & 0.7 \\ 0.1 & 0.1 & 0.3 & 0.2 & 0.25 & 0.49 \end{bmatrix}$$

$$\begin{bmatrix} 0.87 & 0.43 & 0.2 & 0.5 \\ 0.74 & 0.52 & 0.56 & 0.36 \\ 0.69 & 0.2 & 0.95 & 0.85 \\ 0.58 & 0.01 & 0.8 & 0.98 \end{bmatrix} \circ_L \begin{bmatrix} x_1 \\ x_2 \\ x_3 \\ x_4 \\ x_5 \\ x_6 \\ x_7 \\ x_8 \\ x_9 \\ x_{10} \end{bmatrix}$$

$$\leq \begin{bmatrix} 0.7 \\ 0.76 \\ 0.6 \\ 0.5 \end{bmatrix}$$

$$x_j \in [0, 1]^n$$

Step 1:

The matrices \bar{A} and \bar{B} are as following

$$\bar{A} = \begin{bmatrix} 0 & 0.84 & 0.63 & 0.05 & 0.83 \\ 1.42 & 0.99 & 1.24 & 1.4 & 1.2 \\ 0.58 & 0.98 & 1.29 & 1.23 & 1.23 \\ 0.77 & 0.85 & 1.35 & 1.25 & 1.4 \\ 0.99 & 0.88 & 0.78 & 1.28 & 1.18 \\ 1.23 & 0.9 & 1.05 & 1.2 & 1.4 \\ 0.93 & 0.23 & 0.86 & 0.2 & 0.4 \\ 1.1 & 1.2 & 1.15 & 0.6 & 0.5 \\ 0.83 & 1.38 & 1.36 & 0.87 & 0.92 \\ 1.1 & 1.4 & 1.4 & 0.65 & 0.75 \\ 0.88 & 0.73 & 1.28 & 1.62 & 1.22 \\ 1.4 & 0.87 & 1.36 & 0.62 & 0.7 \end{bmatrix}$$

$$\bar{B} = \begin{bmatrix} 1.7 & 0.76 & 1.01 & 1.21 & 1.2 \\ 1.74 & 0.76 & 0.94 & 1.17 & 0.87 \\ 1.5 & 0.87 & 1.2 & 1.23 & 1.15 \\ 1.4 & 1.4 & 1.2 & 1.3 & 1.25 \\ 1.19 & 0.83 & 1.27 & 1.5 & 1.2 \\ 1 & 1.02 & 1.24 & 1.2 & 1.4 \\ 0.9 & 0.91 & 1.4 & 0.65 & 0.75 \\ 1.01 & 0.92 & 1.49 & 0.7 & 0.52 \end{bmatrix}$$

Step 2:

There is no $i \in I^1$ such that $a_{ij} < d_i^1, \forall j \in J$ therefore we can go to step 3.

Step 3:

$$\bar{x} = \begin{bmatrix} 1 & 0.76 & 0.94 & 1 & 0.87 \\ 0.9 & 0.83 & 1 & 0.65 & 0.52 \end{bmatrix}$$

Step 4:

Since $d_1^1 = 0$, then first row from matrix \bar{A} is removed.

Step 5:

In according to this step, \bar{A} is converted to as following:

$$\bar{A} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0 \\ 0.58 & 0 & 0 & 0 & 0 \\ 0.77 & 0 & 0 & 0 & 0 \\ 0.99 & 0 & 0.78 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.6 & 0.5 \\ 0.83 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.65 & 0 \\ 0.88 & 0.73 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.62 & 0 \end{bmatrix}$$

Step 6:

There is no $i \in I^1$ such that $\bar{a}_{ij} = 0, \forall j \in J$ therefore we can go to step 7.

Step 7:

The second, fourth, fifth and eighth columns in according with this step are removed and we have $x(e_0)_2 = x(e_0)_4 = x(e_0)_5 = x(e_0)_8 = 0$, by the way matrix \bar{A} is converted to following:

$$\bar{A} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0.6 & 0.5 \\ 0.58 & 0 & 0.83 & 0 & 0 & 0 \\ 0.77 & 0 & 0 & 0 & 0.65 & 0 \\ 0.99 & 0.78 & 0.88 & 0.73 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0.62 & 0 \end{bmatrix}$$

Step 8:

Since $J_5^* \neq \emptyset$, then we can delete fifth row, then we get to

$$\bar{A} = \begin{bmatrix} 0 & 0 & 0 & 0 & 0.6 & 0.5 \\ 0.58 & 0 & 0.83 & 0 & 0 & 0 \\ 0.77 & 0 & 0 & 0 & 0.65 & 0 \\ 0 & 0 & 0 & 0 & 0.62 & 0 \end{bmatrix}$$

Step 9:

Since $c_3, c_7 < 0$ then, we can remove third and seventh columns and we get to

$$\bar{A} = \begin{bmatrix} 0 & 0 & 0.6 & 0.5 \\ 0.58 & 0.83 & 0 & 0 \\ 0.77 & 0 & 0.65 & 0 \\ 0 & 0 & 0.62 & 0 \end{bmatrix}$$

Also, we have $x(e_0)_3 = x(e_0)_7 = 0$.

Step10:

In the attained matrix, first and ninth columns dominate sixth and tenth columns, respectively. By removing sixth and tenth columns, matrix \bar{A} is converted to

$$\bar{A} = \begin{bmatrix} 0 & 0.6 \\ 0.58 & 0 \\ 0.77 & 0.65 \\ 0 & 0.62 \end{bmatrix}$$

Also, we have $x(e_0)_6 = x(e_0)_{10} = 0$.

Step11:

In the new matrix, we have $J_2^{new} = \{9\}$, $J_3^{new} = \{1\}$, $J_4^{new} = \{1, 9\}$ and $J_6^{new} = \{9\}$. For $e_1 = (9, 1, 1, 9)$, $x(e_1)_1 = 0.77$ and $x(e_1)_9 = 0.62$, then

$$x(e_1) = (0.77, 0, 0, 0, 0, 0, 0, 0, 0.62, 0)$$

Also, $e_2 = (9, 1, 9, 9)$ results in $x(e_2)_1 = 0.58$ and $x(e_2)_9 = 0.65$, then

$$x(e_2) = (0.58, 0, 0, 0, 0, 0, 0, 0, 0.65, 0)$$

Therefore minimal solutions are $x(e_1)$ and $x(e_2)$. Since $c^{+t}x(e_1) \geq c^{+t}x(e_2)$, then $x(e_0) = x(e_2) = (0.58, 0, 0, 0, 0, 0, 0, 0, 0.65, 0)$ is optimal solution for (4a).

Step12:

Since $x(e_0)$ optimizes the problem with objective function $c^{+t}x$ then

$$x^* = (0.58, 0.76, 0.94, 0, 0.87, 0, 0.83, 0, 0.65, 0)$$

6 Conclusion

In this paper, we studied the linear optimization problem with fuzzy relational inequalities constraints defined by max-Lukasiewicz operator. First, we discussed the feasibility region characterization, then; by introducing a new simplification technique the usual difficulty of finding the minimal solutions that optimize the problem with objective function $c^{+t}x$ was solved. In this relation an algorithm together with some simplification operations to accelerate the problem resolution was presented. At last, we gave an example to more illustrate of the problem.

Acknowledgements

The authors would like to thank the anonymous referees for their constructive suggestions and comments.

References

- [1] K. P. Adlassnig, *Fuzzy set theory in medical diagnosis*, IEEE Trans. Systems Man Cybernet 16 (1986) 260-265.
- [2] A. Berrached M. Beheshti A. de Korvin R. Al, *Applying Fuzzy Relation Equations to Threat Analysis*, Proceedings of the 35th Hawaii International Conference on System Sciences - 2002.
- [3] M. M. Brouke and D.G. Fisher, *Solution Algorithms for Fuzzy Relation Equations with Max-product composition*, Fuzzy sets and Systems 94 (1998) 61-69.
- [4] E. Czogala, J. Drewniak, W. Pedrycz, *Fuzzy relation equations on a finite set*, Fuzzy Sets and systems 7 (1982) 89-101.
- [5] E. Czogala, W. Predrycz, *On Identification in Fuzzy Systems and its applications in control problem*, Fuzzy Sets and systems 6 (1999) 73-83.
- [6] F. Di Martino, S. Sessa, *Digital watermarking in coding/decoding processes with fuzzy relation equations*, Soft. Comput. 10 (2006) 238-243.
- [7] A. Di Nola, C. Russo, *Lukasiewicz transform and its application to compression and reconstruction of digital images*, Information Sciences 177 (2007) 1481-1498.
- [8] A. Di Nola, S. Sessa, W. Pedrycz, and E. Sanchez, *Fuzzy Relational Equations and Their Applications in Knowledge Engineering*, Dordrecht: Kluwer Academic Press (1989).
- [9] D. Dubois, H. Prade, *Fuzzy sets and systems: Theory and Applications* Academic Press, New York, 1980.
- [10] S. C. Fang, G. Li, *Solving fuzzy relations equations with a linear objective function*, Fuzzy Sets and systems 103 (1999) 107-113.
- [11] S. C. Fang, S. Puthenpura, *Linear Optimization and Extensions: Theory and Algorithm*, Prentice-Hall, placeCity Englewood Cliffs, NJ, 1993.

- [12] M. J. Fernandez, P. Gil, *some specific types of fuzzy relation equations*, Information Sciences 164 (2004) 189-195.
- [13] S. Z. Guo, P. Z. Wang, A. Di Nola, S. Sessa, *Further contributions to the study of finite fuzzy relation equations*, Fuzzy Sets and systems 26 (1988) 93-104.
- [14] F. F. Guo, Z. Q. Xia, *An Algorithm for solving optimization Problems with one linear objective Function and Finitely Many Constraints of Fuzzy Relation Inequalities*, Fuzzy optimization and Decision making 5 (2006) 33- 47.
- [15] M. M. Gupta and J. Qi, *Design of Fuzzy Logic Controllers Based on Generalized t-operators*, Fuzzy Sets and Systems 40 (1991) 473-486.
- [16] S. M. Guu, and Y. K. Wu, *Minimizing a Linear Objective Function with Fuzzy Relation Equation Constraints*, Fuzzy Optimization and Decision Making 12 (2002) 1568-4539.
- [17] S. Z. Han, A. H. Song, and T. Sekiguchi, *Fuzzy Inequality Relation System Identification via Sign Matrix Method*, Proceeding of 1995 IEEE International Conference 3, (1995) 1375-1382.
- [18] M. Higashi, G. J. Klir, *Resolution of finite fuzzy relation equations*, Fuzzy Sets and systems 13 (1984) 65-82.
- [19] C. F. Hu, *Generalized Variational Inequalities with Fuzzy Relation*, Journal of Computational and Applied Mathematics 146 (1998) 198-203.
- [20] E. P. Klement, R. Mesiar, E. Pap, *Triangular norms. Position paper I: basic analytical and algebraic properties*, Fuzzy Sets and Systems 143 (2004) 5-26.
- [21] J. Loetamonphong, and S.-C. Fang, *Optimization of Fuzzy Relation Equations with Max-product Composition*, Fuzzy Sets and Systems 118 (2001) 509-517
- [22] J. Loetamonphong, S. C. Fang, R. E. Young, *Multi-objective optimization problems with Fuzzy Relation Equation constraints*, Fuzzy Sets and Systems 127 (2002) 141-164
- [23] V. Loia, S. Sessa, *Fuzzy relation equations for coding / decoding processes of images and videos*, Information Sciences 171 (2005) 145-172.
- [24] T. Mitsuishi, T. Terashima, Y. Shidama, *Optimization of SIRMs Fuzzy Model Using Lukasiewicz Logic*, Neural Information Processing 7664 (2012) 108-116.
- [25] A. A. Molai, *Two new algorithms for solving optimization problems with one linear objective function and finitely many constraints of fuzzy relation inequalities*, Journal of Computational and Applied Mathematics 233 (2010) 2090-2103.
- [26] A. A. Molai, *The quadratic programming problem with fuzzy relation inequality constraints*, Computers & Industrial Engineering 62 (2012) 256-263.
- [27] H. Nobuhara, B. Bede, K. Hirota, *On various eigen fuzzy sets and their application to image reconstruction*, Information Sciences 176 (2006) 2988-3010.
- [28] H. Nobuhara, K. Hirota, F. Dimartino, W. Pedrycz, S. Sessa, *Fuzzy Relation Equations for Compression/ Decompression Processes of Colour Images in the RGB and YUV Colour Spaces*, Fuzzy Optimization and Decision Making 4 (2005) 235-246.
- [29] A. Di Nola, C. Russo, *Lukasiewicz transform and its application to compression and reconstruction of digital images*, Information Sciences 177 (2007) 1481-1498.
- [30] W. Pedrycz, *On Generalized fuzzy relational equations and their applications*, Journal of Mathematical Analysis and Applications 107 (1985) 520-536.
- [31] W. Pedrycz, *An approach to the analysis of fuzzy systems*, Int. J. Control 34 (1981) 403-421.
- [32] I. Perfilieva and Vilém Novák *System of fuzzy relation equations as a continuous model of IF-THEN rules*, Information Sciences 177 (2007) 3218-3227
- [33] M. Prevot, *Algorithm for the solution of fuzzy relations*, Fuzzy Sets and systems 5 (1985) 319-322.

- [34] E. Sanchez, *Solution in composite fuzzy relation equations: Application to medical diagnosis in Brouwerian logic*, In *Fuzzy Automata and Decision Processes*, (Edited by M. M. Gupta , G. N. Saridis and B R Games), pp.221-234, North-Holland, New York , (1977).
- [35] B. S. Shieh, *Solutions of fuzzy relation equations based on continuous t-norms*, *Information Sciences* 177 (2007) 4208-4215.
- [36] E. Shivanian, *An Algorithm for Finding Solutions of Fuzzy Relation Equations with max-Lukasiewicz Composition*, *Mathware & Soft Computing* 17 (2010) 15-26.
- [37] E. Shivanian, E. Khorram, *Optimization of linear objective function subject to Fuzzy relation inequalities constraints with max-average composition*, *Iranian journal of fuzzy system* 4 (2007) 15-29.
- [38] E. Shivanian, E. Khorram, *Optimization of linear objective function subject to Fuzzy relation inequalities constraints with max-product composition*, *Iranian journal of fuzzy system* 7 (2010) 51-71.
- [39] E. Shivanian, M. Keshtkar, E. Khorram, *Geometric Programming Subject to System of Fuzzy Relation Inequalities*, *Applications and Applied Mathematics* 7 (2012) 261-282.
- [40] E. Shivanian, E. Khorram, *Monomial geometric programming with fuzzy relation inequality constraints with max-product composition*, *Computers & Industrial Engineering* 56 (2009) 1386-1392.
- [41] W. B. Vasantha, Kandasamy, F. Smarandache, *Fuzzy relational maps and neutrosophic relational maps*, hexis church rock 2004. (Chapter two)
- [42] P. Z. Wang, *How many lower solutions of finite fuzzy relation equations*, *Fuzzy mathematics (Chinese)* 4 (1984) 67-73.
- [43] P. Z. Wang, *Lattecized Linear Programming and Fuzzy Relation Inequalities*, *Journal of Mathematical Analysis and Applications* 159 (1991) 72-87.
- [44] F. Wenstop, *Deductive verbal models of organizations*, *Int. J. Man-Machine Studies* 8 (1976) 293-311.
- [45] L. A. Zadeh, *Fuzzy sets*, *Inform. Control* 8 (1965) 338-353.
- [46] Lotfi A. Zadeh, *Toward a generalized theory of uncertainty (GTU)—an outline*, *Information Sciences* 172 (2005) 1-40
- [47] H. T. Zhang, H. M. Dong, and R. H. Ren, *Programming Problem with Fuzzy Relation Inequality Constraints*, *Journal of Liaoning Noramal University* 3 (2003) 231-233.



Elyas Shivanian was born in Zanjan province, Iran in August 26, 1982. He started his master course in applied mathematics in 2005 at Amirkabir University of Technology and has finished MSc. thesis in the field of fuzzy linear programming in 2007. After his Ph.D. in the field of prediction of multiplicity of solutions of boundary value problems, at Imam Khomeini International University in 2012, he became an Assistant Professor at the same university. His research interests are analytical and numerical solutions of ODEs, PDEs and IEs. He has several papers on these subjects. He also has published some papers in other fields, for more information see <http://scholar.google.com/citations?user=MFncks8AAAAJ&hl=en>