

Available online at http://ijim.srbiau.ac.ir/

Int. J. Industrial Mathematics (ISSN 2008-5621) Vol. 8, No. 4, 2016 Article ID IJIM-00782, 7 pages Research Article



Extensions of Regular Rings

SH. A. Safari Sabet *, M. Farmani^{†‡}

Received Date: 2015-11-06 Revised Date: 2016-09-06 Accepted Date: 2016-11-02

Abstract

Let R be an associative ring with identity. An element $x \in R$ is called $\mathbb{Z}G$ -regular (resp. strongly $\mathbb{Z}G$ -regular) if there exist $g \in G$, $n \in \mathbb{Z}$ and $r \in R$ such that $x^{ng} = x^{ng}rx^{ng}$ (resp. $x^{ng} = x^{(n+1)g}$). A ring R is called $\mathbb{Z}G$ -regular (resp. strongly $\mathbb{Z}G$ -regular) if every element of R is $\mathbb{Z}G$ -regular (resp. strongly $\mathbb{Z}G$ -regular). In this paper, we characterize $\mathbb{Z}G$ -regular (resp. strongly $\mathbb{Z}G$ -regular) rings. Furthermore, this paper includes a brief discussion of $\mathbb{Z}G$ -regularity in group rings.

Keywords : Group ring; π -Regular; $\mathbb{Z}G$ -Regular; Strongly $\mathbb{Z}G$ -regular.

1 Introduction

R Ecall that an element x in R is said to be reg-ular if xyx = x, for some $y \in R$, the ring R is regular if every element of R is regular and an element $x \in R$ is said to be strongly (Von Neumann) regular if there exists $y \in R$ such that $x = x^2 y$, the ring R is strongly regular if each of elements R is strongly regular. More properties of regular and strongly regular rings can be found for example in [2, 7, 10]. An element $a \in R$ is said to be π -regular if there exist $b \in R$ and a positive integer n such that $a^n = a^n b a^n$. An element $a \in R$ is said to be strongly π -regular if $a^n = a^{n+1}b$. The ring R is π -regular if every element of R is π -regular and is strongly π -regular if every element of R strongly π -regular. By a result of Azumaya [3] and Dischinger [9], the element acan be chosen to commute with b. In particular this definition is left-right symmetric. π -regular

and strongly π -regular rings, are studied in particular in [3, 2, 4, 5, 6, 8]. Denote by $\mathbb{Z}G$ the integral group ring of a finite group G. An element $x \in R$ is said to be *G*-regular if there exist $y \in R$ and $g \in G$ such that $x^g = x^g y x^g$. The ring R is G-regular if each elements of R is G-regular. An element $x \in R$ is said to be strongly *G*-regular if there exist an element $y \in R$ and $q \in G$ such that $x^{g} = x^{2g}y$, with this property that $(x^{2})^{g} = (x^{g})^{2}$. A ring R is strongly G-regular if every element of is strongly G-regular. A ring R is abelian if every idempotent element of R is central. A ring R is called locally finite if every finite subset in it generates a finite semigroup multiplicatively. A group is locally finite if every finitely generated subgroup in it, is finite. The $n \times n$ full triangular matrix ring, the $n \times n$ upper triangular matrix ring, the $n \times n$ lower triangular matrix ring over denote by $M_n(R)$, $U_n(R)$, $L_n(R)$ respectively. In Section 2 we define $\mathbb{Z}G$ -regular and strongly $\mathbb{Z}G$ regular rings and investigate some characterization of them. Let G be a group and X a set. Then a group action (or just action) of G on Xis a binary operation:

$$\mu: X \times G \longrightarrow X$$

^{*}Department of Mathematics, Central Tehran Branch, Islamic Azad University, Tehran, Iran.

[†]Corresponding author. mino.farmani@gmail.com

[‡]Young Researchers and Elite Club, Roudehen Branch, Islamic Azad University, Roudehen, Iran.

(If there is no fear of confusion, we write $\mu(x,g)$ simply as by x^g) such that

- (i) $(x^g)^h = x^{gh}$ for all $x \in X$ and $g, h \in G$,
- (ii) $x^1 = x$ for all $x \in X$.

If S is a subset of R and $\prod_{i \in I} R_i$ is a finite direct product of $\{R_i\}_{i \in I}$, then we define: $a^{ng} = (a^g)^n, (x_i)_{i \in I}^g = (x_i^g)_{i \in I}, S^g = \{x^g | x \in S\}$. For each $(x_i)_{i \in I} \in \prod_{i \in I} R_i, g \in G, n \in \mathbb{Z}$. The main purpose of Section 3 is to characterize $\mathbb{Z}G$ regular and strongly $\mathbb{Z}G$ -regular group rings.

2 Preliminarie

Definition 2.1 An element $x \in R$ is called $\mathbb{Z}G$ regular (resp. Strongly $\mathbb{Z}G$ -regular) if there exist $g \in G, n \in \mathbb{Z}$ and $r \in R$ such that $x^{ng} = x^{ng}rx^{ng}$ (resp. $x^{ng} = x^{(n+1)g}r$). A ring R is called $\mathbb{Z}G$ regular (resp. Strongly $\mathbb{Z}G$ -regular) if every element of R is $\mathbb{Z}G$ -regular (resp. strongly $\mathbb{Z}G$ regular). So an element $x \in R$ is $\mathbb{Z}G$ -regular (resp. Strongly $\mathbb{Z}G$ -regular) if there exists $g \in$ G such that x^g is π -regular (resp. Strongly π regular).

Now we define a $\mathbb{Z}G$ -regular ideal as follows: Let $\mu : R \times G \longrightarrow R$ be a group action and I be a two-sided ideal of R. Then group G acts on R/Iby the rule $\mu(r + I, g) = \mu(r, g) + I$.

Definition 2.2 Let J be a two-sided ideal of a ring R. J is a $\mathbb{Z}G$ -regular ideal of R in case for any $x \in J$, there exist $n \in \mathbb{Z}$ and $y \in R$ such that $x^{ng} = x^{ng}yx^{ng}$.

Theorem 2.1 Any factor ring of a $\mathbb{Z}G$ -regular (resp. Strongly $\mathbb{Z}G$ -regular) ring is $\mathbb{Z}G$ -regular (resp. Strongly $\mathbb{Z}G$ -regular). In particular a homomorphic image of a $\mathbb{Z}G$ -regular (resp. Strongly $\mathbb{Z}G$ -regular) ring is $\mathbb{Z}G$ -regular (resp. Strongly $\mathbb{Z}G$ -regular).

Proof. Let R be $\mathbb{Z}G$ -regular (resp. Strongly $\mathbb{Z}G$ -regular) and I be a two-sided ideal of R. Let $\overline{x} = x + I \in R/I$. Since R is $\mathbb{Z}G$ -regular, then there exist $g \in G$, $n \in \mathbb{Z}$ and $r \in R$ such that $x^{ng} = x^{ng}rx^{ng}$ (resp. $x^{ng} = x^{(n+1)g}r$). This implies $\overline{x^{ng}} = \overline{x^{ng}}\overline{r}\overline{x^{ng}}$ (resp. $\overline{x^{ng}} = \overline{x^{ng}}\overline{r}$). Thus by definition we have $\overline{x}^{ng} = \overline{x}^{ng}\overline{r}\overline{x}^{ng}$ (resp. $\overline{x}^{ng} = \overline{x}^{ng}\overline{r}\overline{x}^{ng}$ (resp. $\overline{x}^{ng} = \overline{x}^{ng}\overline{r}\overline{x}^{ng}$).

Theorem 2.2 Let R be a ring. Then the following conditions are equivalent:

- (i) R is strongly $\mathbb{Z}G$ -regular.
- (ii) R/N is strongly ZG-regular that is the prime radical N of R.
- (iii) Every prime factor ring of R is strongly ZGregular.

Proof. It suffices to show that (iii) implies (i). Suppose R contains an element a that is not strongly $\mathbb{Z}G$ -regular. Then by Zorn s lemma, there exists an ideal I of R that is maximal with respect to the property that \overline{a} is not strongly $\mathbb{Z}G$ regular in $\overline{R} = R/I$. Since I can not be prime, there exist ideals K, L properly containing I such that $KL \subseteq I$ Then we can find a $n \in \mathbb{Z}$ such that $a^{ng_1} - a^{(n+1)g_1}x \in K$ and $a^{ng_2} - a^{(n+1)g_2}y \in L$ for some $x, y \in R, g_1, g_2 \in G$. But

 $a^{n(g_1+g_2)} - a^{(n+1)(g_1+g_2)}(a^{-g_1}y + a^{-g_2}x + xy) = (a^{ng_1} - a^{(n+1)g_1}x)(a^{ng_2} - a^{(n+1)g_2}y) \in KL \subseteq I$

Which is a contradiction.

Lemma 2.1 Let G be a group acts on the ring R by this property that $(xy)^g = x^g y^g$ for each $x, y \in$ R. If $x, y \in R$, $g \in G$ and $x' = x^{ng} - x^{ng} y x^{ng}$, and if $x'^{n'h} = x'^{n'h} a^{n'} x'^{n'h}$ for some $a \in R$ and some $h \in G$. Then $x^{ng} = x^{ng} b x^{ng}$ for some $b \in R$.

Proof. We have

$$\begin{aligned} x^{ng} &= x' + x^{ng}yx^{ng} \\ &= (x'^{n'h}a^{n'}x'^{n'h})^{n'^{-1}h^{-1}} + x^{ng}yx^{ng} \\ &= x'a^{h^{-1}}x' + x^{ng}yx^{ng} \\ &= (x^{ng} - x^{ng}yx^{ng})a^{h^{-1}}(x^{ng} - x^{ng}yx^{ng}) \\ &+ x^{ng}yx^{ng} \\ &= (x^{ng}a^{h^{-1}} - x^{ng}yx^{ng}a^{h^{-1}})(x^{ng} - x^{ng}yx^{ng}) \\ &+ x^{ng}yx^{ng} \\ &= x^{ng}a^{h^{-1}}x^{ng} - x^{ng}yx^{ng}a^{h^{-1}}x^{ng} \\ &- x^{ng}a^{h^{-1}}x^{ng}yx^{ng} + x^{ng}yx^{ng}a^{h^{-1}}x^{ng}yx^{ng} \\ &+ x^{ng}yx^{ng} \\ &= x^{ng}(a^{h^{-1}} - yx^{ng}a^{h^{-1}} - a^{h^{-1}}x^{ng}y \\ &+ yx^{ng}a^{h^{-1}}x^{ng}y + y)x^{ng} \end{aligned}$$

Now by taking $b = a^{h^{-1}} - yx^{ng}a^{h^{-1}} - a^{h^{-1}}x^{ng}y + yx^{ng}a^{h^{-1}}x^{ng}y + y$ we have $x^{ng} = x^{ng}bx^{ng}$.

Theorem 2.3 Let $J \subseteq K$ be two sided ideals in a ring R. So J and K/J are both $\mathbb{Z}G$ -regular if and only if K is $\mathbb{Z}G$ -regular. **Proof.** Assume that J and K/J are both $\mathbb{Z}G$ -regular. Given $x \in J$, it follows from the regularity of K/J that $x^{ng} - x^{ng}yx^{ng} \in J$ for some $y \in K$ and $n \in \mathbb{Z}$. Consequently, $(x^{ng} - x^{ng}yx^{ng})^{n'h} = (x^{ng} - x^{ng}yx^{ng})^{n'h}z(x^{ng} - x^{ng}yx^{ng})^{n'h}$ for some $z \in J$, from which by lemma 2.1 we conclude that $x^{ng} = x^{ng}wx^{ng}$ for some $w \in K$. Thus, Kis $\mathbb{Z}G$ -regular. Conversely, assume that K is a $\mathbb{Z}G$ -regular ring. Clearly K/J is $\mathbb{Z}G$ -regular. It suffices to show that J is $\mathbb{Z}G$ -regular. Since Kis a $\mathbb{Z}G$ -regular ring then for any $\forall x \in J$, there exist $g \in G$ and $y \in K$ and $n \in \mathbb{Z}$ such that

$$x^{ng} \in J, x^{ng} = x^{ng}yx^{ng}$$

Now by taking $z = yx^{ng}y \in J$ we have:

$$x^{ng} = x^{ng} z x^{ng}$$

Therefore, J is a $\mathbb{Z}G$ -regular ideal.

Lemma 2.2 A finite direct product $\prod_{i \in I} R_i$ (*I* is a finite set) of $\mathbb{Z}G$ -regular rings $\{R_i\}_{i \in I}$ is $\mathbb{Z}G$ -regular.

Proof. At first we prove that direct product of two $\mathbb{Z}G$ -regular rings is $\mathbb{Z}G$ -regular. Let R_1 and R_2 be two $\mathbb{Z}G$ -regular rings. Then for every $(a_1, a_2) \in R_1 \times R_2$ there exist $g_1, g_2 \in G$, $(r_1, r_2) \in R_1 \times R_2$ and $n_1, n_2 \in \mathbb{Z}$ such that $a_1^{n_1g_1} = a_1^{n_1g_1}r_1a_1^{n_1g_1}$ and $a_2^{n_2g_2} = a_2^{n_2g_2}r_2a_2^{n_2g_2}$. Now by setting $ng = n_1n_2g_1g_2$ we have

$$(a_{1}, a_{2})^{ng} = (a_{1}^{ng}, a_{2}^{ng})$$

= $((a_{1}^{n_{1}g_{1}})^{n_{2}g_{2}}, (a_{2}^{n_{2}g_{2}})^{n_{1}g_{1}})$
= $((a_{1}^{n_{1}g_{1}}r_{1}a_{1}^{n_{1}g_{1}})^{n_{2}g_{2}}, (a_{2}^{n_{2}g_{2}}r_{2}a_{2}^{n_{2}g_{2}})^{n_{1}g_{1}})$
= $(a_{1}, a_{2})^{ng}(r_{1}^{n_{2}g_{2}}, r_{2}^{n_{1}g_{1}})(a_{1}, a_{2})^{ng}$
(2.2)

Thus by induction any finite direct product of $\mathbb{Z}G$ -regular rings is $\mathbb{Z}G$ -regular.

- **Theorem 2.4 (i)** Let $x \in R$ be $\mathbb{Z}G$ -regular, then there exist $g \in G$, $n \in \mathbb{Z}$ and $r \in R$ such that $x^{ng}r$ is idempotent.
- (ii) If an element $x \in R$ is π -regular, then it is $\mathbb{Z}G$ -regular by taking G to be trivial group.
- (iii) An element $x \in R$ is $\mathbb{Z}G$ -regular if there exist $g \in G$, $n \in \mathbb{Z}$ such that x^{ng} is Von Neumann.

Proof. (i) Since $x \in R$ is $\mathbb{Z}G$ -regular thus there exist $g \in G$, $n \in \mathbb{Z}$ and $r \in R$ such that $x^{ng} = x^{ng}rx^{ng}$ therefore $x^{ng}r = x^{ng}rx^{ng}r = (x^{ng}r)^2$. (ii), (iii) are trivial.

Theorem 2.5 Let S be the center of $\mathbb{Z}G$ -regular ring R with the property that $S^{ng} \subseteq S$, for any $g \in G$, $n \in \mathbb{Z}$. Then S is $\mathbb{Z}G$ -regular.

Proof. Let R be a ring with center S, and let $x \in S$. There exist $y \in R$, $n \in \mathbb{Z}$ and $g \in G$ such that $x^{ng}yx^{ng} = x^{ng}$, and we set $z = yx^{ng}y$. Note that

$$x^{ng}zx^{ng} = x^{ng}yx^{ng}yx^{ng} = x^{ng}yx^{ng}z^{ng} = x^{ng}z^{ng}z^{ng}z^{ng} = x^{ng}z^{ng}z^{ng}z^{ng}z^{ng}z^{ng} = x^{ng}z^{$$

For any $r \in R$, we have

$$zr = yx^{ng}yr$$

$$= y^{2}rx^{ng}$$

$$= y^{2}rx^{ng}yx^{ng}$$

$$= y^{2}rx^{ng}yr^{ng}y$$

$$= yx^{ng}yrx^{ng}y$$

$$= yx^{ng}yr^{ng}ry$$

$$= yx^{ng}ry$$

$$(2.3)$$

Similarly we have $rz = yrx^{ng}y$, so $rz = yrx^{ng}y = yx^{ng}ry = zr$, therefore $z \in S$. Thus S is also $\mathbb{Z}G$ -regular.

Proposition 2.1 A ring R is strongly $\mathbb{Z}G$ regular, if and only if R satisfies the descending chain condition on principal right ideals of the form $a^{g}R \supseteq a^{2g}R \supseteq ...$, for every $a \in R$ and an element $g \in G$.

Proof. One direction is clear. Assume R is not strongly $\mathbb{Z}G$ -regular. Then there exists an element $a \in R$ such that $x^{ng} \neq x^{(n+1)g}r$ for any $r \in R$ and $g \in G$ and $n \in \mathbb{Z}$. We have a descending chain $a^g R \supset a^{2g} R \supset \ldots$ of ideals of R which does not terminate, which is a contradiction.

Lemma 2.3 Let R be a ring. If R is locally finite and $a \in R$, then a^t is an idempotent for some positive t.

Proof. see [11].

Theorem 2.6 Let R be a ring. If R is a locally finite ring, then R is strongly $\mathbb{Z}G$ -regular.

By lemma 2.3, a locally finite ring R satisfies the descending chain condition on principal right ideals of form $aR \supseteq a^2R \supseteq ...$, for every a in R; then R satisfies the descending chain condition on principal right ideals of form $a^gR \supseteq a^{2g}R \supseteq ...$, for every $a \in R$ and $g \in G$. Therefore R is strongly $\mathbb{Z}G$ -regular by proposition 2.1.

Proposition 2.2 Let the $n \times n$ full triangular matrix ring over R be $\mathbb{Z}G$ -regular. Then R is $\mathbb{Z}G$ -regular.

Proof. It is obvious. We introduced π -regular (resp. Strongly π -regular) rings as an example of $\mathbb{Z}G$ -regular (resp. Strongly $\mathbb{Z}G$ -regular) rings by taking G to be trivial group. Lee and Kim showed in [12], that the n by n full matrix rings over strongly π -regular ring R, need not be strongly π -regular (see: Example 2.1), so we conclude that the $n \times n$ full triangular matrix ring over R need not be strongly $\mathbb{Z}G$ -regular rings.

Theorem 2.7 For a ring R and a positive integer m, the following conditions are equivalent.

- (a) R is locally finite.
- (b) $M_n(R)$ is locally finite.
- (c) $U_n(R)$ is locally finite.
- (d) $L_n(R)$ is locally finite.

Proof. see [11].

Example 2.1 Let R be a locally finite ring. $M_n(R), U_n(R), L_n(R)$ R are examples of strongly $\mathbb{Z}G$ -regular rings by Theorems 2.6 and 2.7.

Lemma 2.4 Let R be an abelian $\mathbb{Z}G$ -regular ring. Then for each $x \in R$, there exist $r \in R, g \in$ G and $n \in \mathbb{Z}$ such that $x^{ng}r = rx^{ng}$.

Since R is $\mathbb{Z}G$ -regular, then by theorem 2.4 (i), for each $x \in R$, there exist $g \in G$, $n \in \mathbb{Z}$, $r \in R$ such that $x^{ng}r, rx^{ng} \in Id(R)$ and since R is abelian then $x^{ng}r, rx^{ng} \in Z(R)$, therefore we have:

$$\begin{aligned}
x^{ng}r &= (x^{ng}rx^{ng})r \\
&= x^{ng}(rx^{ng})r \\
&= x^{ng}r(rx^{ng}) = r(x^{ng}rx^{ng}) \\
&= rx^{ng}
\end{aligned}$$
(2.4)

Definition 2.3 An element $x \in R$ is said unit $\mathbb{Z}G$ -regular if there exist $g \in G$ and $u \in U(R)$ and $n \in \mathbb{Z}$ depending on x such that $x^{ng} = x^{ng}ux^{ng}$. R is unit $\mathbb{Z}G$ -regular if every element of R is unit $\mathbb{Z}G$ -regular.

Theorem 2.8 Let R be an abelian $\mathbb{Z}G$ -regular ring. Then R is unit $\mathbb{Z}G$ -regular.

Since R is abelian $\mathbb{Z}G$ -regular by lemma 2.4, for each $x \in R$, there exist $g \in G$, $y \in R$ and $n \in \mathbb{Z}$, such that $x^{ng}y = yx^{ng}$. Let $u = x^{ng} + x^{ng}y - 1$ and $v = x^{ng}y + x^{ng}y^2 - 1$. Since $x^{ng}y = yx^{ng}$, then we have:

$$uv = (x^{ng} + x^{ng}y - 1)(x^{ng}y + x^{ng}y^2 - 1)$$

$$= x^{ng}(x^{ng}y) + x^{ng}x^{ng}y^2 - x^{ng}$$

$$+ (x^{ng}y)(x^{ng}y) + (x^{ng}y)(x^{ng}y^2)$$

$$-x^{ng}y - x^{ng}y - x^{ng}y^2 + 1$$

$$= x^{ng}(yx^{ng}) + (x^{ng}yx^{ng})y - x^{ng}$$

$$+ (x^{ng}yx^{ng})y + (x^{ng}yx^{ng})y^2$$

$$-x^{ng}y - x^{ng}y - x^{ng}y^2 + 1$$

$$= x^{ng} + x^{ng}y - x^{ng}y + x^{ng}y$$

$$+ x^{ng}y^2 - x^{ng}y - x^{ng}y - x^{ng}y^2 + 1 = 1$$

(2.5)

And

$$vu = (x^{ng}y + x^{ng}y^2 - 1)(x^{ng} + x^{ng}y - 1)$$

$$= x^{ng}yx^{ng} + x^{ng}yx^{ng}y - x^{ng}y$$

$$+ x^{ng}y^2x^{ng} + x^{ng}y^2x^{ng}y$$

$$- x^{ng}y^2 - x^{ng} - x^{ng}y + 1$$

$$= x^{ng}yx^{ng} + (x^{ng}yx^{ng})y - x^{ng}y$$

$$+ x^{ng}y(yx^{ng}) + x^{ng}y(yx^{ng})y - x^{ng}y^2 - x^{ng} - x^{ng}y + 1$$

$$= x^{ng} + x^{ng}y - x^{ng}y + (x^{ng}yx^{ng})y$$

$$+ x^{ng}yx^{ng}y^2 - x^{ng}y^2 - x^{ng} - x^{ng}y + 1$$

$$= x^{ng} + x^{ng}y - x^{ng}y + x^{ng}y$$

$$+ x^{ng}y^2 - x^{ng}y^2 - x^{ng}y + x^{ng}y + 1 = 1$$

$$= x^{ng} + x^{ng}y^2 - x^{ng}y^2 - x^{ng}y + 1 = 1$$

$$(2.6)$$

Therefore, uv = vu = 1. Moreover,

$$\begin{aligned} x^{ng}vx^{ng} &= x^{ng}(x^{ng}y \\ &+ x^{ng}y^2 - 1)x^{ng} \\ &= x^{ng}x^{ng}yx^{ng} \\ &+ x^{ng}x^{ng}y^2x^{ng} - x^{ng}x^{ng} \\ &= x^{ng}x^{ng} + x^{ng}x^{ng}yyx^{ng} \\ &= x^{ng}x^{ng} \\ &= (x^{ng}yx^{ng})yx^{ng} \\ &= x^{ng}yx^{ng} \\ &= x^{ng}yx^{ng} \\ &= x^{ng}yx^{ng} \end{aligned}$$
(2.7)

Theorem 2.9 Let R be an abelian $\mathbb{Z}G$ -regular ring, and $x \in R$. Then there exist $g \in G$, $n \in \mathbb{Z}$ such that $x^{ng} = eu$, for some $e \in Id(R)$ and $u \in U(R)$.

Proof. By theorem 2.8, R is unit $\mathbb{Z}G$ -regular. Thus there exists $v \in U(R)$ such that $x^{ng} = x^{ng}vx^{ng}$. Let u be the multiplicative inverse of v in R, then $x^{ng} = x^{ng}uv = x^{ng}vu = eu$. Since $e = x^{ng}v \in Id(R)$. Thus $x^{ng} = eu$ for some $e \in Id(R)$ and $u \in U(R)$.

Theorem 2.10 Let R be an abelian ring. Then the following statements are equivalent: (i) R is a unit $\mathbb{Z}G$ -regular ring.

- (ii) For every a ∈ R, there exist g ∈ G, n ∈ Z such that a^{ng} can be written as a product of a unit, and an idempotent of R.
- (iii) For every $a \in R$, there exist $g \in G$, $n \in \mathbb{Z}$ such that a^{ng} can be written as a product of an idempotent and a unit of R.

Proof. (i \Rightarrow ii) By theorem 2.9, is clear.

(ii \Rightarrow i) Suppose there exists $g \in G$, $n \in \mathbb{Z}$ such that $a^{ng} = ve$ where $v \in U(R)$ and $e^2 = e$. The latter implies $v^{-1}a^{ng}=v^{-1}a^{ng}v^{-1}a^{ng}$, so $a^{ng} = a^{ng}v^{-1}a^{ng}$, as desired.

3 $\mathbb{Z}G$ -regular group ring

Let R be a ring and G a group. We shall denote the group ring of G over R as RG. The augmentation ideal of RG is generated by $\{1, g\}$. We shall use Δ to denote the augmentation ideal of RG. It is known that R is a homomorphic image of RG. Since $RG/\Delta \cong R$. For any element $x = \sum_{g \in G} x_g g \in RG$, the support of x, written as Supp(x), is the subset of G consisting of all those $g \in G$ such that $x_g \neq 0$. Since $x_g \neq 0$ for only finitely many $g \in G$, so Supp(x) is a finite subset of G.

Corollary 3.1 Let R be a ring and G a group. If RG is a $\mathbb{Z}G$ -regular ring, then R is a $\mathbb{Z}G$ regular ring.

Proof. Since R is homomorphic image of RG, then R is $\mathbb{Z}G$ -regular by theorem 2.1. For any idempotent e in a ring, we have the following peirce decomposition:

$$R = eRe \oplus eRf \oplus fRe \oplus fRf$$

Where f = 1 - e is the complementary idempotent to e. Two ring eRe and fRf be characterized by the equation:

 $eRe = \{rR : er = r = re\},\$ $fRf = \{rR : fr = r = rf\}$

Lemma 3.1 *e is a central idempotent iff* eRf = fRe = 0.

Proof. For $r \in R$, erf = 0 and fre = 0 amount to er = ere = re.

Proposition 3.1 Let $e \neq 0$ be any central idempotent in R. If eRe and fRf are $\mathbb{Z}G$ -regular, then R is a $\mathbb{Z}G$ -regular ring.

Since e is a central idempotent, then we have the peirce decomposition:

$$R = eRe \oplus fRf$$

Thus by lemma 2.2, since eRe and fRf are $\mathbb{Z}G$ -regular then R is $\mathbb{Z}G$ -regular.

Theorem 3.1 Let $e_1 + \ldots + e_n = 1$ be a decomposition of 1 into sums of orthogonal idempotents. If $e_i Re_i$ is $\mathbb{Z}G$ -regular for each i, then R is $\mathbb{Z}G$ -regular.

Proof. It is obvious from Lemma 2.2 and Proposition 3.1.

Theorem 3.2 Let R be a commutative semiperfect ring and G a group, and let (eRe)G be $\mathbb{Z}G$ regular for each local idempotent e in R. Then RG is $\mathbb{Z}G$ -regular.

Proof. Since R is semiperfect, so by theorem 6.27 of [1], R has a complete orthogonal set $e_1, e_2, ..., e_n$ of idempotent R. So e_i is a local idempotent for each $i \in \{0, ..., n\}$. Now by hypothesis, $(e_i R e_i)G$ is $\mathbb{Z}G$ -regular. Since, $(e_i R e_i)G \cong e_i(RG)e_i$ for each i, it follows that $e_i(RG)e_i$ is $\mathbb{Z}G$ -regular. Hence RG is $\mathbb{Z}G$ -regular by proposition 3.1.

Theorem 3.3 Let R be a ring in which 2 is invertible and $G = \{1, g\}$ be a group. Then RG is $\mathbb{Z}G$ -regular if and only if R is $\mathbb{Z}G$ -regular.

Proof. If RG is $\mathbb{Z}G$ -regular, then by corollary 3.1, R is $\mathbb{Z}G$ -regular. Conversely, since R is $\mathbb{Z}G$ -regular and 2 is invertible in R, then $RG \cong R \times R$ via the map $a + bg \iff (a + b, a - b)$. Hence RG is $\mathbb{Z}G$ -regular by lemma 2.2.

Theorem 3.4 Let R be a ring and G a group. Then RG is strongly $\mathbb{Z}G$ -regular if and only if (R/P)G is strongly $\mathbb{Z}G$ -regular for every prime ideal P of R.

Proof. If RG is strongly $\mathbb{Z}G$ -regular, and I is an ideal of R, then since

$$(R/I)G\cong RG/IG$$

and homomorphic images of strongly $\mathbb{Z}G$ -regular rings strongly $\mathbb{Z}G$ -regular, it follows that (R/I)Gis strongly $\mathbb{Z}G$ -regular.

Conversely, suppose to the contrary that RG is not strongly $\mathbb{Z}G$ -regular. Then there exists an element $x \in RG$ such that for any $n \in \mathbb{Z}$ and $g \in G$, $x^{ng} \neq x^{(n+1)g}y$ for any $y \in RG$. Therefore the sequence

 $x^g RG \supseteq x^{2g} RG$

 $\supseteq \ldots \supseteq x^{ng}RG \supseteq x^{(n+1)g}RG \supseteq \ldots$

ideals of does not terminate. Let \Im be the set of all ideals I of R such that the sequence $(x + IG)^g (RG/IG)$

 $\supseteq (x + IG)^{2g} (RG/IG) \supseteq \dots$

does not terminate. Note that $\Im \neq \emptyset$, since (0) $\in \Im$. Furthermore, \Im is partially ordered by inclusion. Let $(I_{\alpha})_{\alpha\in\Omega}$ be a chain of elements of \Im and let $J = \bigcup_{\alpha\in\Omega}I_{\alpha}$. Clearly, J is an ideal of R and $I_{\alpha} \subset J$ for all $\alpha \in \Omega$. We show that $J \in \Im$. Suppose that $J \notin \Im$. Then $z = x^{ng} - x^{(n+1)g}r \in JG$ for some $r \in RG, g \in G$ and $n \in \mathbb{Z}$. Since Supp(z) is finite, there exists some $\alpha \in \Omega$ such that $z \in I_{\alpha}G$. It follows that the sequence $(x + I_{\alpha}G)^g(RG/I_{\alpha}G)$

 $\supseteq (x + I_{\alpha}G)^{2g}(RG/I_{\alpha}G) \supseteq \dots$

terminates, which is a contradiction. Therefore $J \in \Im$ and thus by Zorn 's Lemma, \Im contains a maximal element M. Since $(R/M)G \cong RG/MG$ is not strongly $\mathbb{Z}G$ -regular, it follows by hypothesis that M is not a prime ideal. Therefore there exist ideals A, B of R such that $AB \subseteq M$ but $A, B \not\subseteq M$. Let A' = M + A and B' = B + M. Then M is strictly contained in A' and B', and we also have that:

$$A'B' = (M+A)(M+B) \subseteq M$$

By the maximality of M in \Im , the sequences $(x + A'G)^{2g}(RG/A'G)$ $\supseteq (x + A'G)^{2g}(RG/A'G) \supseteq \dots$ And $(x + B'G)^{2g}(RG/B'G)$

 $\supseteq (x + B'G)^{2g} (RG/B'G) \supseteq \dots$ both terminate. Hence there exists $m \in \mathbb{Z}$ such that $(x^{mg} + A'G)(RG/A'G) = (x^{(2m+1)g} + A'G)(RG/A'G)$ and $(x^{mg} + B'G)(RG/B'G) = (x^{(2m+1)g} + B'G)(RG/B'G).$

It follows that $x^{mg} - x^{(2m+1)g}s \in A'G$ and $x^{mg} - x^{(2m+1)g}t \in B'G$ for some $s, t \in RG$. Therefore: $(x^{mg} - x^{(2m+1)g}s)(x^{mg} - x^{(2m+1)g}t)$ $\in (A'B')G \subseteq MG$

Form which it follows that $x^{mg} - x^{(2m+1)g}w \in MG$ for some $w \in RG$. Hence the sequence: $(x + MG)^g (RG/MG)$

 $\supseteq (x + MG)^{2g} (RG/MG) \supseteq \dots$

terminates; contradicting the fact that $M \in \mathfrak{T}$. We thus have that RG must be a strongly $\mathbb{Z}G$ ring. **Theorem 3.5** Let R be a ring with artinian prime factors and G be a locally finite group. Then RG is strongly $\mathbb{Z}G$ -regular.

Proof. Let P be a prime ideal of R and $x = \sum_{g \in G} r_g g \in (R/P)G$. Let H_x be the subgroup of G generated by the support of x. Since Supp(x) is finite and G is locally finite, it follows that H_x is finite. It is clear that $x \in (R/P)H_x$ is strongly $\mathbb{Z}G$ -regular. Indeed, since R/P is artinian and H_x is finite, so $(R/P)H_x$ is artiniar; hence strongly $\mathbb{Z}G$ -regular. Since x is arbitrary in (R/P)G, so (R/P)G is also strongly $\mathbb{Z}G$ -regular. By theorem 3.3, it follows that RG is strongly $\mathbb{Z}G$ -regular.

Theorem 3.6 Let R be a ring, G be a group and $U_n(RG)$ be strongly $\mathbb{Z}G$ -regular for $n \ge 2$. Then R is strongly $\mathbb{Z}G$ -regular.

Proof. As $U_n(RG)$ is strongly $\mathbb{Z}G$ -regular, so by example 2.1, RG is strongly $\mathbb{Z}G$ -regular. Hence by Corallary 3.1, R is strongly $\mathbb{Z}G$ -regular.

4 Examples

Here we give some examples of $\mathbb{Z}G$ -regular rings.

Example 4.1 It is clear that if G is a trivial group (group with only one element) then R is $\mathbb{Z}G$ -regular for $n \geq 1$ iff R is π -regular.

Example 4.2 One easily checks that $\mathbb{Z}/4\mathbb{Z}$ and $\mathbb{Z}/6\mathbb{Z}$ are $\mathbb{Z}G$ -regular rings because they are π -regular rings.

Example 4.3 Let G = U(R) (where U(R) is the group of units in R) and X a set, then a regular action μ of G on X is a function:

$$\mu: X \times G \longrightarrow X: (x,g) = gx \tag{4.8}$$

And conjugate action is a function:

$$\mu: X \times G \longrightarrow X: (x,g) = gxg^{-1} \qquad (4.9)$$

Example 4.4 Let G = U(R). An element $x \in R$ is said unitary π -regular (resp. strongly unitary R-regular) if there exist $g \in G$ and $r \in R$ and $n \in \mathbb{Z}$ such that $(gx)^n = (gx)^n r(gx)^n$ (resp. $(gx)^n = (gx)^{(n+1)}r$). R is unitary π -regular (resp. strongly unitary π -regular) if every element of R is unitary π -regular (resp. strongly unitary π -regular). **Example 4.5** Let G = U(R). An element $x \in R$ is said conjugate π -regular (resp. strongly conjugate π -regular) if there exist $g \in G$ and $r \in R$ and $n \in \mathbb{Z}$ such that $(x)^n = (x)^n g^{-1} rg(x)^n$ (resp. $(x)^n = (x)^{n+1} gxg)$. R is conjugate π -regular (resp. strongly conjugate π -regular) if every element of R is conjugate π -regular (resp. strongly conjugate π -regular).

Example 4.6 Let Aut(R) be automorphism group of R. An element $x \in R$ is called Automorphic π -regular ((Aut) π -regular) if there exist $\alpha \in Aut(R), r \in R$ and $n \in \mathbb{Z}$ such that $(x^{\alpha})^n =$ $(x^{\alpha})^n r(x^{\alpha})^n$. R is Automorphic π -regular every element of R is automorphic π -regular.

5 Conclusion

Ring theory is a subject of central importance in algebra. Historically, some of major discoveries in ring theory have helped shape the course of development of abstract algebra. In the moment, ring theory is a fertile meeting ground for group rings. In this paper, we characterized $\mathbb{Z}G$ -regular and strongly $\mathbb{Z}G$ -regular group rings.

References

- F. W. Anderson, K. R. Fuller, *Rings and categories of modules*, Second edition, Graduate Texts in Mathematics, Springer-Verlang, New York, 13 (1992).
- [2] P. Ara, π-regular rings have stable range one, Proc. Amer. Math. Soc. 11 (1996) 3293-3298.
- [3] G. Azumaya, Strongly π-regular rings, J.
 Fac. Sci. Hokkaido Univ. 13 (1954) 34-39.
- [4] A. Badawi, Semicommutative π-regular rings, Commun. Algebra 1 (1993) 151-157.
- [5] A. Badawi, Abelian π-regular rings, Common. Algebra 4 (1997) 1009-1021.
- [6] W. D. Burgess, P. Menal, Strongly π-regular rings and homomorphisms into them, Commun. Algebra 16 (1988) 1701-1725.
- [7] V. Camillo, Y. Xiao, *Regular rings*, Commun. Algebra, 10 (1994), 4095-4112.

- [8] A. Y. M. Chin, H. V. Chen, On strongly πregular, Southeast Asian Bulletin of mathematics 26 (2002) 387-390.
- [9] M. F. Dischinger, Sur les anneaux fortement π-rguliers, C. R. Acad. Sci. Paris, Ser. A 283 (1976) 571-573.
- [10] K. R. Goodearl, Von Neumann regular rings, Pitman, London (1979).
- [11] Ch. Huh, N. K. Kim, Y. Lee, Example of strongly π-regular rings, Pure Appl. Algera 189 (2004) 195-210.
- [12] N. K. Kim, Y. Lee, On strong π -regularity and π -regular, Commun. Algebra 39 (2011) 4470-4482.



Shaaban Ali Safari Sabet is an Associate Professor at the Department of Mathematice in Islamic Azad University, Central Branch, Tehran, Iran. He received his Master in Amirkabir University of Technology, Tehran, Iran and re-

ceived PHD in Islamic Azad University, Science and Research Branch, Tehran, Iran. Now, He is Assistant of Planning and Knowledge-based economic affairs of Central Tehran Branch. His research interests includes Noncommutative Rings Theory and Graphs Theory.



Marzieh Farmani is an Assistant Professor at the Department of Mathematice in Islamic Azad University, Roudehen Branch, Tehran, Iran and member of Young Researchers Club. She received his Master and PHDin Islamic Azad

University, Science and Research Branch, Tehran, Iran and Central Tehran, Iran respectively. Her research interests includes Noncommutative Rings Theory and Groups Theory.

337

Extensions of Regular Rings

SH. A. Safari Sabet, M. Farmani

توسيعي از حلقه هاى منظم

چکیدہ:

فرض کنید R یک حلقه شرکت پذیر با عنصر همانی باشد. یک عنصر $x \in R$ ، $x \in \mathbb{Z}$ -منظم نامیده می شود اگر یک عنصر $r \in R$ یک حلقه $g \in G \in \mathbb{Z}$ موجود باشد به طوری که $n^{g} = x^{ng}rx^{ng}$. یک حلقه $g_{\mathbb{Z}}$ -منظم نامیده می شود اگر هر عنصر آن $g_{\mathbb{Z}}$ -منظم باشد. در این مقاله ما حلقه های $g_{\mathbb{Z}}$ -منظم (قویاً $g_{\mathbb{Z}}$ -منظم) را بررسی خواهیم نمود. علاوه بر آن، این مقاله شامل یک بحث مختصری راجع به $g_{\mathbb{Z}}$ -منظم بودن گروه حلقه ها می باشد.