



Some Improvements of The Cordero-Torregrosa Method for The Solution of Nonlinear Equations

M. Mohamadi Zadeh ^{*}, T. Lotfi ^{†‡}, M. Amirfakhriyan ^{§¶}

Received Date: 2020-01-24 Revised Date: 2020-05-17 Accepted Date: 2021-02-15

Abstract

In this paper, two adaptive methods with memory are improved based on the Cordero-Torregrosa method. The technique of adaptive method increases the efficiency index as high as possible. The new proposed derivative free methods have possessed the convergence order 7.46315 and 7.99315, and they use the information from the last two iterations. Finally, we provide convergence analysis and numerical examples to illustrate the efficiency and applicability of the proposed methods.

Keywords : Nonlinear equation; Iterative methods; The method without and with memory; Efficiency index; Adaptive method; Convergence order.

1 Introduction

Finding the solution of nonlinear equation $f(x) = 0$, where $f : D \subset R \rightarrow R$ which is a scalar function at an open interval, is important and challenging a problem in the field of computational mathematics. To approximate the simple root of $f(x) = 0$, it is appropriate to use iterative methods where it starts with initial guess x_0 and conjectures by generating new iterations, for example 1, $x_{n+1} = \varphi(x_n)$. The first well-known iterative method was

introduced by the Newton about 300 years ago and it is defined as

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)},$$

and continued with new published methods [1, 9, 13, 14, 15, 16, 17, 18]. Kung and Traub in [5] proposed the following method with memory based on the Newton method:

$$\begin{cases} w_n = x_n + \gamma f(x_n), \\ \gamma = -\frac{1}{f[x_n, x_{n-1}]}, \\ x_{n+1} = x_n - \frac{f(x_n)}{f[x_n, w_n]}, \end{cases} \quad n = 0, 1, 2, \dots \quad (1.1)$$

with the order of convergence 2.414, where $f[x_n, w_n] = \frac{f(x_n) - f(w_n)}{x_n - w_n}$ signifies a divided difference. Kung and Traub proposed that each n-step method without memory is optimal if it uses $n + 1$ functional evaluations with conver-

^{*}Department of Mathematics, Hamedan Branch, Islamic Azad University, Hamedan, Iran.

[†]Corresponding author. lotfitaher@yahoo.com, Tel:+98(81)344163085.

[‡]Department of Mathematics, Hamedan Branch, Islamic Azad University, Hamedan, Iran.

[§]Department of Mathematics, Central Tehran Branch, Islamic Azad University, Tehran, Iran.

[¶]Department of Computer Sciences, University of Calgary, Canada.

gence order 2^n [5]. According to this conjecture, there have been many efforts to construct such optimal methods without memory. Certain authors used the Newtons interpolation polynomial to construct the self-accelerating parameter and gave some effective iterative methods with memory [4, 12, 15]. The acceleration in convergence is based on the utilizing of a variation of at least one free non-zero parameter in each iterative step. This parameter is calculated using information from the current and previous iterations. Despite many successful and genuine attempts in developing methods with memory, a motivation for studying adaptive method with memory is needed. Previous scientists just use the information of the current and the previous iterations while it is possible to reuse all the information of the previous iterations to increase the efficiency as high as possible. In this paper, two adaptive methods have been introduced based on the Cordero-Torregrosa method with high-efficiency index.

This paper is organized as follows: In Section 2, based on the Cordero-Torregrosa method [1], an optimal fourth-order method is proposed for solving nonlinear equations, then two iterative methods with memory are constructed two adaptive methods. Using the new technique, the new iterative methods with memory reach the desired convergence order. Numerical examples are given in Section 3 to display the implementation of the new method, which confirms the theoretical results. Section 4 is tendered a short conclusion.

2 The method and analysis of convergence

In this section, we consider the following two-step method without memory that is kind of Cordero-Torregrosa method [1],

$$\begin{cases} w_n = x_n + \gamma f(x_n), \\ y_n = x_n - \frac{f(x_n)}{f[x_n, w_n]}, \\ x_{n+1} = y_n - \frac{f(y_n)f[x_n, w_n]}{f[x_n, y_n]f[y_n, w_n]}, \\ n = 0, 1, 2, \dots \end{cases} \quad (2.2)$$

when x_0 is given. This method is optimal with convergence order four that uses three functional

evaluations.

Theorem 2.1. Method (2.2) has convergence order four if x_0 is close enough to the zero. Its error equation is

$$e_{n+1} = (1 + f'(\alpha)\gamma)^2 c_2(2c_2^2 - c_3)e_n^4 + O(e_n^5).$$

Proof. Instead of involving reporting some algebraic expressions, we prefer to present the prove according to implementation in mathematica. The commands are self-explained and simple to understand, so we neglect to decipher them.

$$\begin{aligned} f[e_] &= f1a * \left(e + \sum_{i=2}^4 c_i * e^i \right); \\ ew &= \gamma * f[e] + e; \\ f[x_, x_] &:= f'[x]; \\ f[x_, y_] &:= (f[x] - f[y]) / (x - y); \\ In[1] := ey &= Series \left[e - \frac{f[e]}{f[e, ew]}, \{e, 0, 4\} \right] \\ & // FullSimplify (*First step*) \\ Out[1] &= c_2(\gamma f1a + 1)e^2 \\ & + (c_2^2(-(\gamma f1a + 2) + 2)) \\ & + c_3 + (\gamma f1a + 1)(\gamma f1a + 2)e^3 \\ & + (c_2^3(\gamma f1a + 3) + 5) + 4 \\ & - c_2 c_3 (\gamma f1a (\gamma f1a (2\gamma f1a + 7) + 10) + 7) \\ & + c_4 + (\gamma f1a + 1)(\gamma f1a + 3) + 3 e^4 + O[e]^5 \\ In[2] := ez &= \frac{f[ew]f[ey]}{(f[ew] - f[ey])f[e, ey]} \\ & // FullSimplify (*Second step*) \\ Out[2] &= (1 + \gamma f1a)^2 c_2 (2c_2^2 - c_3) e^4 + O[e]^5. \end{aligned}$$

This completes our proof. □

To develop a method with memory, it is necessary to introduce a parameter that can be approximated as the procedure follows, so we modify Method (2.2) as follows:

$$\begin{cases} w_n = x_n + \gamma f(x_n) \\ y_n = x_n - \frac{f(x_n)}{f[x_n, w_n] + \beta f(w_n)}, \\ x_{n+1} = y_n - \frac{f(y_n)f[x_n, w_n]}{f[x_n, y_n]f[y_n, w_n]}, \\ n = 0, 1, 2, \dots, \end{cases} \quad (2.3)$$

Table 1: Numerical results for $f_1(x)$ by the methods with memory

Methods	$ x_1 - \alpha $	$ x_2 - \alpha $	$ x_3 - \alpha $	r_c	EI
Method (3.27) with (3.28)	0.9027(-3)	0.1827(-10)	0.7583(-41)	3.9489	1.5806
Method (3.27) with (3.29)	0.9027(-3)	0.1787(-10)	0.6924(-41)	4.1477	1.6067
Method (3.27) with (3.30)	0.9027(-3)	0.4948(-13)	0.4048(-51)	3.9116	1.5756
Method (3.27) with (3.31)	0.9027(-3)	0.3226(-10)	0.1607(-85)	3.9783	1.5845
Method (3.32)	0.2415	0.6840(-2)	0.7002(-3)	6.4360	1.8601
Method (2.4)	0.1501(-2)	0.3601(-19)	0.2462(-129)	7.0001	1.9129
Method (2.15)	0.1501(-2)	0.3781(-19)	0.7333(-129)	7.5234	1.9594

Table 2: Numerical results for $f_2(x)$ by the methods with memory

Methods	$ x_1 - \alpha $	$ x_2 - \alpha $	$ x_3 - \alpha $	r_c	EI
Method (3.27) with (3.28)	0.1284(-1)	0.6392(-6)	0.2098(-22)	3.9306	1.5781
Method (3.27) with (3.29)	0.1288(-6)	0.6157(-6)	0.1592(-22)	4.1242	1.6036
Method (3.27) with (3.30)	0.1284(-1)	0.3521(-7)	0.1663(-27)	4.2464	1.6193
Method (3.27) with (3.31)	0.1616(-1)	0.1044(-7)	0.4112(-37)	4.4466	1.6444
Method (3.32)	0.2415	0.6840(-2)	0.7002(-3)	6.3360	1.8504
Method (2.4)	0.1181(-1)	0.8712(-64)	0.3771(-447)	7.3223	1.9418
Method (2.15)	0.1182(-1)	0.5791(-10)	0.1491(-63)	7.7432	1.9783

By adding the parameter β in the method (2.2) remains optimal with following error equation.

Theorem 2.2. The error equation of method (2.3) is given by

$$e_{n+1} = (1 + f'(\alpha)\gamma)^2(\beta + c_2) (c_2(\beta + 2c_2) - c_3)e_n^4 + O(e_n^5).$$

To derive a method with memory, we suppose that $\gamma = \frac{1}{f'(\alpha)} \approx \frac{-1}{N_3'(x_k)}$ and $\beta = -c_2 = -\frac{f''(\alpha)}{2f'(\alpha)} \approx -\frac{N_4''(w_k)}{2N_4'(w_k)}$, where $N_3(x_k)$ and $N_4(w_k)$ are Newton's interpolating polynomials of two and third degree on the nodes $(x_k, y_{k-1}, x_{k-1}, w_{k-1})$ and $(x_k, w_k, y_{k-1}, x_{k-1}, w_{k-1})$, respectively.

Here we affirmative that albeit this method is with memory but it is not adaptive yet. It consumes information on the last two iterations. To develop this method to an adaptive method with memory we update γ_n and β_n based on all the available information from the first iteration to the current iteration. So in each iteration, we use

the following accelerators:

$$\begin{cases} \gamma_n = \frac{-1}{N_{3n}'(x_n)}, \\ \beta_n = -\frac{N_{3n+1}''(w_n)}{2N_{3n+1}'(w_n)}, \\ w_n = x_n + \gamma_n f(x_n), \\ y_n = x_n - \frac{f(x_n)}{f[e_n, w_n] + \beta_n f(w_n)}, \\ x_{n+1} = y_n - \frac{f(y_n)f[x_n, w_n]}{f[x_n, y_n]f[y_n, w_n]}, \\ n = 0, 1, 2, \dots, \end{cases} \quad (2.4)$$

where x_0 and γ_0 are given suitably.

Lemma 2.1. If $\gamma_n = \frac{1}{N_{3n}'(x_n)}$, and

$$\beta_n = \frac{-N_{3n+1}''(w_n)}{N_{3n+1}'(w_n)}, \text{ then}$$

$$1 + \gamma_n f'(\alpha) \sim \prod_{k=0}^{n-1} e_k e_{k,y} e_{k,w},$$

$$c_2 + \beta_n \sim \prod_{k=0}^{n-1} e_k e_{k,y} e_{k,w}, \quad n = 1, 2, \dots$$

where

$$e_k = x_k - \alpha, \quad e_{k,y} = y_k - \alpha, \quad e_{k,w} = w_k - \alpha.$$

Proof. The proof is very similar to Lemma 4, [14] and Lemma 1, [19]. \square

The convergence order of the adaptive Method (2.4) is given in the following theorem:

Theorem 2.3. *If $\gamma_n = \frac{1}{N'_{3n}(x_n)}$ and, $\beta_n = \frac{-N''_{3n+1}(w_n)}{N'_{3n+1}(w_n)}$ then convergence order of method (2.3) is obtained from solution of the following nonlinear system of equations:*

$$\begin{cases} r^{n+1} - 3(1 + q + p)(1 + r + \dots + r^{n-1}) \\ -4r^n = 0, \\ r^n q - 2(1 + q + p)(1 + r + \dots + r^{n-1}) \\ -2r^n = 0, \\ r^n p - (1 + q + p)(1 + r + \dots + r^{n-1}) \\ -r^n = 0, \quad n = 1, 2, \dots \end{cases} \quad (2.5)$$

Proof. Let x_n, y_n and w_n have convergent orders r, p and q respectively, i.e.,

$$e_{n+1} \sim e_n^r \sim e_{n-1}^{r^2} \sim \dots \sim e_0^{r^{n+1}}, \quad (2.6)$$

$$e_{n,y} \sim e_n^q \sim e_{n-1}^{rq} \sim \dots \sim e_0^{r^n q}. \quad (2.7)$$

$$e_{n,w} \sim e_n^p \sim e_{n-1}^{rp} \sim \dots \sim e_0^{r^n p}. \quad (2.8)$$

Also, considering Lemma (2.1), we have

$$\begin{aligned} 1 + \gamma_n f'(\alpha) &\sim \prod_{k=0}^{n-1} e_k e_{k,y} e_{k,w} \\ &= (e_0 e_{0,y} e_{0,w})(e_1 e_{1,y} e_{1,w}) \dots (e_{n-1} e_{n-1,y} e_{n-1,w}) \\ &= (e_0 e_0^q e_0^p)(e_0^r e_0^r q e_0^r p) \dots (e_0^{r(n-1)} e_0^{rq(n-1)} e_0^{rp(n-1)}) \\ &= e_0^{(1+q+p)(1+r+r^2+\dots+r^{n-1})}. \end{aligned} \quad (2.9)$$

Therefore,

$$\begin{aligned} e_{n,w} &\sim (1 + \gamma_n f'(\alpha)) e_n \\ &= e_0^{(1+q+p)(1+r+r^2+\dots+r^{n-1})} e_0^{rn}, \end{aligned} \quad (2.10)$$

$$\begin{aligned} e_{n,y} &\sim (1 + \gamma_n f'(\alpha))(c_2 + \beta) e_n^2 \\ &= e_0^{2(1+q+p)(1+r+r^2+\dots+r^{n-1})} e_0^{2rn}, \end{aligned} \quad (2.11)$$

and

$$\begin{aligned} e_{n+1} &\sim (1 + \gamma_n f'(\alpha))^2 (c_2 + \beta) \\ &\quad (c_2(\beta + 2c_2) - c_3) e_n^4 \\ &\sim e_0^{3(1+q+p)(1+r+r^2+\dots+r^{n-1})} e_0^{4pn}. \end{aligned} \quad (2.12)$$

Combining the right-hand-side (2.6)-(2.12), (2.7)-(2.11) and (2.8)-(2.10) we have the following system of nonlinear equations:

$$\begin{cases} r^{n+1} - 3(1 + q + p)(1 + r + \dots + r^{n-1}) \\ -4r^n = 0, \\ r^n q - 2(1 + q + p)(1 + r + \dots + r^{n-1}) \\ -2r^n = 0, \\ r^n p - (1 + q + p)(1 + r + \dots + r^{n-1}) \\ -r^n = 0 \end{cases} \quad (2.13)$$

This completes the proof. \square

Remark 2.1. *We have solved the nonlinear system of equations (2.13) by varying n . If $n = 1$, we use the information from the last iteration and the convergence order is 7.0000, and for $n = 2$ the convergence order is 7.40515, while if $n = 3$ the convergence order is 7.45635, we use the information from the last two iterations, and so on.*

Now we consider the following two-step method without memory by three parameters

$$\begin{cases} w_n = x_n + \gamma f(x_n), \\ y_n = x_n - \frac{f(x_n)}{f[x_n, w_n] + \beta f(w_n)}, \\ x_{n+1} = y_n - \frac{f(y_n) f[x_n, w_n]}{b}, \\ n = 0, 1, 2, \dots, \end{cases} \quad (2.14)$$

where

$$b = f[x_n, y_n] f[y_n, w_n] + \lambda(y_n - x_n)(y_n - w_n)$$

This method is optimal with convergence order 4.

Theorem 2.4. *The error equation of method (2.14) is given by*

$$e_{n+1} = \frac{d}{f'(\alpha)^2} + o(e_n^5).$$

where

$$d = (1 + f'(\alpha)\gamma)^2 (\beta + c_2) (\lambda + f'(\alpha)^2 (c_2(\beta + 2c_2) - c_3)) e_n^4$$

Table 3: Numerical results for $f_3(x)$ by the methods with memory

Methods	$ x_1 - \alpha $	$ x_2 - \alpha $	$ x_3 - \alpha $	r_c	EI
Method (3.27) with (3.28)	0.1973(-1)	0.4185(-1)	0.7094(-3)	2.6296	1.3802
Method (3.27) with (3.29)	0.1973(-1)	0.2466(-1)	0.6695(-4)	2.8418	1.4164
Method (3.27) with (3.30)	0.1973(-1)	0.3578(-1)	0.5356(-5)	3.6013	1.5328
Method (3.27) with (3.31)	0.1973(-1)	0.7848(-1)	0.5077(-3)	4.3339	1.6303
Method (3.32)	0.1345	0.5325(-7)	0.1349(-51)	5.9941	1.8165
Method (2.4)	0.1921(-3)	0.5121(-22)	0.4932(-152)	7.0404	1.9166
Method (2.15)	0.1941(-3)	0.5593(-22)	0.8862(-168)	7.4334	1.9516

Table 4: Numerical results for $f_4(x)$ by the methods with memory

Methods	$ x_1 - \alpha $	$ x_2 - \alpha $	$ x_3 - \alpha $	r_c	EI
Method (3.27) with (3.28)	0.1814(-1)	0.2164(-2)	0.3004(-9)	3.7760	1.5571
Method (3.27) with (3.29)	0.1814(-1)	0.5435(-2)	0.9601(-8)	4.0711	1.5965
Method (3.27) with (3.30)	0.1814(-1)	0.5592(-2)	0.3220(-8)	4.4546	1.6453
Method (3.27) with (3.31)	0.1814(-1)	0.1762(-1)	0.7697(-7)	5.29261	1.7427
Method (3.32)	0.1991	0.6470(-6)	0.5272(-44)	6.1360	1.8307
Method (2.4)	0.1441(-4)	0.1341(-34)	0.6462(-240)	7.2688	1.9371
Method (2.15)	0.1612(-2)	0.7771(-12)	0.2281(-51)	7.9733	1.9977

To derive an adaptive method with memory, we suppose that $\gamma_n = -\frac{1}{f'(\alpha)} \approx -\frac{1}{N'_{3n}(x_n)}$, $\beta_n = -c_2 = -\frac{f''(\alpha)}{2f'(\alpha)} \approx -\frac{N''_{3n+1}(w_n)}{2N'_{3n+1}(w_n)}$ and $\lambda_n = f'(\alpha)c_3 = \frac{f'''(\alpha)}{6f'(\alpha)} \approx \frac{N'''_{3n+2}(y_n)}{6N'_{3n+2}(y_n)}$, where $N'_{3n}(x_n)$, $N''_{3n+1}(w_n)$, and $N'''_{3n+2}(y_n)$ are Newton's interpolation polynomials around the nodes $\{x_n, x_{n-1}, w_{n-1}, y_{n-1}\}$, $\{w_n, x_n, x_{n-1}, w_{n-1}, y_{n-1}\}$ and $\{y_n, w_n, x_n, x_{n-1}, w_{n-1}, y_{n-1}\}$, respectively.

So in each iteration, we use the following accelerators:

$$\begin{cases} \gamma_n = \frac{-1}{N'_{3n}(x_n)}, \\ \beta_n = -\frac{N''_{3n+1}(w_n)}{2N'_{3n+1}(w_n)}, \\ \lambda_n = \frac{N'''_{3n+2}(y_n)}{6N'_{3n+2}(y_n)}, \\ w_n = x_n + \gamma_n f(x_n), \\ y_n = x_n - \frac{f(x_n)}{f[e_n, w_n] + \beta_n f(w_n)}, \\ x_{n+1} = y_n - \frac{f(y_n)f[x_n, w_n]}{g}. \end{cases} \quad n = 0, 1, 2, \dots \tag{2.15}$$

where

$g = f[x_n, y_n]f[y_n, w_n] + \lambda_n(y_n - x_n)(y_n - w_n)$, x_0 and γ_0 are given suitably.

Lemma 2.2. If $\gamma_n = \frac{-1}{N'_{3n}(x_n)}$, $\beta_n = \frac{-N''_{3n+1}(w_n)}{2N'_{3n+1}(w_n)}$ and $\lambda_n = \frac{N'''_{3n+2}(y_n)}{6N'_{3n+2}(y_n)}$, then

$$1 + \gamma_n f'(\alpha) \sim \prod_{k=0}^{n-1} e_k e_{k,y} e_{k,w},$$

$$c_2 + \beta_n \sim \prod_{k=0}^{n-1} e_k e_{k,y} e_{k,w},$$

$$(\lambda_n + f'(\alpha)^2(c_2(\beta_n + 2c_2) - c_3) \sim \prod_{k=0}^{n-1} e_k e_{k,y} e_{k,w}.$$

where

$$e_k = x_k - \alpha, \quad e_{k,y} = y_k - \alpha, \quad e_{k,w} = w_k - \alpha.$$

Proof. The proof is very similar to Lemma 4, [14] and Lemma 1, [19]. \square

The convergence order of the adaptive method (2.15) is given in the following theorem:

Theorem 2.5. If $\gamma_n = \frac{-1}{N'_{3n}(x_n)}$,
 $\beta_n = \frac{-N''_{3n+1}(w_n)}{2N'_{3n+1}(w_n)}$ and $\lambda_n = \frac{N'''_{3n+2}(y_n)}{6N'_{3n+2}(y_n)}$,

then convergence order of method (2.15) is obtained from solution of the following nonlinear system of equations:

$$\begin{cases} r^{n+1} - 4(1+q+p)(1+r+\dots+r^{n-1}) \\ -4r^n = 0, \\ r^n q - 2(1+q+p)(1+r+\dots+r^{n-1}) \\ -2r^n = 0, \\ r^n p - (1+q+p)(1+r+\dots+r^{n-1}) \\ -r^n = 0, \quad n = 1, 2, \dots \end{cases} \quad (2.16)$$

Proof. Let x_n, y_n and w_n have convergent orders r, p and q respectively, i.e.,

$$e_{n+1} \sim e_n^r \sim e_{n-1}^{r^2} \sim \dots \sim e_0^{r^{n+1}}. \quad (2.17)$$

$$e_{n,y} \sim e_n^q \sim e_{n-1}^{rq} \sim \dots \sim e_0^{r^n q}. \quad (2.18)$$

$$e_{n,w} \sim e_n^p \sim e_{n-1}^{rp} \sim \dots \sim e_0^{r^n p}. \quad (2.19)$$

Also, considering Lemma (2.2), we have

$$\begin{aligned} 1 + \gamma_n f'(\alpha) &\sim \prod_{k=0}^{n-1} e_k e_{k,y} e_{k,w} \\ &= (e_0 e_{0,y} e_{0,w}) (e_1 e_{1,y} e_{1,w}) \dots (e_{n-1} e_{n-1,y} e_{n-1,w}) \\ &= (e_0^q e_0^p) (e_0^r e_0^q e_0^p) \dots (e_0^{r(n-1)} e_0^{rq(n-1)} e_0^{rp(n-1)}) \\ &= e_0^{(1+q+p)(1+r+r^2+\dots+r^{n-1})}, \end{aligned} \quad (2.20)$$

and

$$\begin{aligned} c_2 + \beta &\sim \prod_{k=0}^{n-1} e_k e_{k,y} e_{k,w} \\ &= e_0^{(1+q+p)(1+r+r^2+\dots+r^{n-1})}, \end{aligned} \quad (2.21)$$

and

$$\begin{aligned} \lambda + f'(\alpha)^2 (c_2(\beta + 2c_2) - c_3) &\sim \prod_{k=0}^{n-1} e_k e_{k,y} e_{k,w} \\ &= e_0^{(1+q+p)(1+r+r^2+\dots+r^{n-1})}. \end{aligned} \quad (2.22)$$

Therefore,

$$\begin{aligned} e_{n,w} &\sim (1 + \gamma_n f'(\alpha)) e_n \\ &= e_0^{(1+q+p)(1+r+r^2+\dots+r^{n-1})} e_0^{r^n}, \end{aligned} \quad (2.23)$$

$$\begin{aligned} e_{n,y} &\sim (1 + \gamma_n f'(\alpha)) (c_2 + \beta_n) e_n^2 \\ &= e_0^{2(1+q+p)(1+r+r^2+\dots+r^{n-1})} e_0^{2r^n}, \end{aligned} \quad (2.24)$$

and

$$\begin{aligned} e_{n+1} &\sim (1 + \gamma_n f'(\alpha))^2 (c_2 + \beta_n) \\ &\quad (\lambda_n + f'(\alpha)^2 (c_2(\beta_n + 2c_2) - c_3)) e_n^4 \\ &\sim e_0^{4(1+q+p)(1+r+r^2+\dots+r^{n-1})} e_0^{4p^n}. \end{aligned} \quad (2.25)$$

Incorporate the right-hand-side of the relations (2.17)-(2.25), (2.18)-(2.24) and (2.19)-(2.23), we reach the following system of nonlinear equations

$$\begin{cases} r^{n+1} - 4(1+q+p)(1+r+\dots+r^{n-1}) \\ -4r^n = 0, \\ r^n q - 2(1+q+p)(1+r+\dots+r^{n-1}) \\ -2r^n = 0, \\ r^n p - (1+q+p)(1+r+\dots+r^{n-1}) \\ -r^n = 0 \end{cases} \quad (2.26)$$

This completes the proof. □

Remark 2.2. By solving the nonlinear system of equations (2.26) by varying n . If $n = 1$, we use the information from the last iteration and the convergence order is 7.530 and $n = 2$, we use the information from the last two iterations the convergence order is 7.94449., while if $n = 3$ the convergence order is 7.99315, and so on.

3 Numerical results and comparisons

In this section, the family of methods with memory (2.13) and (2.25) are tested using four examples of nonlinear equations. The errors $|x_k - \alpha|$ of approximations to the sought zeros, are given in Tables where $a(-b)$ shows $a * 10^{-b}$. Computational order of convergence COC , in Tables 1 – 4 computed by follow expression [12]

$$r_c = \frac{\log \left| \frac{f(x_n)}{f(x_{n-1})} \right|}{\log \left| \frac{f(x_{n-1})}{f(x_{n-2})} \right|}.$$

Moreover, the following test functions are used:

$$\begin{aligned}
 f_1(x) &= (x - 2)(x^6 + x^3 + 1)e^{-x^2}, \\
 x_0 &= 1.8, \alpha = 2, \\
 f_2(x) &= e^{(x^2-3x)} \sin(x) + \log(x^2 + 1), \\
 x_0 &= 0.35, \alpha = 0, \\
 f_3(x) &= \sin(t)e^{(t^2+t \cos(t)-1)} + t \log(t + 1), \\
 x_0 &= 0.6, \alpha = 0, \\
 f_4(x) &= e^{(t)} \sin(t) + \log(t^4 - 3t + 1), \\
 x_0 &= 0.4, \alpha = 0.
 \end{aligned}$$

Also, we compare our methods to some of the existing methods given as follows:

Example 3.1. Xiaofeng Wang method method with order $2\sqrt{6} \approx 4.4495$ [16]:

$$\begin{cases}
 w_n = x_n + f(x_n) \\
 z_n = x_n - \frac{f(x_n)}{f[x_n, w_n]}, \\
 y_n = z_n - \frac{\lambda_n(z_n - x_n)^2}{1 + \lambda_n(z_n - x_n)}, \\
 x_{n+1} = y_n - \frac{f(y_n)f[x_n, w_n]}{f[x_n, y_n]f[y_n, w_n]},
 \end{cases} \quad (3.27)$$

where

$$\lambda_n = \frac{z_{n-1} - z_n}{(z_{n-1} - x_{n-1})^2}. \quad (3.28)$$

$$\lambda_n = \frac{z_{n-1} - x_n}{(z_n - x_{n-1})(x_n - x_{n-1})}. \quad (3.29)$$

$$\begin{aligned}
 \lambda_n &= \frac{(z_{n-1} - x_n)(y_{n-1} - x_{n-1})}{(x_n - x_{n-1})^3} \\
 &\times \left\{ \frac{2y_{n-1} - x_n - w_{n-1}}{y_{n-1} - w_{n-1}} \right\} \\
 &+ \frac{2z_n - x_n - w_n}{(y_{n-1} - x_n)(z_{n-1} - x_{n-1})} \\
 &+ \frac{(z_n - z_{n-1})^2(w_{n-1} - x_{n-1})}{(z_{n-1} - x_{n-1})^2(z_n - w_{n-1})(w_{n-1} - z_{n-1})}, \quad (3.30)
 \end{aligned}$$

$$\begin{aligned}
 \lambda_n &= \frac{(x_n - z_{n-1})(2x_n - y_{n-1} - z_{n-1})}{(z_n - x_{n-1})(z_{n-1} - x_{n-1})^2} \\
 &- \frac{2z_n - x_n - w_n}{(x_n - y_{n-1})(z_{n-1} - x_{n-1})} \\
 &+ \frac{(w_{n-1} - y_{n-1})(x_n - z_{n-1})}{(x_n - x_{n-1})(z_{n-1} - w_{n-1})(z_n - x_{n-1})} \\
 &- \frac{(z_n - z_{n-1})^2(w_{n-1} - x_{n-1})^2}{(z_{n-1} - w_{n-1})^2(z_{n-1} - x_{n-1})^3}. \quad (3.31)
 \end{aligned}$$

Example 3.2. Alicia Cordero et al. method method with order 7 [2]:

$$\begin{cases}
 x_0, \gamma_0, \lambda_0 & w_0 = x_0 + \gamma_0 f(x_0), \\
 \gamma_k = -\frac{1}{N_3'(x_k)}, & \lambda_k = -\frac{N_4''(w_k)}{2N_4'(w_k)}, \\
 w_k = x_k + \gamma_k f(x_k), & k \geq 1, \\
 y_k = x_k - \frac{f(x_k)}{f[x_k, w_k] + \lambda_k f(w_k)}, \\
 x_{k+1} = y_k - \frac{f(y_k)}{f[y_k, x_k] + f[y_k, x_k, w_k](y_k - x_k)}.
 \end{cases} \quad (3.32)$$

4 Conclusion

In this study, we introduced two adaptive methods with memory based on the variant of the Cordero-Torregrosa method. The technique of adaptive method uses all the available information from the first to current iterations. Consequently, it increases the efficiency index as high as possible. The new derivative free methods have possessed the swift convergence order 7.46315 and 7.99315, and they only use the information from the last two iterations. Observing the tables and examples, we found that the proposed new adaptive methods have higher convergence compared to the Wang method [16] and other methods. For the future work, similar to [6, 10], local and semi local analysis of the proposed methods can be carried out.


Acknowledgements

This work has been partially supported by Islamic Azad University-Hamedan Branch.

References

- [1] A. Cordero, J. L. Hueso, E. Martnez, J. R. Torregrosa, A new technique to obtain derivative-free optimal iterative methods for solving nonlinear equations, *J. Comp. Appl. Math.* 252 (2013) 95-102.
- [2] A. Cordero, T. Lotfi, P. Bakhtiari, J.R. Torregrosa, An efficient two-parametric family

- with memory for nonlinear equations, *Numer. Algor.* 68 (2014) 323-335.
- [3] A. M. Ostrowski, Solution of equations and systems of equations, Academic press, *New York*, (1964).
- [4] F. Soleymani, T. Lotfi, E. Tavakoli, F. Khaksar Haghani, Several iterative methods with memory using self-accelerators, *Appl. Math. Comp.* 254 (2015) 452-458.
- [5] H. T. Kung, J. F. Traub, Optimal order of one-point and multipoint iteration, *J. Assoc. Comp. Mach.* 21 (1974) 643-651.
- [6] H. Veisheh, T. Lotfi, T. Allahviranloo, A study on the local convergence and dynamics of the two-step and derivative-free Kung-Traub's method, *J. Comput. Appl. Math.* 37 (2018) 2428-2444.
- [7] J. F. Traub, Iterative Methods for the Solution of Equations, Prentice Hall, *New York*, (1964).
- [8] M. A. Fariborzi Araghi, T. Lotfi, V. Torkashvand, A general efficient family of adaptive method with memory for solving nonlinear equations, *Bull. Math. Soc. Sci. Math. Roumanie* 11 (2019) 37-49.
- [9] M. Mohamadi Zadeh, T. Lotfi, M. Amirfakhrian, Developing two efficient adaptive Newton-type methods with memory, *Math Meth Appl Sci.* 17 (2019) 5687-5695.
- [10] M. Moccari, T. Lotfi, On a two-step optimal Steffensen-type method: Relaxed local and semi-local convergence analysis and dynamical stability, *J. Math. Anal. Appl.* 468 (2018) 240-269.
- [11] M. S. Petkovic, B. Neta, L. D. Petkovic, J. Dzunic, Multipoint methods for solving nonlinear equations: a survey, *Appl. Math. Comp.* 226 (2014) 635-660.
- [12] M. S. Petkovi, S. Ili, J. Duni, Derivative free two-point methods with and without memory for solving nonlinear equations, *Appl. Math. Comp.* 217 (2010) 1887-1895.
- [13] T. Lotfi, P. Bakhtiari, A. Cordero, K. Mahdiani, J. R. Torregrosa, Some new efficient multipoint iterative methods for solving nonlinear systems of equations, *J. Comp. Math.* 92 (2015) 1921-1934.
- [14] T. Lotfi, F. Soleymani, Z. Noori, A. Kman, F. Khaksar Haghani, Efficient Iterative Methods with and without Memory Possessing High Efficiency Indices, *Discrete Dynamics in Nature and Society*, Article ID 912796, (2014) 9 pages.
- [15] V. Torkashvand, T. Lotfi, M. A. F. Araghi, A new family of adaptive methods with memory for solving nonlinear equations, *Math. Sci.* 18 (2019) 1-20.
- [16] X. Wang, A new accelerating technique applied to a variant of Cordero-Torregrosa method, *J. Comp. Appl. Math.* 330 (2018) 695-709.
- [17] X. Wang, T. Zhang, A new family of Newton-type iterative methods with and without memory for solving nonlinear equations, *Calcolo*, 51 (2014) 1-15.
- [18] X. Wang, An Ostrowski-type method with memory using a novel self-accelerating parameter, *J. Comp. Appl. Math.* 330 (2018) 710-720.
- [19] X. Wang, T. Zhang, High-order Newton-type iterative methods with memory for solving nonlinear equations, *Math. Commun.* 19 (2014) 91-109.



Maryam Mohamadi Zade got her PhD degree in applied mathematics from the department of applied mathematics, Hamedan Branch, in 2019. Her research interest in numerical analysis includes approximating numerical solutions using

iterative methods for nonlinear systems of equations.



Taher Lotfi is an associate professor in applied mathematics and numerical analysis. His research interest includes iterative methods for approximating numerical solution of nonlinear systems of equations.



Majid Amirfakhrian is full professorship of Applied Mathematics at Azad University, Central Tehran Branch (IAUCTB). He is currently a Visiting Researcher at the University of Calgary, Canada. His main research interests lie in Numerical Analysis and Approximation. Some of his specific interests are solving the problems of Fuzzy Approximation, Data Sciences, Multivariate Approximation, Image Processing and Numerical Partial Differential Equations. He uses approximation techniques in some applied problems, but at the same time, he works on the theoretical aspects of the mathematical problems. He has side interests in Optimization and Statistics. He uses computational methods in his research.